

Advanced Mathematics

for Rwanda Schools

Teacher's Guide

Senior Six

Authors

Emmanuel Ngezahayo

Pacifique Icyingeneye

FOUNTAIN PUBLISHERS
www.fountainpublishers.co.ug

Fountain Publishers Rwanda Ltd

P.O. Box 6567

Kigali, Rwanda

E-mail: fountainpublishers.rwanda@gmail.com

sales@fountainpublishers.co.ug;

publishing@fountainpublishers.co.ug

Website: www.fountainpublishers.co.ug

© Emmanuel Ngezahayo, Pacifique Icyingeneye 2017

First Published 2017

All rights reserved; no part of this book may be reproduced, stored in a retrieval system or transmitted in any form or by any means; electronic, mechanical, photocopying, recording or otherwise without prior written permission from the authors and the publishers.

ISBN: 978-9970-19-420-9

Contents

Section 1:	General introduction	ix
Section 2:	Content map	xvii
Section 3:	Sample lesson plan	xxii
Section 4:	Units description	xxviii

Unit 1 Complex numbers..... 1

Lesson 1.1.	Concepts of complex numbers	3
Lesson 1.2.	Definition and properties of the number i	5
Lesson 1.3.	Geometric representation of a complex number	7
Lesson 1.4.	Modulus of a complex number.....	8
Lesson 1.5.	Loci related to distances on Argand diagram	11
Lesson 1.6.	Equality of two complex numbers.....	13
Lesson 1.7.	Addition and subtraction of complex numbers.....	14
Lesson 1.8.	Conjugate and opposite of a complex number	16
Lesson 1.9.	Multiplication of complex numbers.....	17
Lesson 1.10.	Inverse and division of complex numbers.....	19
Lesson 1.11.	Square root of a complex number	20
Lesson 1.12.	Linear equations	22
Lesson 1.13.	Quadratic equations	24
Lesson 1.14.	Polynomials in set of complex numbers.....	25
Lesson 1.15.	Argument of a complex number	29
Lesson 1.16.	Loci related to the angles	32
Lesson 1.17.	Polar form of a complex number	36
Lesson 1.18.	Multiplication and division of complex numbers in polar form	38
Lesson 1.19.	Powers of complex number in polar form ..	41
Lesson 1.20.	n th root of a complex number.....	43
Lesson 1.21.	Graphical representation of n th roots of a complex number	45
Lesson 1.22.	Construction of regular polygon.....	49
Lesson 1.23.	Exponential form of a complex number	52
Lesson 1.24.	Trigonometric number of a multiple of an angle	53
Lesson 1.25.	Linearisation of trigonometric expressions	55
Lesson 1.26.	Solving equation of the form $a \cos x + b \sin x = c$, $a, b, c \in \mathbb{R}$ and $a \cdot b \neq 0$.	57
Lesson 1.27.	Alternating current problem.....	59

Unit 2 Logarithmic and Exponential Functions75

Lesson 2.1.	Domain and range of natural logarithmic functions	77
Lesson 2.2.	Limit and asymptotes for natural logarithmic functions	79
Lesson 2.3.	Derivative of natural logarithmic functions.	81
Lesson 2.4.	Variation and curve sketching of natural logarithmic functions	83
Lesson 2.5.	Domain and range of logarithmic function with any base.....	88
Lesson 2.6.	Limit of logarithmic function with any base	91
Lesson 2.7.	Logarithmic differentiation.....	94
Lesson 2.8.	Further logarithmic differentiation	95
Lesson 2.9.	Variation and curves of logarithmic functions with any base	97
Lesson 2.10.	Domain and range of exponential functions with base " e "	104
Lesson 2.11.	Limit of exponential functions with base " e "	105
Lesson 2.12.	Derivative of exponential functions with base " e ".....	108
Lesson 2.13.	Variation and curve of exponential functions with base " e "	109
Lesson 2.14.	Domain and range of exponential functions with any base	113
Lesson 2.15.	Limit of exponential functions with any base.....	114
Lesson 2.16.	Derivative of exponential functions with any base.....	116
Lesson 2.17.	Variation and curve of exponential functions with any base	117
Lesson 2.18.	Compound interest problems.....	122
Lesson 2.19.	Mortgage amount problems.....	124
Lesson 2.20.	Population growth problems	126
Lesson 2.21.	Depreciation value problems	127
Lesson 2.22.	Earthquake problems	128
Lesson 2.23.	Carbon-14 dating problems.....	130

Unit 3	Taylor and Maclaurin's Expansions	141
Lesson 3.1.	Finite series.....	143
Lesson 3.2.	Infinite series	144
Lesson 3.3.	Tests for convergence of series	147
Lesson 3.4.	Power series	149
Lesson 3.5.	Taylor and Maclaurin series	151
Lesson 3.6.	Taylor series by using Maclaurin series ..	153
Lesson 3.7.	Calculation of limits	156
Lesson 3.8.	Estimation of the number e	157
Lesson 3.9.	Estimation of the number π	158
Lesson 3.10.	Estimation of trigonometric number of an angle.....	161
Lesson 3.11.	Estimation of an irrational number	162
Lesson 3.12.	Estimation of natural logarithm of a number.....	164
Lesson 3.13.	Estimation of roots of equations.....	166
Unit 4	Integration.....	175
Lesson 4.1.	Differentials.....	177
Lesson 4.2.	Definition of indefinite integrals.....	179
Lesson 4.3.	Properties of integrals	180
Lesson 4.4.	Integration by substitution.....	183
Lesson 4.5.	Integration of rational functions where numerator is expressed in terms of derivative of denominator.....	185
Lesson 4.6.	Integration of rational functions where degree of numerator is greater or equal to the degree of denominator.....	187
Lesson 4.7.	Integration of rational functions where denominator is factorised into linear factors	188
Lesson 4.8.	Integration of rational functions where denominator is a quadratic factor.....	192
Lesson 4.9.	Integrals of the form $\int \sin mx \cos nx \, dx, \int \cos mx \cos nx \, dx, \int \sin mx \sin nx \, dx$.	195
Lesson 4.10.	Integrals of the form $\int \sin^m x \cos^n x \, dx$	197
Lesson 4.11.	Integrals of the form $\int \tan^m x \sec^n x \, dx$	199
Lesson 4.12.	Integrals containing $\sin x, \cos x, \tan x$ on denominator	201
Lesson 4.13.	Integrals containing $\sin^2 x, \cos^2 x$ on denominator.....	203
Lesson 4.14.	Integrals containing $\sqrt[n]{ax+b}$	204
Lesson 4.15.	Integrals containing $\sqrt{ax^2+bx+c}$	205

Lesson 4.16. Integration by parts 209
 Lesson 4.17. Integration by reduction formulae 210
 Lesson 4.18. Integration by Maclaurin series 212
 Lesson 4.19. Definite integrals 213
 Lesson 4.20. Properties of definite integrals..... 215
 Lesson 4.21. Improper integrals, Infinite limits of
 integration..... 217
 Lesson 4.22. Discontinuous integrand 219
 Lesson 4.23. Calculation of area of plane surfaces 221
 Lesson 4.24. Calculation of volume of a solid of
 revolution..... 225
 Lesson 4.25. Calculation of arc length of a curved
 surface 229

Unit 5 Differential Equations 245

Lesson 5.1. Definition and classification..... 247
 Lesson 5.2. Differential equations with separable
 variables 249
 Lesson 5.3. Simple homogeneous equations..... 250
 Lesson 5.4. Linear differential equations 252
 Lesson 5.5. Particular solution 254
 Lesson 5.6. Second order differential equations:
 Definition 256
 Lesson 5.7. Second order differential equations
 with constant coefficient: two distinct
 real roots 257
 Lesson 5.8. Characteristic equation with a
 double root..... 259
 Lesson 5.9. Characteristic equation with
 complex roots 261
 Lesson 5.10. Non- homogeneous linear differential
 equations of the second order with
 constant coefficients..... 262
 Lesson 5.11. Differential equations of the second order
 with the right hand side $r(x) = Pe^{ax}$ 265
 Lesson 5.12. Differential equations of the second
 order with the right hand side
 $r(x) = Pe^{ax} \cos \beta x + Qe^{ax} \sin \beta x$ 267
 Lesson 5.13. Applications: Newton's law of cooling..... 269
 Lesson 5.14. Applications: Electrical circuits 271

Unit 6	Intersection and Sum of Subspaces.....	281
	Lesson 6.1. Definition of subspaces.....	282
	Lesson 6.2. Intersection of subspaces.....	283
	Lesson 6.3. Dimensions of intersection of subspaces .	285
	Lesson 6.4. Sum of subspaces.....	286
	Lesson 6.5. Dimension of sum of subspaces.....	288
	Lesson 6.6. Grassmann's formula of dimension for subspaces.....	289
Unit 7	Transformation of Matrices.....	295
	Lesson 7.1. Kernel and range.....	297
	Lesson 7.2. Elementary row/column operations.....	298
	Lesson 7.3. Eigenvalues and eigenvectors.....	300
	Lesson 7.4. Diagonalisation of a matrix.....	304
	Lesson 7.5. Echelon matrix.....	307
	Lesson 7.6. Matrix inverse.....	309
	Lesson 7.7. Rank of matrix.....	312
	Lesson 7.8. Solving system of linear equations.....	313
	Lesson 7.9. Power of matrix.....	315
Unit 8	Conics.....	325
	Lesson 8.1. Generalities on conic sections.....	326
	Lesson 8.2. Definition and equation of a parabola.....	329
	Lesson 8.3. Tangent line and normal line on a parabola.....	334
	Lesson 8.4. Definition and equation of an ellipse.....	337
	Lesson 8.5. Tangent line and normal line on ellipse...	342
	Lesson 8.6. Definition and equation of a hyperbola ...	344
	Lesson 8.7. Tangent line and normal line on hyperbola.....	349
	Lesson 8.8. Definition of polar coordinates.....	351
	Lesson 8.9. Polar equation of a conic.....	353
	Lesson 8.10. Polar equation of a straight line.....	355
	Lesson 8.11. Polar form of a circle.....	357
	Lesson 8.12. Applications of conics.....	359
Unit 9	Random Variables.....	375
	Lesson 9.1. Probability density function of a discrete random variable.....	376
	Lesson 9.2. Cumulative distribution of discrete random variable.....	378
	Lesson 9.3. Expected value, variance and standard deviation of a discrete random variable...	380
	Lesson 9.4. Binomial distribution.....	382

Lesson 9.5.	Expected value, variance and standard deviation of a binomial distribution	384
Lesson 9.6.	Poisson distribution	385
Lesson 9.7.	Probability density function of a continuous random variable	388
Lesson 9.8.	Cumulative distribution of continuous random variable.....	391
Lesson 9.9.	Variance and standard deviation of a continuous random variable	392
<i>Answers for Summative Evaluation One.....</i>		<i>403</i>
<i>Answers for Summative Evaluation Two.....</i>		<i>416</i>
<i>Answers for Summative Evaluation Three.....</i>		<i>425</i>
<i>Answers for Summative Evaluation Four.....</i>		<i>438</i>
References.....		450

Section 1: General introduction

1.1. Instruction to the user

This Teacher's Guide is developed to help the teacher and learners to use the Advanced Mathematics Learner's Book Senior 6. Generally, this Teacher's Guide provides:

- ✔ Content map.
- ✔ Sample lesson plan.
- ✔ All lessons to be taught for each unit, recommended periods to each lesson and development of each lesson.
- ✔ Generic competence and cross-cutting issues to be addressed for each lesson.
- ✔ Summary for each lesson.
- ✔ Guidance on how to do each activity.
- ✔ Solutions to all activities.
- ✔ Answers (final answers) to all exercises.

Notice

- ✔ Number of lessons given is a proposal, the teacher can combine two or more lessons depending on students' level of understanding or number of periods a class has per day.
- ✔ The teacher must do exercises or activities instead of copying answers given in this book.
- ✔ This book is a guide, the teacher must prepare and use her/his innovation based on the guidance provided herein.

1.2. General introduction to the new curriculum

The curriculum for Rwandan schools at primary and secondary levels has been changed from knowledge and content based to competence based. The overall objective is to raise the standards of education through providing all the knowledge, skills, attitudes and values that constitute the competencies that are relevant to real life, enabling them to succeed in the world of work and of further learning.

1.3. Competences to be developed

Competence is defined as ability to use appropriate combination of knowledge, skills, attitudes and values in order to accomplish a particular task successfully. That is, the ability to apply learning with confidence in a range of situations. There are basic competences and generic competences

Basic competences

These are essential competences highlighted in national policy documents. Their descriptors give an orientation about priority subjects to be taught, and the kind of learner envisaged at the end of every cycle. These are:

- ✔ Literacy
- ✔ Numeracy
- ✔ ICT and digital
- ✔ Citizenship and nationality identity
- ✔ Entrepreneurship and business development
- ✔ Science and technology

Generic competence

Generic competences apply across the curriculum and can be developed in all subjects. They are transferrable and applicable to a range of situations including employment. These are;

- ✔ **Critical thinking:** Helps learners become capable of critical and open-minded questioning and reasoning.
- ✔ **Communication:** Through group discussion, learners develop their communication skills. Also through presentation, learners communicate and convey information and ideas through speaking when they are presenting their work.
- ✔ **Self confidence:** Learners will gain self confidence competence when they are presenting their work.
- ✔ **Cooperation, interpersonal management and life skills:** Through group discussion, learners will learn how to

work with others, participate and collaborate. It gives them the opportunity to debate their opinions, take turns, and work together towards a common goal.

- ✔ **Research:** Learners will gain the knowledge of using the internet. They will also learn how to do a research.

1.4. Techniques to develop competences

The following are some techniques that will be used to develop competences:

- ✔ **Group work:** Learners form groups of at least 5 learners. They discuss something in groups and report back on what the group discussed. In this way, they learn from each other, and how to work together as a group to address or solve a problem.
- ✔ **Pairing:** Learners work in pairs (two learners). They exchange ideas and write down results.
- ✔ **Practical work:** Learners form groups and do the activity practically such as curve sketching, drawing figures and then present their results.
- ✔ **Research work:** Learners form groups and do research either by reading textbooks or using the internet.

1.5. Cross cutting issues to be addressed

Cross cutting issues are the link between what is taught in school and real life. The following are integrated in Mathematics subject:

- ✔ **Peace and values education:** By offering learners the chance of working in groups, they develop peace and value education by the fact of being convinced without fighting.
- ✔ **Inclusive education:** The lesson covers all categories of learners.
- ✔ **Gender education:** Both girls and boys have equal opportunity to study mathematics.
- ✔ **Financial education:** Learners will be able to use mathematics to solve problems in finance.

1.6. Equipments needed for the subject

Learners will need geometric instruments for sketching curves and scientific calculators for some calculations.

1.7. General guidance on activities and exercises

In Core mathematics-Learners Book Six, there are nine units. There are many activities to be done by learners before a new lesson. Some activities will be done in pairs (exactly two learners) and others in groups (more than two learners; at least five learners). This will help learners to understand well the lesson. For each activity, there is an icon to show the teacher what kind the activity is.

At the end of the lesson, there is a series of exercises which summarise the lesson taught. As a teacher, let learners do those exercises and correct them on the chalkboard. Also at the end of each unit, there is a series of exercises which summarise the whole unit.

Icons

The icons used in this book are as follows:



Practical Activity icon

The hand indicates a practical activity such as curve sketching, draw figures and then presents the results or comments. The activity is done in groups.



Group Work icon

Group work means that learners are expected to discuss something in groups and report back on what their group discussed. In this way, they learn from each other and also learn how to work together as a group to address or solve a problem.



Pairing Activity icon

This means that they are required to do the activity in pairs, exchange ideas and write down the results.



Research Activity icon

Some activities require you to do research either by reading textbooks or using the internet.

When organising groups or pairs:

- ✔ Kind of activity: Depending on the icon shown for the activity.
- ✔ If the activity is a pairing activity, request learners to form groups of exactly two learners each.
- ✔ If the activity is a group work, practical or research activity, request learners to form groups of at least five learners each.
- ✔ For each lesson, we provided the topics (prerequisites) that learners have to recall before studying the lesson or doing the activity. So, as teacher help learners, especially time takers, to recall those topics.
- ✔ Decide who will be working together: Learners who are sitting together should sit face to face.
- ✔ Give the learners roles; manager/leader, resource collector, reporter, ...
- ✔ Arrange a stop signal: Decide on a signal that tells learners when you want them to stop talking and listen to the teacher such as clapping rhythm/1, 2, 3 look at me/ shaker/ 5, 4, 3, 2, 1/ hands up/ mobile ringtone.
- ✔ Make a noise monitor to show learners what level of noise is OK.
- ✔ Monitor the groups: Move around the classroom monitoring the groups to check whether everyone is working and intervene where necessary.
- ✔ Choose some groups or learners to come and present the result of their work.
- ✔ Harmonise the answers given by learners.

1.8. General guidance on assessment

Assessment is regarded as those formal and informal procedures that teachers and learners employ in gathering information on learning and making judgment about what learners know and can do.

Competence based assessment is an assessment process in which a learner is confronted with a complex situation relevant to his/her everyday life and asked to look for a solution by applying what he/ she has learnt (knowledge, skills, competences and attitudes). The purpose of this is to evaluate what learners can do or what changes/ behavior they should have. Evaluation is when facts from the assessment are used to make a decision.

Types of assessment

- ④ Formative or continuous assessment
- ④ Summative assessment.

Formative assessment is commonly referred to as assessment for learning, in which focus is on monitoring learner response to and progress with instruction. Formative assessment provides immediate feedback to both the teacher and the learner regarding the learning process. Formative and summative assessments contribute in different ways to the larger goals of the assessment process.

Purpose of formative assessment

- a) Determine the extent to which learning objectives and competences are being achieved.
- b) Diagnose or detect learning errors.
- c) Decide on the next steps in terms of progression.
- d) Keep records and measure progress.
- e) Identify learners with extra learning abilities.
- f) Motivate learners to learn and succeed.
- g) Check effectiveness of teaching methods (variety, appropriateness, relevance or need for new approaches/ strategies).

- h) Provide feedback to learners, parents and teachers.
- i) Guide learners to take control of their own learning.

Purpose of summative assessment

- a) Mainly concerned with appraisal of work in terms of units of work completed.
- b) Comes at the end of the unit course or program.
- c) Used for:
 - ✔ Selection
 - ✔ Certification
 - ✔ Curriculum control
 - ✔ Guidance on future courses
 - ✔ Promotion
 - ✔ Accountability

When to assess

Assessment should be clearly visible in lesson, unit, term and yearly plans.

- ✔ **Before learning (diagnostic):** At the beginning of a new lesson, find out what learners already know and can do, and check whether they are at the same level.
- ✔ **During learning (formative/continuous):** When learners appear to be having difficulty in some of the work; by using on-going assessment (continuous). The assessment aims at giving learners support and feedback.
- ✔ **After learning (summative):** At the end of a section of work or a learning unit, you have to assess the learners. This is also known as **assessment of learning** to establish and record overall progress of learners towards full achievement.

What to assess

The assessment should focus on correctness of answers, coherence of ideas, logical reasoning and understanding. It should also focus on the learner's approach to a situation, appreciation of the task given, impression of a situation, manipulation, reasoning, persistence and tolerance. Learners should show evidence of the ability to perform and accomplish a given task through aptitude, and/or use practical tests and evaluation of the final outcome learning.

Note that when assessing, if the assessment is a question paper, the questions should be:

- ✔ Clear, simple and straight forward.
- ✔ Short and precise.
- ✔ Free of bias.
- ✔ Readable.
- ✔ Original.
- ✔ Indicate marks for each.
- ✔ Follow order of difficulty.
- ✔ Contain a variety of action verbs.

Section 2: Content map

	Unit 1	Unit 2
Unit Title	Complex numbers	Logarithmic and exponential functions
Number of Periods	36 + homework	28 + homework
Key Unit competence	Perform operations on complex numbers in different forms and use complex numbers to solve related problems in Physics (voltage and current in alternating current), computer Science (fractals), Trigonometry (Euler’s formula to transform trigonometric expressions).	Extend the concepts of functions to investigate fully logarithmic and exponential functions, finding the domain of definition, the limits, asymptotes, variations, graphs , and model problems about interest rates, population growth or decay, magnitude of earthquake, etc
Equipment, learning and teaching materials required	Instruments of geometry and Scientific calculator	Instruments of geometry and Scientific calculator
List of generic competence practiced	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills • Research 	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills
Activities/ Techniques	Group work, pairing, practical, research	Group work, pairing, practical, research
List of cross cutting issues to be addressed	<ul style="list-style-type: none"> • Peace and values education • Inclusive education 	<ul style="list-style-type: none"> • Peace and values education • Inclusive education • Financial education

Assessments strategies of the key unit competence	<ul style="list-style-type: none"> • Formative assessments through activities • Summative assessments through exercises and end of unit assessments. 	<ul style="list-style-type: none"> • Formative assessments through activities • Summative assessments through exercises and end of unit assessments.
----------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	Unit 3	Unit 4
Unit Title	Taylor and Maclaurin's Expansions	Integration
Number of Periods	14 + homework	42 + homework
Key Unit competence	Use Taylor and Maclaurin's expansion to solve problems about approximations, limits, integration,... Extend the Maclaurin's expansion to Taylor series.	Use integration as the inverse of differentiation and as the limit of a sum, then apply it to find area of plane surfaces, volumes of solid of revolution, lengths of curved lines.
Equipment, learning and teaching materials required	Instruments of geometry and Scientific calculator	Instruments of geometry and Scientific calculator
List of generic competence practiced	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills 	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills • Research
Activities/ Techniques	Group work, pairing	Group work, pairing, practical
List of cross cutting issues to be addressed	<ul style="list-style-type: none"> • Peace and values education • Inclusive education 	<ul style="list-style-type: none"> • Peace and values education • Inclusive education

Assessments strategies of the key unit competence	<ul style="list-style-type: none"> • Formative assessments through activities • Summative assessments through exercises and end of unit assessments. 	<ul style="list-style-type: none"> • Formative assessments through activities. • Summative assessments through exercises and end of unit assessments.
----------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	Unit 5	Unit 6
Unit Title	Differential equations	Intersection and Sum of vector Subspaces
Number of Periods	21 + homework	14 + homework
Key Unit competence	Use ordinary differential equations of first and second order to model and solve related problems in Physics, Economics, Chemistry, Biology.	Relate the sum and the intersection of subspaces of a vector space by the dimension formula.
Equipment, learning and teaching materials required	Scientific calculator	Scientific calculator
List of generic competence practiced	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills • Research 	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills
Activities/ Techniques	Group work, pairing, research	Group work, pairing
List of cross cutting issues to be addressed	<ul style="list-style-type: none"> • Peace and values education • Inclusive education 	<ul style="list-style-type: none"> • Peace and values education • Inclusive education

Assessments strategies of the key unit competence	<ul style="list-style-type: none"> • Formative assessments through activities. • Summative assessments through exercises and end of unit assessments. 	<ul style="list-style-type: none"> • Formative assessments through activities. • Summative assessments through exercises and end of unit assessments.
----------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	Unit 7	Unit 8
Unit Title	Transformation of Matrices	Conics
Number of Periods	29 + homework	35 + homework
Key Unit competence	Transform matrices to an echelon form or to diagonal matrix and use the results to solve simultaneous linear equations or to calculate the n th power of a matrix.	Determine the characteristics and the graph of a conic given by its Cartesian, parametric or polar equation. Find the Cartesian, parametric and polar equations of a conic from its characteristics.
Equipment, learning and teaching materials required	Scientific calculator	Instrument of geometry, Scientific calculator
List of generic competence practiced	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills 	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills
Activities/ Techniques	Group work, pairing	Group work, pairing, practical, research
List of cross cutting issues to be addressed	<ul style="list-style-type: none"> • Peace and values education • Inclusive education 	<ul style="list-style-type: none"> • Peace and values education • Inclusive education

Assessments strategies of the key unit competence	<ul style="list-style-type: none"> • Formative assessments through activities. • Summative assessments through exercises and end of unit assessments. 	<ul style="list-style-type: none"> • Formative assessments through activities. • Summative assessments through exercises and end of unit assessments.
----------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	Unit 9
Unit Title	Random variables
Number of Periods	33 + homework
Key Unit competence	Calculate and interpret the parameters of a random variable (discrete or continuous) including binomial and the Poisson distributions.
Equipment, learning and teaching materials required	Instrument of geometry, Scientific calculator
List of generic competence practiced	<ul style="list-style-type: none"> • Critical thinking • Communication • Cooperation, interpersonal management and life skills • Research
Activities/ Techniques	Group work, pairing, practical
List of cross cutting issues to be addressed	<ul style="list-style-type: none"> • Peace and values education • Inclusive education
Assessments strategies of the key unit competence	<ul style="list-style-type: none"> • Formative assessments through activities. • Summative assessments through exercises and end of unit assessments.

Section 3: Sample lesson plan

School: Academic year:

Teacher's name:

Term	Date	Subject	Class	Unit No	Lesson No	Duration	Class size
1	Mathematics	S6 MEG	1	1 of 27	40 minutes	35
Type of Special Educational Needs and number of learners				<ul style="list-style-type: none"> • 3 low vision learners: Give them places where they are able to see what is written on the blackboard. Avoid making their own group otherwise it can be considered as segregation. • 4 learners with extra abilities: To encourage them to explain, to each other and help their classmates. 			
Unit title		Complex numbers					
Key Unit Competence:		Perform operations on complex numbers in different forms and use complex numbers to solve related problems in Physics (voltage and current in alternating current), computer Science (fractals), Trigonometry (Euler's formula to transform trigonometric expressions).					
Title of the lesson		Concepts of complex numbers					
Instructional objective		Through examples, learners should be able to define a complex number, show real part and imaginary part of a complex number and show that two complex numbers are equal or not equal accurately.					
Plan for this Class		Location: Classroom Learners are organised into groups					
Learning Materials		Exercise book, pen, calculator, ruler					
References		Advanced Mathematics for Rwanda Schools, Learner's Book Senior 6					

Timing for each step	DESCRIPTION OF TEACHING AND LEARNING ACTIVITY		Competences and cross cutting issues to be addressed
	Teacher's activities	Learners' activities	
Introduction 5 minutes	Ask a question, on how to solve quadratic equations in set of real numbers using discriminant method (Including case where the discriminant is negative).	<p>Questions</p> <p>By using discriminant method, solve in \mathbb{R}</p> <ol style="list-style-type: none"> $x^2 + 7x + 10 = 0$ $x^2 + 4x + 4 = 0$ $x^2 + x + 4 = 0$ <p>Solution</p> <ol style="list-style-type: none"> $x^2 + 7x + 10 = 0$ $\Delta = 49 - 40 = 9$ $x_1 = \frac{-7+3}{2} = -2, x_2 = \frac{-7-3}{2} = -5$ $S = \{-5, -2\}$ $x^2 + 4x + 4 = 0$ $\Delta = 16 - 16 = 0$ $x_1 = x_2 = \frac{-4}{2} = -2$ $S = \{-2\}$ $x^2 + x + 4 = 0$ $\Delta = 1 - 16 = -15 < 0$ No real solution 	Students are developing communication skills when they are explaining and sharing ideas.

<p>Development of the lesson</p> <p>5 minutes</p>	<p>Step 1:</p> <p>Form groups</p> <ul style="list-style-type: none"> • Ask learners to do activity 1.1 in Learner's Book page 2 in their groups. • Go round to check the progress of the discussion, and intervene where necessary. • Guide learners with special educational needs on how to do activity. 	<p>In their groups, learners will do activity 1.1. In their exercise book using the fact that $\sqrt{-1} = i$, they will find two numbers, a and b, whose sum is 6 and whose product is 18.</p>	<p>Cooperation and interpersonal management developed through working in groups.</p>
<p>10 minutes</p>	<p>Step 2:</p> <p>Ask groups to present their work on the chalkboard.</p>	<p>Secretary presents the work.</p> <p>Learners interact through questions and comments.</p> <p>Answers</p> <p>Recall that a quadratic equation is written as</p> $x^2 - sx + p = 0,$ <p>where s and p are the sum and product of two roots respectively.</p> <p>Then, we need to solve the equation</p> $x^2 - 6x + 18 = 0$ $\Delta = (-6)^2 - 4(18)$ $= 36 - 72$ $= -36$	<p>Communication: Learners communicate and convey information and ideas through speaking when they are presenting their work.</p> <p>Self confidence: Learners will gain self confidence competence when they are presenting their work.</p> <p>In group activities, the fact of being convinced without fighting, peace and education values are developed too.</p>

		$x_1 = a = \frac{6 + \sqrt{-36}}{2}$ <p>and</p> $x_2 = b = \frac{6 - \sqrt{-36}}{2}$ <p>Now, if $\sqrt{-1} = i$ or $i^2 = -1$, we have</p> $a = \frac{6 + \sqrt{36 \times i^2}}{2}$ $b = \frac{6 - \sqrt{36 \times i^2}}{2}$ <p>or</p> $a = 3 + 3i, b = 3 - 3i$	
<p>Conclusion 5 minutes</p>	<p>Ask learners to give the main points of the learned lesson in summary.</p>	<p>Summarise the learned lesson:</p> <p>A complex number is a number that can be put in the form $a + bi$, where a and b are real numbers and $i = \sqrt{-1}$ (i being the first letter of the word “imaginary”).</p> <p>The set of all complex numbers is denoted by \mathbb{C} and is defined as</p> $\mathbb{C} = \left\{ z = a + bi : a, b \in \mathbb{R} \right. \\ \left. \text{and } i^2 = -1 \right\}$ <p>The number a of the complex number $z = a + bi$ is called the real part of z and</p>	<p>Learners develop critical thinking through generating a summary.</p> <p>Through group activities, cooperation is developed.</p>

<p>5 minutes</p>	<p>Request learners to do questions 1.a) and 2.a) of exercise 1.1 in their respective groups.</p> <p>Move around the class checking the progress of the discussion, and intervene where necessary.</p>	<p>denoted by $Re(z)$ or $\Re(z)$; the number b is called the imaginary part and denoted by $Im(z)$ or $\Im(z)$. A complex number whose real part is zero, is said to be purely imaginary, whereas a complex number whose imaginary part is zero, is said to be a real number or simply real.</p> <p>Thus,</p> $\forall x \in \mathbb{R}, x \in \mathbb{C},$ <p>which gives that $\mathbb{R} \subset \mathbb{C}$.</p> <p>Learners will do questions 1.a) and 2.a) of exercise 1.1, in Learner's Book page 4 in their respective groups.</p>	<p>Through presentation on chalkboard, communication skills are developed.</p>
<p>5 minutes</p>	<p>Request some learners to answer to questions 1.b), and 2.b) of exercise 1.1 on chalkboard.</p> <p>Ensures that learners understand the learned lesson and decide whether to repeat the lesson or to start a new a lesson next time.</p>	<p>Learners will present answers of questions 1.b) and 2.b) of exercise 1.1, in Learner's Book page 4, on chalkboard.</p>	

5 minutes	Give learners an individual evaluation (quiz) and homework in regard to the learned lesson. Lead into next lesson: Request learners to do activity 1.2 at home.	Learners will do questions 1.c) and 2.c) of exercise 1.1, in Learner's Book page 4 as individual quiz; questions 1.d) and 2.d) as assignment.	
Teacher's self evaluation	Even if the objective has been achieved, some learners don't remember how to solve a quadratic equation using discriminant method. The time management has been disturbed by revising how to use discriminant method. For this reason, next time before any activity, learners will be given a task of revising the topics related to the given activity as homework.		

General Methodology

Follow the following three steps when teaching any lesson.

a) Introduction

Review previous lesson through asking learners some questions. If there is no previous lesson, ask them pre-knowledge questions on the day lesson.

b) Body of the lesson

Give an activity to learners that will be done in groups or pairs. Invite one or more groups/pairs for presentation of their work to other groups/pairs. After activities, capture the main points from the presentation of the learners, summarise them and answer to their questions.

c) Conclusion

Ask learners what they learned in the day lesson. Request them to do exercises in their respective groups and to correct exercises on the chalkboard. Give them individual evaluation. Remember to give homework to learners. Give them two home works: one for the lesson of the day and another which will be activity for the next lesson.

Section 4: Units description

Unit 1 Complex Numbers

Unit 2 Logarithmic and Exponential Equations

Unit 3 Taylor and Maclaurin's Expansions

Unit 4 Integration


Unit 5 Differential Equations

Unit 6 Intersection and Sum of Subspaces

Unit 7 Transformation of Matrices

Unit 8 Conics

Unit 9 Random Variables



Complex numbers

Learner's Book pages 1 - 79

Key unit competence

Perform operations on complex numbers in different forms and use complex numbers to solve related problems in Physics (voltage and current in alternating current), Computer Science (fractals), Trigonometry (Euler's formulae to transform trigonometric expressions), ...

Vocabulary or key words concepts

Argand diagram: Plane representing complex plane where x -axis is called real axis and y -axis is called imaginary axis.

Affix: Coordinates of a complex number in Argand diagram.

Modulus: The distance from origin to the affix of the complex number.

Argument: Argument of complex number z is the angle the segment oz makes with the positive real x -axis.

Polar form: A way of expressing a complex number using its modulus, its argument and basic trigonometric ratios (sine and cosine).

Exponential form: A way of expressing a given complex number using its modulus, its argument and the number e .

Guidance on the problem statement

The problem statement is “Solve in set of real number the following equations $x^2 + 6x + 8 = 0$ and $x^2 + 4 = 0$ ”.

For first equation, we have solution in \mathbb{R} but second equation does not have solution in \mathbb{R} . Therefore, the set \mathbb{R} is not sufficient to contain solutions of some equations.

Since the square root of -4 does not exist in set of real number, for second equation we introduce new kind of number i such that $i^2 = -1$ and we write $x^2 = -4 = 4 \times i^2$ so that $x = \pm\sqrt{-4} = \pm\sqrt{4 \times i^2} = \pm 2i$.

List of lessons

No	Lesson title	Number of periods
1	Concepts of complex numbers	1
2	Definition and properties of number i	1
3	Geometric representation of a complex number	1
4	Modulus of a complex number	1
5	Loci related to distances	1
6	Equality of complex numbers	1
7	Addition and subtraction of complex numbers	1
8	Conjugate and opposite of a complex number	1
9	Multiplication of complex numbers	1
10	Inverse and division of complex numbers	1
11	Square root of a complex number	1
12	Linear equations in set of complex numbers	1
13	Quadratic equations in set of complex numbers	1
14	Polynomials in set of complex numbers	2
15	Argument of a complex number	2
16	Loci related to angles	1
17	Polar form of a complex number	1

18	Multiplication and division of complex numbers in polar form	1
19	Powers of complex number in polar form	1
20	N^{th} roots of a complex number	2
21	Graphical representation of n^{th} roots of a complex number	2
22	Construction of regular polygon	2
23	Exponential form of a complex number	1
24	Trigonometric number of a multiple of an angle	2
25	Linearisation of trigonometric expressions	2
26	Solving equation of the form $a \cos x + b \sin x = c$	2
27	Alternating current problem	2
Total periods		36

Lesson development

Lesson 1.1. Concepts of complex numbers

Learning objectives

Through examples, learners should be able to define a complex number, show real part and imaginary part of a complex number and show that two complex numbers are equal or not equal accurately.

Prerequisites

- ⑦ Find the square root of a positive real number.
- ⑦ The square root of a negative real number does not exist.

Teaching Aids

Exercise book, pen and calculator



Activity 1.1 Learner's Book page 2

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Here, we need to solve the equation $x^2 - 6x + 18 = 0$
Calculating the discriminant for $x^2 - 6x + 18 = 0$, we get

$$\Delta = (-6)^2 - 4(18) = 36 - 72 = -36$$

$$x_1 = a = \frac{6 + \sqrt{-36}}{2} \quad \text{and} \quad x_2 = b = \frac{6 - \sqrt{-36}}{2}$$

As $\Delta < 0$, $x^2 - 6x + 18 = 0$ has no real solutions.

2. Now, if $\sqrt{-1} = i$ or $i^2 = -1$, we get

$$a = \frac{6 + \sqrt{-36}}{2} = \frac{6 + \sqrt{36i^2}}{2} = \frac{6 + 6i}{2} = 3 + 3i$$

$$\text{and } b = \frac{6 - \sqrt{-36}}{2} = \frac{6 - \sqrt{36i^2}}{2} = \frac{6 - 6i}{2} = 3 - 3i.$$

3. a and b , are not elements of \mathbb{R} or $a, b \notin \mathbb{R}$

Synthesis

As conclusion, a complex number is a number that can be put in the form $a + bi$, where a and b are real numbers and $i = \sqrt{-1}$ (i being the first letter of the word "imaginary").

Exercise 1.1 Learner's Book page 4

1. a) $\operatorname{Re}(z) = 0, \operatorname{Im}(z) = 45$ b) $\operatorname{Re}(z) = -3, \operatorname{Im}(z) = 0$
 c) $\operatorname{Re}(z) = -1, \operatorname{Im}(z) = 3$ d) $\operatorname{Re}(z) = -10, \operatorname{Im}(z) = 7$
2. a) Real b) Purely imaginary
 c) Purely imaginary
 d) neither real nor purely imaginary

Lesson 1.2. Definition and properties of the number i **Learning objectives**

Given powers of the number i with natural exponents, learners should be able to simplify them accurately.

Prerequisites

- ▼ Properties of powers in set of real numbers.

Teaching Aids

Exercise book, pen and calculator

Activity 1.2 Learner's Book page 4

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ▼ Critical thinking
- ▼ Self confidence
- ▼ Communication
- ▼ Cooperation, interpersonal management and life skills
- ▼ Peace and values education
- ▼ Inclusive education

Answers

1. $i^3 = i^2 i^1 = -1i = -i$
 $i^4 = i^2 i^2 = (-1) \times (-1) = 1$
 $i^5 = i^2 i^2 i^1 = (-1) \times (-1) \times i = i$

$$i^6 = i^2 i^2 i^2 = (-1) \times (-1) \times (-1) = -1$$

$$i^7 = i^2 i^2 i^2 i = (-1) \times (-1) \times (-1) i = -i$$

$$i^8 = i^2 i^2 i^2 i^2 = (-1) \times (-1) \times (-1) \times (-1) = 1$$


$$i^9 = i^2 i^2 i^2 i^2 i = (-1) \times (-1) \times (-1) \times (-1) i = i$$

2. In general;

$$i^{4k} = 1, i^{4k+1} = i, i^{4k+2} = -1, i^{4k+3} = -i \quad k = 0, 1, 2, 3, 4, 5, \dots$$

Synthesis

The imaginary unit, i , “cycles” through 4 different values each time we multiply as it is illustrated in the following figure .



The powers of imaginary unit can be generalised as follows:

$$i^{4k} = 1, i^{4k+1} = i, i^{4k+2} = -1, i^{4k+3} = -i$$

Exercise 1.2 Learner's Book page 5

1. -1

2. i

3. 1

4. $-i$

5. i

6. $-i$

Lesson 1.3. Geometric representation of a complex number

Learning objectives

Given complex numbers and using a ruler, learners should be able to represent those complex numbers in Argand diagram accurately.

Prerequisites

- ☛ Remember how to represent a point (x, y) in Cartesian plane.

Teaching Aids

Exercise book, pen, instruments of geometry


Activity 1.3 Learner's Book page 5



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ☛ Critical thinking
- ☛ Communication
- ☛ Self confidence
- ☛ Cooperation, interpersonal management and life skills
- ☛ Peace and values education
- ☛ Inclusive education


Answers



Synthesis

A complex number $z = a + bi$ can be visually represented as a pair of numbers (a, b) forming a vector from the origin or point on a diagram called **Argand diagram**.

Exercise 1.3 Learner's Book page 7



Lesson 1.4. Modulus of a complex number

Learning objectives

Given a complex number, learners should be able to find its modulus correctly.

Prerequisites

- ④ Finding distance between two points in Cartesian plane.

Teaching Aids

Exercise book, pen and calculator




Activity 1.4 Learner's Book page 7

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

Curve:



1. $z = -8 = (-8, 0)$

Distance from origin is $\sqrt{(-8-0)^2 + 0} = 8$

2. $z = 2i = (0, 2)$

Distance from origin is $\sqrt{0 + (2-0)^2} = 2$

3. $z = -3 + 7i = (-3, 7)$

Distance from origin is $\sqrt{(-3-0)^2 + (7-0)^2} = \sqrt{58}$

4. $z = 3 - 4i = (3, -4)$

Distance from origin is $\sqrt{(3-0)^2 + (-4-0)^2} = 5$

Synthesis

As conclusion, the distance from the origin to point (x, y) corresponding to the complex number $z = x + yi$ is called the modulus of z and is denoted by $|z|$ or $|x + iy|$:

$$r = |z| = \sqrt{x^2 + y^2} .$$




Figure 1.1: Modulus of a complex number

Exercise 1.4 Learner's Book page 9

1) $\sqrt{5}$

2) 5

3) 1

4. $\frac{\sqrt{2}}{2}$

5) 1

6) $5\sqrt{5}$

Lesson 1.5. Loci related to distances on Argand diagram

Learning objectives

Given a condition, learners should be able to determine the locus on Argand plane precisely.

Prerequisites

- ⑦ Finding modulus of a complex number.
- ⑦ The general form of equation of a circle, a straight line, ... in Cartesian plane.

Teaching Aids

Exercise book, pen, calculator and instrument of geometry



Activity 1.5 Learner's Book page 9

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers


Let $z = x + yi$, we have

$$|x + yi - 1 + 3i| = 2 \Leftrightarrow |x - 1 + i(y + 3)| = 2$$

$$\Leftrightarrow \sqrt{(x-1)^2 + (y+3)^2} = 2 \Leftrightarrow (x-1)^2 + (y+3)^2 = 2^2$$

which is the circle of centre $(1, -3)$ or $1 - 3i$ and radius $R = 2$.

Curve



Synthesis

As conclusion, $|z| = R$ represents a circle with centre P and radius R , $|z - z_1| = R$ represents a circle with centre z_1 and radius R and $|z - z_1| = |z - z_2|$ represents a straight line, the perpendicular bisector (mediator) of the segment joining the points z_1 and z_2 .

Exercise 1.5 Learner's Book page 12

1. Circle: $3x^2 + 3y^2 + 4x + 1 = 0$, radius is $\frac{1}{3}$ and centre is $\left(-\frac{2}{3}, 0\right)$
2. a) Circle: $x^2 + y^2 = 4$, radius 2, centre at origin; (0,0)
 - b) Interior of the circle: $x^2 + y^2 = 4$, radius 2, centre at origin
 - c) Exterior of the circle: $x^2 + y^2 = 4$, radius 2, centre at origin
 - d) Circle: $(x+1)^2 + y^2 = 1$, radius 1, centre $(-1,0)$
 - e) Vertical line: $z_1 = -1$, mediator of the line segment joining points $z_1 = -1$ and $z_2 = 1$
 - f) Circle: $(x-1)^2 + (y+3)^2 = 4$, radius 2, centre $(1,-3)$

Lesson 1.6. Equality of two complex numbers**Learning objectives**

Given two complex numbers, learners should be able to show that they are equal or not and to use this concept to solve some equation accurately.

Prerequisites

- ⑦ Solving a linear equation with one unknown.
- ⑦ Solving a system of two linear equations with two unknown.

Teaching Aids

Exercise book, pen and calculator



Activity 1.6 Learner's Book page 12


In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $3 + 2i - 1 = 3 - 1 + 2i = 2 + 2i$ and $2 + 4i - 2i = 2 + 2i$.

Argand diagram



The two complex numbers are represented by the same point in Argand diagram.

From their real and imaginary parts, the two quantities have equal real parts and equal imaginary parts, so they are equal.

2. Combine like terms on the right:

$$x + 2i = x + (2x - 3)i. \text{ Since the imaginary parts}$$

$$\text{must be equal, } 2 = 2x - 3 \Rightarrow x = \frac{5}{2}.$$

3. This is interesting: we have only one equation, but two variables; it doesn't seem like there is enough information to solve.

But since we can break this into a real part and an imaginary part, we can create two equations:

$x = 3y$, $y = 2x - 4$. Doing substitution gives us

$$y = 6y - 4 \Rightarrow y = \frac{5}{5}, \text{ which gives } x = \frac{12}{5}.$$

Synthesis

As conclusion, if two complex numbers, say $a + bi$ and $c + di$ are equal, then their real parts are equal and their imaginary parts are equal. That is, $a + bi = c + di \Leftrightarrow a = c$ and $b = d$.

Exercise 1.6 Learner's Book page 13

- | | |
|-----------------------|-----------------------|
| 1. $x = 4$, $y = -3$ | 2. $x = 5$, $y = 6$ |
| 3. $x = 3$, $y = 3$ | 4. $x = -6$, $y = 9$ |
| 5. $x = 2$, $y = 5$ | 6. $x = 3$, $y = 1$ |
| 7. $x = 6$, $y = -6$ | 8. $x = 8$, $y = 14$ |

Lesson 1.7. Addition and subtraction of complex numbers

Learning objectives

Given two complex numbers, learners should be able to add and subtract them correctly.

Prerequisites

- ✔ Simplification by combining like terms.

Teaching Aids

Exercise book, pen and calculator



Activity 1.7 Learner's Book page 13

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence

- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. $z_1 + z_2 = (2 + 3i) + (5 - 4i) = 2 + 5 + 3i - 4i = 7 - i$
 $z_1 - z_2 = (2 + 3i) - (5 - 4i) = 2 - 5 + 3i + 4i = -3 + 7i$
2. $\operatorname{Re}(z_1 + z_2) = 7, \operatorname{Im}(z_1 + z_2) = -1$
 $\operatorname{Re}(z_1 - z_2) = -3, \operatorname{Im}(z_1 - z_2) = 7$

Synthesis

As conclusion, two complex numbers are added (or subtracted) by adding (or subtracting) separately the two real and the two imaginary parts.

Exercise 1.7 Learner's Book page 14

1. $z_1 + z_2 = -12, z_1 - z_2 = 12 + 6i$
2. $z_1 + z_2 = 16i, z_1 - z_2 = -10 + 8i$
3. $z_1 + z_2 = 5 + 3i, z_1 - z_2 = 1 + 5i$
4. $z_1 + z_2 = -2 - 24i, z_1 - z_2 = -44 - 4i$
5. $z_1 + z_2 = -2 + 8i, z_1 - z_2 = 8 + 12i$
6. $z_1 + z_2 = -2 - 24i, z_1 - z_2 = -44 - 4i$
7. $z_1 + z_2 = 35 - 13i, z_1 - z_2 = -9 - 15i$
8. $z_1 + z_2 = 4 + 9i, z_1 - z_2 = 2 - 11i$

Lesson 1.8. Conjugate and opposite of a complex number

Learning objectives

Given a complex number, learners should be able to find the conjugate and opposite moderately.

Prerequisites

- ✔ Plot a complex number in Argand plane
- ✔ Adding two complex numbers

Teaching Aids

Exercise book, pen and instruments of geometry




Activity 1.8 Learner's Book page 14

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Argand diagram of complex numbers



$$2. \text{ a) } \frac{1}{2}(z_1 + z_2) = \frac{1}{2}(4 + 3i + 4 - 3i) = 4$$

$$\text{b) } \frac{1}{2i}(z_1 - z_2) = \frac{1}{2i}(4 + 3i - 4 + 3i) = 3$$

$$3. \frac{1}{2}(z_1 + z_2) = \text{Re}(z_1) \text{ and } \frac{1}{2i}(z_1 - z_2) = \text{Im}(z_1)$$

Synthesis

As conclusion, the conjugate of the complex number $z = x + yi$, denoted by \bar{z} or z^* , is obtained by changing the sign of the imaginary part. Hence, the complex conjugate of $z = x + yi$ is $\bar{z} = x - yi$.

Exercise 1.8 Learner's Book page 16

$$1. -76$$

$$2. 9i$$

$$3. 12 + 4i$$

$$4. 3 - i$$

$$5. -8 - 10i$$

$$6. 3 + i$$

$$7. 3 + 5i$$

$$8. -5 - 5i$$

Lesson 1.9. Multiplication of complex numbers

Learning objectives

Given two complex numbers, learners should be able to multiply them perfectly.

Prerequisites

- ✔ Distributive property.
- ✔ Multiplication is distributive over addition.
- ✔ Relation $i^2 = -1$.

Teaching Aids

Exercise book, pen and calculator



Activity 1.9 Learner's Book page 17

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$\begin{aligned}
 1. \quad z_1 \times z_2 &= (2 - 3i)(3 + 2i) \\
 &= 6 + 4i - 9i - 6i^2 \\
 &= 6 + 6 - 5i \\
 &= 12 - 5i
 \end{aligned}$$

$$2. \quad \operatorname{Re}(z_1 \times z_2) = 12, \quad \operatorname{Im}(z_1 \times z_2) = -5$$

Synthesis

As conclusion, the multiplication of two complex numbers $z_1 = a + bi$ and $z_2 = c + di$ is defined by the following formula:

$$\begin{aligned}
 z_1 \times z_2 &= (a + bi)(c + di) \\
 &= (ac - bd) + (bc + ad)i
 \end{aligned}$$

Exercise 1.9 Learner's Book page 17

- | | | |
|---------------|----------------|--------------|
| 1. $9 - 36i$ | 2. $-73 + 40i$ | 3. $10 + 5i$ |
| 4. $-3 - 10i$ | 5. $10 - 41i$ | 6. $-4 - 7i$ |

Lesson 1.10. Inverse and division of complex numbers

Learning objectives

Given complex numbers, learners should be able to find the inverse of a complex number and divide two complex numbers accurately.

Prerequisites

- ✔ Multiplication of complex numbers.

Teaching Aids

Exercise book, pen and calculator



Activity 1.10 Learner's Book page 18

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad z_1 \cdot \bar{z}_1 = (2+i)(2-i) = 2^2 - 2i + 2i + 1^2 = 2^2 + 1^2 = 5$$

Hence, if $z = a + bi$ then, $\bar{z} = a - bi$ and $z \cdot \bar{z} = a^2 + b^2$.

$$2. \quad z_1 \cdot \bar{z}_1 = 5 \Rightarrow \bar{z}_1 = \frac{5}{z_1} \Rightarrow \frac{\bar{z}_1}{5} = \frac{1}{z_1}$$

Then,

$$\frac{1}{z_1} = \frac{\bar{z}_1}{5} \quad \text{or} \quad \frac{1}{z_1} = \frac{\bar{z}_1}{a^2 + b^2}$$

$$3. \quad \frac{1}{z_1} = \frac{\bar{z}_1}{a^2 + b^2}$$

Multiplying both sides by z_2 , we get $\frac{z_2}{z_1} = \frac{z_2 \bar{z}_1}{a^2 + b^2}$

Synthesis

The **inverse** of $z = a + bi$ is given by $z^{-1} = \frac{\bar{z}}{z \cdot \bar{z}} = \frac{\bar{z}}{a^2 + b^2}$
and the **division** of two complex

numbers is $\frac{z_1}{z_2} = \frac{z_1 \cdot \bar{z}_2}{z_2 \cdot \bar{z}_2} = \frac{a + bi}{c + di} = \left(\frac{ac + bd}{c^2 + d^2} \right) + i \left(\frac{bc - ad}{c^2 + d^2} \right)$

Exercise 1.10 Learner's Book page 21

1. $\frac{1}{z_1} = -\frac{i}{3}, \frac{1}{z_2} = -\frac{4}{51} + \frac{i}{51}, \frac{z_1}{z_2} = -\frac{1}{17} - \frac{4}{17}i$
2. $\frac{1}{z_1} = -\frac{5}{169} - \frac{12}{169}i, \frac{1}{z_2} = \frac{5}{41} - \frac{4}{41}i, \frac{z_1}{z_2} = \frac{23}{41} + \frac{80}{41}i$
3. $\frac{1}{z_1} = \frac{3}{25} - \frac{4}{25}i, \frac{1}{z_2} = \frac{2}{5} + \frac{i}{5}, \frac{z_1}{z_2} = \frac{2}{5} + \frac{11}{5}i$
4. $\frac{1}{z_1} = -\frac{23}{725} + \frac{14}{725}i, \frac{1}{z_2} = \frac{21}{541} + \frac{10}{541}i, \frac{z_1}{z_2} = -\frac{343}{541} - \frac{524}{541}i$
5. $\frac{1}{z_1} = \frac{1}{10} + \frac{3}{10}i, \frac{1}{z_2} = -\frac{1}{5} - \frac{2}{5}i, \frac{z_1}{z_2} = -\frac{7}{5} + \frac{1}{5}i$
6. $\frac{1}{z_1} = -\frac{2}{5} - \frac{1}{5}i, \frac{1}{z_2} = -\frac{5}{29} - \frac{2}{29}i, \frac{z_1}{z_2} = \frac{12}{29} - \frac{1}{29}i$
7. $\frac{1}{z_1} = -\frac{1}{2} - \frac{1}{2}i, \frac{1}{z_2} = \frac{1}{2} + \frac{1}{2}i, \frac{z_1}{z_2} = -1$
8. $\frac{1}{z_1} = -\frac{1}{74} - \frac{3}{37}i, \frac{1}{z_2} = \frac{1}{101} - \frac{10}{101}i, \frac{z_1}{z_2} = \frac{118}{101} + \frac{32}{101}i$

Lesson 1.11. Square root of a complex number**Learning objectives**

Given a complex number, learners should be able to find its square root accurately.

Prerequisites

- ✔ Solve a system of two linear equations with two unknowns.
- ✔ Use of the identity $(a + b)^2 = a^2 + 2ab + b^2$.

Teaching Aids

Exercise book, pen, calculator

**Activity 1.1 | Learner's Book page 21**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$(x + yi)^2 = 8 - 6i$$

$$\Leftrightarrow x^2 - y^2 + 2xyi = 8 - 6i$$

$$\Leftrightarrow \begin{cases} x^2 - y^2 = 8 \\ 2xy = -6 \end{cases}$$

Squaring both sides of each equation and adding two equations, gives

$$\Leftrightarrow \begin{cases} x^4 - 2x^2y^2 + y^4 = 64 \\ 4x^2y^2 = 36 \end{cases}$$

$$\underline{\hspace{10em}}$$

$$x^4 + 2x^2y^2 + y^4 = 100$$

Using algebraic identity, gives

$$\Rightarrow (x^2 + y^2)^2 = 100$$

$$\Rightarrow x^2 + y^2 = 10$$

Now,

$$\begin{cases} x^2 - y^2 = 8 \\ x^2 + y^2 = 10 \end{cases}$$

$$\underline{\hspace{10em}}$$

$$2x^2 = 18 \Rightarrow x^2 = 9 \text{ or } x = \pm 3$$

But $x^2 + y^2 = 10$, then, $y^2 = 10 - x^2 = 10 - 9 = 1$ or $y = \pm 1$

Thus, the square root of $z = 8 - 6i$ is $3 - i$ or $-3 + i$

We take different sign (for x and y) since the product xy is negative.

Synthesis

To get a square root of the complex number $a + bi$, we let $x + iy$ be a square root of the complex number $a + bi$, and solve the simultaneous equation

$$\begin{cases} (x + iy)^2 = a + bi \\ |(x + iy)^2| = |a + bi| \end{cases} \Rightarrow \begin{cases} x^2 - y^2 = a \\ 2xy = b \\ x^2 + y^2 = \sqrt{a^2 + b^2} \end{cases}$$

$$\Leftrightarrow \begin{cases} x^2 + y^2 = \sqrt{a^2 + b^2} \\ x^2 - y^2 = a \end{cases} \text{ and } 2xy = b$$

Notice:

In writing square root of the complex number $a + bi$, that is, $x + iy$, x and y must satisfy the condition $2xy = b$.

Exercise 1.11 Learner's Book page 23

- | | | |
|--------------------------------|--------------------|-------------------|
| 1) $\pm(\sqrt{7} + i\sqrt{7})$ | 2) $\pm(-4 - 6i)$ | 3) $\pm(-4 - 6i)$ |
| 4) $\pm(-3 + 10i)$ | 5) $\pm(3 + 2i)$ | 6) $\pm(-6 + 2i)$ |
| 7) $\pm(-6 - 2i)$ | 8) $\pm(-5 - 12i)$ | |

Lesson 1.12. Linear equations

Learning objectives

Given linear equations with complex coefficients, learners will be able to solve them in the set of complex numbers accurately.

Prerequisites

- ⑦ Solving linear equations in set or real numbers.

Teaching Aids

Exercise book, pen and calculator



Activity 1.12 Learner's Book page 24

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $z + 3i - 4 = 0 \Rightarrow z = 4 - 3i$
2. $4 - i + iz = 4z - 3i \Rightarrow iz - 4z = -3i - 4 + i$
 $\Rightarrow z(i - 4) = -4 - 2i$
 $\Rightarrow z = \frac{-4 - 2i}{i - 4}$
 $\Rightarrow z = \frac{14}{17} + \frac{12}{17}i$
3. $(1 + i)(i + z) = 4i \Rightarrow i + z - 1 + iz = 4i$
 $\Rightarrow z(i + 1) = 4i + 1 - i$
 $\Rightarrow z(i + 1) = 1 + 3i$
 $\Rightarrow z = \frac{1 + 3i}{i + 1} = 2 + i$
4. $(1 - i)z = 2 + i \Rightarrow z = \frac{2 + i}{1 - i} \Rightarrow z = \frac{1 + 3i}{2}$

Synthesis

As conclusion, in complex numbers also, we may need to find the complex number z that satisfies the given linear equation.

Exercise 1.12 Learner's Book page 25

1. $2 + 2i$
2. $-4 - 2i$
3. $-7 + 6i$
4. $\frac{9}{5} + \frac{3}{5}i$

Lesson 1.13. Quadratic equations

Learning objectives

Given quadratic equations, learners should be able to solve them in the set of complex numbers correctly.

Prerequisites

- ✔ Discriminant method used to solve a quadratic equation.
- ✔ Relation $i = \sqrt{-1}$.

Teaching Aids

Exercise book, pen and calculator



Activity 1.13 Learner's Book page 25

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad x^2 + 2x + 3 = 0$$

$$\Delta = 4 - 12 = -8$$

$$x_1 = \frac{-2 + \sqrt{-8}}{2} = \frac{-2 + 2i\sqrt{2}}{2} = -1 + i\sqrt{2}$$

$$x_2 = \frac{-2 - \sqrt{-8}}{2} = \frac{-2 - 2i\sqrt{2}}{2} = -1 - i\sqrt{2}$$

$$S = \{-1 + i\sqrt{2}, -1 - i\sqrt{2}\}$$

$$2. \quad x^2 + 2x + 1 + i = 0$$

$$\Delta = 4 - 4(1+i) = 4 - 4 - 4i = -4i$$

$$\sqrt{\Delta} = \sqrt{-4i}$$

$$\sqrt{\Delta} = \sqrt{2} - i\sqrt{2} \text{ or } -\sqrt{2} + i\sqrt{2}$$

$$x_1 = \frac{-2 + \sqrt{2} - i\sqrt{2}}{2} = \frac{-2 + \sqrt{2}}{2} - i\frac{\sqrt{2}}{2}$$

$$x_2 = \frac{-2 - \sqrt{2} + i\sqrt{2}}{2} = \frac{-2 - \sqrt{2}}{2} + i\frac{\sqrt{2}}{2}$$

$$S = \left\{ \frac{-2 + \sqrt{2}}{2} - i\frac{\sqrt{2}}{2}, \frac{-2 - \sqrt{2}}{2} + i\frac{\sqrt{2}}{2} \right\}$$

Synthesis

In solving equation $az^2 + bz + c = 0$ where a , b and c are real numbers ($a \neq 0$), we get either:

- ④ Two real roots (if $\Delta > 0$); $z_1 = \frac{-b + \sqrt{\Delta}}{2a}$ and $z_2 = \frac{-b - \sqrt{\Delta}}{2a}$
- ④ One double real root (if $\Delta = 0$); $z_1 = z_2 = \frac{-b}{2a}$ or
- ④ Two conjugate complex roots (if $\Delta < 0$):

$$z_1 = \frac{-b + i\sqrt{-\Delta}}{2a} \text{ and } z_2 = \frac{-b - i\sqrt{-\Delta}}{2a}.$$

Exercise 1.13 Learner's Book page 27

1. $S = \left\{ \frac{2 + i\sqrt{26}}{3}, \frac{2 - i\sqrt{26}}{3} \right\}$
2. $S = \{5 + 3i, 5 - 3i\}$
3. $S = \left\{ \frac{3}{2} + \frac{\sqrt{11}}{2}i, \frac{3}{2} - \frac{\sqrt{11}}{2}i \right\}$

Lesson 1.14. Polynomials in set of complex numbers

Learning objectives

Given a polynomial with complex coefficients, learners should be able to factorise completely it in set of complex numbers accurately.

Prerequisites

- ④ Finding zero of a polynomial.
- ④ Use of synthetic division.

Teaching Aids

Exercise book, pen and calculator

**Activity 1.14 Learner's Book page 28**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$1. \quad a) \quad (z - 2 - 3i)(z + 3 + i) = z^2 + 3z + iz - 2z - 6 - 2i - 3zi - 9i + 3$$

$$= z^2 + (1 - 2i)z - 3 - 11i$$

$$b) \quad (z - i)(z + 3i)(z - 4i) = (z^2 + 3zi - iz + 3)(z - 4i)$$

$$= (z^2 + 2iz + 3)(z - 4i)$$

$$= z^3 - 4iz^2 + 2iz^2 + 8z + 3z - 12i$$

$$= z^3 - 2iz^2 + 11z - 12i$$

$$2. \quad P(z) = z^3 + (-2 - i)z^2 + (2 + 2i)z - 4 \text{ is divisible by } z + i \text{ if and only if } P(-i) = 0.$$

$$P(-i) = (-i)^3 + (-2 - i)(-i)^2 + (2 + 2i)(-i) - 4$$

$$= i + 2 + i - 2i + 2 - 4$$

$$= 0$$

Thus, $P(z) = z^3 + (-2 - i)z^2 + (2 + 2i)z - 4$ is divisible by $z + i$.

Now, using synthetic division

	1	$-2 - i$	$2 + 2i$	-4
$-i$		$-i$	$2i - 2$	4
	1	$-2 - 2i$	$4i$	0

$$\begin{aligned}
 P(z) &= z^3 + (-2-i)z^2 + (2+2i)z - 4 \\
 &= (z+i)[z^2 + (-2-2i)z + 4i]
 \end{aligned}$$

Again, we factorise $z^2 + (-2-2i)z + 4i$ since 2 is a root, then

	1	$-2-2i$	$4i$
2		2	$-4i$
	1	$-2i$	0

$$z^2 + (-2-2i)z + 4i = (z-2)(z-2i)$$

$$\text{Thus, } P(z) = (z+i)(z-2)(z-2i)$$

3. If $P(z) = z^3 - 2z^2 + (7+2i)z - 6(2-i)$

$2-i$ is a factor of $-6(2-i)$,

$$\begin{aligned}
 P(2-i) &= (2-i)^3 - 2(2-i)^2 + (7+2i)(2-i) - 6(2-i) \\
 &= 2 - 11i - 6 + 8i + 16 - 3i - 12 + 6i \\
 &= 0
 \end{aligned}$$

Other values are: $z = 3i, z = -2i$

All roots can be found as follows:

$$P(z) = z^3 - 2z^2 + (7+2i)z - 12 + 6i$$

$$P(z) = 0 \Leftrightarrow z^3 - 2z^2 + (7+2i)z - 12 + 6i = 0$$

$z = 3i$ is a root since

$$\begin{aligned}
 P(3i) &= (3i)^3 - 2(3i)^2 + (7+2i)(3i) - 12 + 6i \\
 &= -27i + 18 + 21i - 6 - 12 + 6i \\
 &= 0
 \end{aligned}$$

Using Synthetic division, we have

	1	-2	$7+2i$	$-12+6i$
$3i$		$3i$	$-6i-9$	$-6i+12$
	1	$-2+3i$	$-2-4i$	0

$$\begin{aligned}
 P(z) &= z^3 - 2z^2 + (7+2i)z - 12 + 6i \\
 &= (z-3i)[z^2 + (-2+3i)z - 2 - 4i]
 \end{aligned}$$

$-2i$ is also a root

	1	$-2 + 3i$	$-2 - 4i$
$-2i$		$-2i$	$4i + 2$
	1	$-2 + i$	0

$$P(z) = z^3 - 2z^2 + (7 + 2i)z - 12 + 6i$$

$$= (z - 3i)(z + 2i)(z - 2 + i)$$

Then, $z = 3i$ or $z = -2i$ or $z = 2 - i$

Synthesis

As conclusion, the process of finding the roots of a polynomial in set of complex numbers is similar to the case of real numbers remembering that the square root of a negative real number exist in set of complex numbers considering $\sqrt{-1} = i$. The methods used are synthetic division and factorisation.

Exercise 1.14 Learner's Book page 31

- $P(z) = (z + 2)(z - 1 + 3i)(z - 1 - 3i); \{-2, 1 - 3i, 1 + 3i\}$
 - $Q(z) = (z - 2)(z - 2 + i)(z - 2 - i); \{2, 2 - i, 2 + i\}$
 - $R(z) = (z - 2 + 3i)(z - 2 - 3i)(z + 2 - \sqrt{2})(z + 2 + \sqrt{2});$
 $\{2 - 3i, 2 + 3i, -2 + \sqrt{2}, -2 - \sqrt{2}\}$
 - $M(z) = (z - 3i)(z + 2i)(z - 2 + i); \{3i, -2i, 2 - i\}$
- $a = 79, b = 29$
- $p(z) = -2z^3 + 8z^2 - 18z + 20$
- $S = \left\{3 - i, -\frac{7}{2} + \frac{7}{2}i\right\}$

Lesson 1.15. Argument of a complex number

Learning objectives

Given a complex numbers, learners should be able to find its argument moderately.

Prerequisites

- ✔ Concepts of trigonometry.

Teaching Aids

Exercise book, pen, scientific calculator and instruments of geometry


Activity 1.15 Learner's Book page 31



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers



For $z_1 = 1 + i$, $\tan \theta = \frac{1}{1} = 1 \Rightarrow \theta = \arctan(1) = \frac{\pi}{4}$. From the figure, this is the needed angle.

For $z_2 = 1 - i$, $\tan \theta = \frac{1}{-1} = -1 \Rightarrow \theta = \arctan(-1) = -\frac{\pi}{4}$. From the figure, this is the needed angle.

For $z_3 = -1 + i$, $\tan \theta = \frac{-1}{1} = -1 \Rightarrow \theta = \arctan(-1) = -\frac{\pi}{4}$. From the figure, this is not the needed angle. The needed angle is $\pi + \theta = \pi - \frac{\pi}{4} = \frac{3\pi}{4}$

For $z_4 = -1 - i$, $\tan \theta = \frac{-1}{-1} = 1 \Rightarrow \theta = \arctan(1) = \frac{\pi}{4}$. From the figure, this is not the needed angle. The needed angle is $\theta - \pi = \frac{\pi}{4} - \pi = -\frac{3\pi}{4}$.

For $z_5 = i$. From the figure, the needed angle is $\theta = \frac{\pi}{2}$

For $z_6 = -i$. From the figure, the needed angle is $\theta = -\frac{\pi}{2}$

Synthesis

Depending on the quadrant in which the argument of complex number $z = x + yi$ lies, we define $\arg(z)$ as follows:

$$\arg(z) = \begin{cases} \arctan \frac{y}{x}, & \text{if } z \text{ lies in } 1^{\text{st}} \text{ or } 4^{\text{th}} \text{ quadrant or on positive } x\text{-axis} \\ \pi + \arctan \frac{y}{x}, & \text{if } z \text{ lies in } 2^{\text{nd}} \text{ quadrant or on negative } x\text{-axis} \\ -\pi + \arctan \frac{y}{x}, & \text{if } z \text{ lies in } 3^{\text{rd}} \text{ quadrant} \\ \frac{\pi}{2}, & \text{if } z \text{ lies on positive } y\text{-axis} \\ -\frac{\pi}{2}, & \text{if } z \text{ lies on negative } y\text{-axis} \\ \text{undefined}, & \text{if } x = 0 \text{ and } y = 0 \end{cases}$$

This is equivalent to

$$\arg(z) = \begin{cases} \arctan \frac{y}{x}, & \text{if } x > 0 \\ \pi + \arctan \frac{y}{x}, & \text{if } x < 0 \text{ and } y \geq 0 \\ -\pi + \arctan \frac{y}{x}, & \text{if } x < 0 \text{ and } y < 0 \\ \frac{\pi}{2}, & \text{if } x = 0, y > 0 \\ -\frac{\pi}{2}, & \text{if } x = 0, y < 0 \\ \text{undefined}, & \text{if } x = 0 \text{ and } y = 0 \end{cases}$$




Figure 1.2: Argument of a complex number

Exercise 1.15 Learner's Book page 34

- | | | |
|---------------------|---------------------|--------------------|
| 1. $-\frac{\pi}{4}$ | 2. $-\frac{\pi}{6}$ | 3. $\frac{\pi}{4}$ |
| 4. $-\frac{\pi}{3}$ | 5. $-\frac{\pi}{6}$ | |

Lesson 1.16. Loci related to the angles

Learning objectives

Given an argument condition, learners should be able to sketch on Argand diagram, the region satisfying that condition accurately.

Prerequisites

- ✔ Drawing angle with a given size.

Teaching Aids

Exercise book, pencil and instruments of geometry



Activity 1.16 Learner's Book page 34


In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers


$$1. \arg(z) = \frac{\pi}{4} \Rightarrow \arg(x + yi) = \frac{\pi}{4}$$

Here, we need all complex numbers lying on the half line passing through $(0,0)$ and makes an angle of $\frac{\pi}{4}$ with positive x -axis.



$$2. \arg(z-4) = \frac{\pi}{3} \Rightarrow \arg(x+yi-4) = \frac{\pi}{3} \Rightarrow \arg(x-4+yi) = \frac{\pi}{3}$$

Here, we need all complex numbers lying on the half line passing through $(4,0)$ and makes an angle of $\frac{\pi}{3}$ with positive x -axis.



Synthesis

As conclusion, $\arg(z) = \theta$ represents the half line through O inclined at an angle θ to the positive direction of x -axis.




Figure 1.3: Locus as a half line through O

$\arg(z-z_1) = \theta$ represents the half line through the point z_1 inclined at an angle θ to the positive direction of x -axis.




Figure 1.4: Locus as a half line through any point

$\theta \leq \arg(z - z_1) \leq \beta$ indicates that the angle between AP and the positive x-axis lies between θ and β , so that P can lie on or within the two half lines as shown in Figure 5.1.






Figure 1.5: Locus between two half lines

Exercise 1.16 Learner's Book page 37


1. a)



b)



2.



a) From the graph, we see that there is only one point of intersection. Thus, there is only one complex number satisfying both conditions.

b) Putting $z = -7 - 4i$, we have

$$|-7 - 4i + 3 + i| = |-4 - 3i| = \sqrt{16 + 9} = 5 \text{ also}$$

$$\arg(-7 - 4i + 3) = \arg(-4 - 4i) = -\pi + \arctan(1) = -\pi + \frac{\pi}{4} = -\frac{3\pi}{4}$$

Thus, $z = -7 - 4i$ verifies both conditions.

Lesson 1.17. Polar form of a complex number

Learning objectives

Given a complex number, learners should be able to express it in polar form accurately.

Prerequisites

- ✔ Finding the modulus of a complex number.
- ✔ Finding argument of a complex number.

Teaching Aids

Exercise book, pen and scientific calculator




Activity 1.17 Learner's Book page 38

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Graph



$$2. r = \sqrt{16+16} = 4\sqrt{2}$$

$$3. \theta = \arctan\left(\frac{4}{4}\right) = \frac{\pi}{4}$$

$$4. \cos \theta = \frac{x}{r} \Rightarrow x = r \cos \theta \Rightarrow 4 = 4\sqrt{2} \cos \frac{\pi}{4}$$

$$\sin \theta = \frac{y}{r} \Rightarrow y = r \sin \theta \Rightarrow 4 = 4\sqrt{2} \sin \frac{\pi}{4}$$

From $z = 4 + 4i$, we have

$$z = 4\sqrt{2} \cos \frac{\pi}{4} + i4\sqrt{2} \sin \frac{\pi}{4} = 4\sqrt{2} \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right)$$

Synthesis

As conclusion, if r and θ are the modulus and principal argument of complex number z respectively, then the polar form of z is $z = r(\cos \theta + i \sin \theta)$.




Figure 1.6: Modulus and argument of a complex number

Exercise 1.17 Learner's Book page 40

1. a) $4 \operatorname{cis} 0$ b) $2 \operatorname{cis} \frac{\pi}{2}$ c) $2 \operatorname{cis}(\pi)$
- d) $5 \operatorname{cis}\left(-\frac{\pi}{2}\right)$ e) $2 \operatorname{cis} \frac{\pi}{6}$ f) $2\sqrt{3} \operatorname{cis}\left(-\frac{\pi}{3}\right)$
- g) $2\sqrt{3} \operatorname{cis}\left(-\frac{\pi}{6}\right)$

Here, remember that the notation $r \operatorname{cis} \theta$ is the same as $r(\cos \theta + i \sin \theta)$.

$$2. \quad \begin{array}{lll} \text{a) } 1+i\sqrt{3} & \text{b) } \sqrt{2}(-2+2i) & \text{c) } 1-i \\ \text{d) } 3i & \text{e) } -4 & \text{f) } \frac{1}{2}(-\sqrt{3}-i) \\ \text{g) } \sqrt{3}-i & & \end{array}$$

Lesson 1.18. Multiplication and division of complex numbers in polar form

Learning objectives

Given two complex numbers, learners should be able to multiply and divide them in polar form exactly.

Prerequisites

- ✔ Putting a complex number in polar form
- ✔ Addition formulae in trigonometry.

Teaching Aids

Exercise book, pen and scientific calculator



Activity 1.18 Learner's Book page 41

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad \begin{aligned} z_1 z_2 &= (1-i)(\sqrt{3}-i) \\ &= \sqrt{3}-i-i\sqrt{3}-1 = \sqrt{3}-1-(\sqrt{3}+1)i \\ |z_1 z_2| &= \sqrt{(\sqrt{3}-1)^2 + (\sqrt{3}+1)^2} = \sqrt{3-2\sqrt{3}+1+3+2\sqrt{3}+1} = \sqrt{8} = 2\sqrt{2} \\ \arg(z_1 z_2) &= \arctan\left(\frac{-\sqrt{3}-1}{\sqrt{3}-1}\right) = -\frac{5\pi}{12} \end{aligned}$$

$$\text{Then, } z_1 z_2 = 2\sqrt{2} \operatorname{cis}\left(\frac{-5\pi}{12}\right)$$

$$2. \quad z_1 = 1 - i$$

$$|z_1| = \sqrt{2}, \quad \arg(z_1) = \arctan(-1) = -\frac{\pi}{4} \Rightarrow z_1 = \sqrt{2} \operatorname{cis}\left(-\frac{\pi}{4}\right)$$

$$z = \sqrt{-i}$$

$$|z_2| = 2, \quad \arg(z_2) = \arctan\left(-\frac{1}{\sqrt{3}}\right) = -\frac{\pi}{6} \Rightarrow z_2 = 2 \operatorname{cis}\left(-\frac{\pi}{6}\right)$$

$$\begin{aligned} z_1 z_2 &= \sqrt{2} \operatorname{cis}\left(-\frac{\pi}{4}\right) 2 \operatorname{cis}\left(-\frac{\pi}{6}\right) = 2\sqrt{2} \left[\cos\left(-\frac{\pi}{4}\right) + i \sin\left(-\frac{\pi}{4}\right) \right] \left[\cos\left(-\frac{\pi}{6}\right) + i \sin\left(-\frac{\pi}{6}\right) \right] \\ &= 2\sqrt{2} \left[\cos\left(-\frac{\pi}{4}\right) \cos\left(-\frac{\pi}{6}\right) + i \cos\left(-\frac{\pi}{4}\right) \sin\left(-\frac{\pi}{6}\right) + i \sin\left(-\frac{\pi}{4}\right) \cos\left(-\frac{\pi}{6}\right) - \sin\left(-\frac{\pi}{4}\right) \sin\left(-\frac{\pi}{6}\right) \right] \\ &= 2\sqrt{2} \left[\cos\left(-\frac{\pi}{4}\right) \cos\left(-\frac{\pi}{6}\right) - \sin\left(-\frac{\pi}{4}\right) \sin\left(-\frac{\pi}{6}\right) + i \left(\cos\left(-\frac{\pi}{4}\right) \sin\left(-\frac{\pi}{6}\right) + \sin\left(-\frac{\pi}{4}\right) \cos\left(-\frac{\pi}{6}\right) \right) \right] \\ &= 2\sqrt{2} \left[\cos\left(-\frac{\pi}{4} + \left(-\frac{\pi}{6}\right)\right) + i \left(\sin\left(-\frac{\pi}{4} + \left(-\frac{\pi}{6}\right)\right) \right) \right], \quad \text{from addition formulae in trigonometry} \\ &= 2\sqrt{2} \left[\cos\left(-\frac{5\pi}{12}\right) + i \cos\left(-\frac{5\pi}{12}\right) \right] = 2\sqrt{2} \operatorname{cis}\left(\frac{-5\pi}{12}\right) \end{aligned}$$

3. The two results are the same

$$\begin{aligned} 4. \quad \frac{z_1}{z_2} &= \frac{1-i}{\sqrt{3}-i} \\ &= \frac{(1-i)(\sqrt{3}+i)}{4} = \frac{\sqrt{3}+i-i\sqrt{3}+1}{4} = \frac{\sqrt{3}+1}{4} + \frac{1-\sqrt{3}}{4}i \end{aligned}$$

$$\left| \frac{z_1}{z_2} \right| = \frac{|z_1|}{|z_2|} = \frac{|1-i|}{|\sqrt{3}-i|} = \frac{\sqrt{2}}{2}$$

$$\arg\left(\frac{z_1}{z_2}\right) = \arctan\left(\frac{\frac{1-\sqrt{3}}{4}}{\frac{\sqrt{3}+1}{4}}\right) = -\frac{\pi}{12}$$

Then,

$$\frac{z_1}{z_2} = \frac{\sqrt{2}}{2} \operatorname{cis}\left(-\frac{\pi}{12}\right)$$

$$5. \quad z_1 = \sqrt{2} \operatorname{cis}\left(-\frac{\pi}{4}\right), \quad z_2 = 2 \operatorname{cis}\left(-\frac{\pi}{6}\right)$$

$$\begin{aligned}
 \frac{z_1}{z_2} &= \frac{\sqrt{2} \operatorname{cis}\left(-\frac{\pi}{4}\right)}{2 \operatorname{cis}\left(-\frac{\pi}{6}\right)} = \frac{\sqrt{2} \left[\cos\left(-\frac{\pi}{4}\right) + i \sin\left(-\frac{\pi}{4}\right) \right]}{2 \left[\cos\left(-\frac{\pi}{6}\right) + i \sin\left(-\frac{\pi}{6}\right) \right]} \\
 &= \frac{\sqrt{2} \left[\cos\left(-\frac{\pi}{4}\right) + i \sin\left(-\frac{\pi}{4}\right) \right] \left[\cos\left(-\frac{\pi}{6}\right) - i \sin\left(-\frac{\pi}{6}\right) \right]}{2 \left[\cos\left(-\frac{\pi}{6}\right) + i \sin\left(-\frac{\pi}{6}\right) \right] \left[\cos\left(-\frac{\pi}{6}\right) - i \sin\left(-\frac{\pi}{6}\right) \right]} \\
 &= \frac{\sqrt{2} \left[\cos\left(-\frac{\pi}{4}\right) \cos\left(-\frac{\pi}{6}\right) - i \sin\left(-\frac{\pi}{6}\right) \cos\left(-\frac{\pi}{4}\right) + i \cos\left(-\frac{\pi}{6}\right) \sin\left(-\frac{\pi}{4}\right) + \sin\left(-\frac{\pi}{4}\right) \sin\left(-\frac{\pi}{6}\right) \right]}{2 \left[\cos^2\left(-\frac{\pi}{6}\right) + \sin^2\left(-\frac{\pi}{6}\right) \right]} \\
 &= \frac{\sqrt{2} \left[\cos\left(-\frac{\pi}{4}\right) \cos\left(-\frac{\pi}{6}\right) + \sin\left(-\frac{\pi}{4}\right) \sin\left(-\frac{\pi}{6}\right) + i \left(\sin\left(-\frac{\pi}{4}\right) \cos\left(-\frac{\pi}{6}\right) - \cos\left(-\frac{\pi}{4}\right) \sin\left(-\frac{\pi}{6}\right) \right) \right]}{2 \times 1} \\
 &= \frac{\sqrt{2} \left[\cos\left(-\frac{\pi}{4} - \left(-\frac{\pi}{6}\right)\right) + i \sin\left(-\frac{\pi}{4} - \left(-\frac{\pi}{6}\right)\right) \right]}{2} \\
 &= \frac{\sqrt{2}}{2} \left[\cos\left(-\frac{\pi}{12}\right) + i \sin\left(-\frac{\pi}{12}\right) \right] = \frac{\sqrt{2}}{2} \operatorname{cis}\left(-\frac{\pi}{12}\right)
 \end{aligned}$$

6. The two results are the same.

Synthesis

Given two complex numbers $z_1 = r_1 (\cos \theta_1 + i \sin \theta_1)$

and $z_2 = r_2 (\cos \theta_2 + i \sin \theta_2)$ then,

$z_1 z_2 = r_1 r_2 (\cos (\theta_1 + \theta_2) + i \sin (\theta_1 + \theta_2))$ and

$\frac{z_1}{z_2} = \frac{r_1}{r_2} (\cos (\theta_1 - \theta_2) + i \sin (\theta_1 - \theta_2))$ with the provision

that 2π may have to be added to, or subtracted from $\theta_1 + \theta_2$ (or $\theta_1 - \theta_2$) if $\theta_1 + \theta_2$ (or $\theta_1 - \theta_2$) is outside the permitted range of the principal argument $]-\pi, \pi]$.

We note that;

$\operatorname{Arg}(z_1 z_2) = \operatorname{Arg}(z_1) + \operatorname{Arg}(z_2)$ and

$\operatorname{Arg}\left(\frac{z_1}{z_2}\right) = \operatorname{Arg}(z_1) - \operatorname{Arg}(z_2)$

Exercise 1.18 Learner's Book page 45

1. a) $zw = 2\sqrt{2} \operatorname{cis}\left(-\frac{11\pi}{12}\right), \frac{z}{w} = \frac{\sqrt{2}}{2} \operatorname{cis}\left(-\frac{7\pi}{12}\right)$
 - b) $zw = 4 \operatorname{cis}\left(\frac{\pi}{3}\right), \frac{z}{w} = 4 \operatorname{cis}(-\pi)$
 - c) $zw = 2\sqrt{2} \operatorname{cis}\left(\frac{5\pi}{12}\right), \frac{z}{w} = \frac{\sqrt{2}}{2} \operatorname{cis}\left(\frac{\pi}{12}\right)$
 - d) $zw = 4\sqrt{6} \operatorname{cis}\left(\frac{5\pi}{12}\right), \frac{z}{w} = \frac{\sqrt{6}}{3} \operatorname{cis}\left(-\frac{11\pi}{12}\right)$
 - e) $zw = \sqrt{2} \operatorname{cis}\left(\frac{\pi}{12}\right), \frac{z}{w} = \frac{\sqrt{2}}{2} \operatorname{cis}\left(\frac{7\pi}{12}\right)$
 - f) $zw = 2 \operatorname{cis}(\pi), \frac{z}{w} = \operatorname{cis}\left(\frac{\pi}{2}\right)$
2. $\sqrt{2} \operatorname{cis}\left(-\frac{5\pi}{12}\right)$

Lesson 1.19. Powers of complex number in polar form**Learning objectives**

Given a complex number, learners should be able to find its powers and use De Moivre's theorem accurately.

Prerequisites

- ⑦ Putting a complex number in polar form.

Teaching Aids

Exercise book, pen and scientific calculator



Activity 1.19 Learner's Book page 45

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad z = \sqrt{3} + i$$

$$|z| = 2, \quad \arg(z) = \arctan\left(\frac{1}{\sqrt{3}}\right) = \frac{\pi}{6}$$

$$z = 2 \operatorname{cis}\left(\frac{\pi}{6}\right)$$

$$2. \quad z^2 = (\sqrt{3} + i)(\sqrt{3} + i) = (\sqrt{3})^2 + 2i\sqrt{3} + (i)^2 = 3 + 2i\sqrt{3} - 1 = 2 + 2i\sqrt{3}$$

$$|z^2| = \sqrt{4 + 12} = 4, \quad \arg(z^2) = \arctan(\sqrt{3}) = \frac{\pi}{3}$$

$$z^2 = 4 \operatorname{cis}\left(\frac{\pi}{3}\right)$$

$$3. \quad z^3 = (2 + 2i\sqrt{3})(\sqrt{3} + i) = 2\sqrt{3} + 2i + 6i - 2\sqrt{3} = 8i$$

$$|z^3| = \sqrt{8^2} = 8, \quad \arg(z^3) = \frac{\pi}{2}$$

$$z^3 = 8 \operatorname{cis}\left(\frac{\pi}{2}\right)$$

4. From 1 to 3, we see that

$$z^2 = 4 \operatorname{cis}\left(\frac{\pi}{3}\right) = 2^2 \operatorname{cis}\left(\frac{2\pi}{6}\right)$$

$$z^3 = 8 \operatorname{cis}\left(\frac{\pi}{2}\right) = 2^3 \operatorname{cis}\left(\frac{3\pi}{6}\right)$$

Hence, $z^n = r^n \operatorname{cis}(n\theta)$ where r is modulus and θ is argument of z .

Synthesis

The power of a complex number in polar form is given

$$\text{by; } z^n = (r(\cos \theta + i \sin \theta))^n = r^n (\cos n\theta + i \sin n\theta); n \in \mathbb{Z}_0$$

where r and θ are modulus and argument of z respectively.

Exercise 1.19 Learner's Book page 46

1. a) $\frac{1}{2}(-1 - i\sqrt{3})$ b) -64 c) $-32i$
 d) $\frac{1}{2}(-1 - i\sqrt{3})$ e) -1 f) -512
 g) $-128 - 128i$
2. $m = 6k, k = 1, 2, 3, 4, \dots$

Lesson 1.20. n^{th} root of a complex number**Learning objectives**

Given a complex number, learners should be able to find the n^{th} roots of that complex number accurately.

Prerequisites

- ✔ Putting a complex number in polar form.
- ✔ Evaluating powers in polar form.

Teaching Aids

Exercise book, pen and scientific calculator

Activity 1.20 Learner's Book page 46

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. |z| = 4, \arg(z) = \arctan(0) = 0 \Rightarrow z = 4 \operatorname{cis} 0$$

$$2. (z_k)^4 = z$$

$$\text{But } (z_k)^4 = (r')^4 \operatorname{cis} 4\theta'$$

$$\text{Then } (r')^4 \operatorname{cis} 4\theta' = 4 \operatorname{cis} 0$$

$$\begin{cases} (r')^4 = 4 \\ 4\theta' = 2k\pi \end{cases} \Rightarrow \begin{cases} r' = \sqrt[4]{4} \Rightarrow r' = \sqrt{2} \\ \theta' = \frac{k\pi}{2} \end{cases}$$

$$\text{Now, } z_k = \sqrt{2} \operatorname{cis} \left(\frac{k\pi}{2} \right)$$

$$\text{If } k = 0, z_0 = \sqrt{2} \operatorname{cis} 0$$

$$\text{If } k = 1, z_1 = \sqrt{2} \operatorname{cis} \left(\frac{\pi}{2} \right)$$

$$\text{If } k = 2, z_2 = \sqrt{2} \operatorname{cis} \pi$$

$$\text{If } k = 3, z_3 = \sqrt{2} \operatorname{cis} \left(\frac{3\pi}{2} \right) = \sqrt{2} \operatorname{cis} \left(-\frac{\pi}{2} \right)$$

Synthesis

To find n^{th} roots of a complex number z , you start by expressing z in polar form $z = r \operatorname{cis} \theta$, where r is modulus of z and θ argument of z .

Then, n^{th} roots of a complex number z is given by

$$z_k = \sqrt[n]{r} \operatorname{cis} \left(\frac{\theta + 2k\pi}{n} \right) \quad k = 0, 1, 2, 3, \dots, n-1$$

Exercise 1.20 Learner's Book page 50

$$1. z_0 = \operatorname{cis} \left(\frac{\pi}{8} \right), z_1 = \operatorname{cis} \left(\frac{5\pi}{8} \right), z_2 = \operatorname{cis} \left(-\frac{7\pi}{8} \right), z_3 = \operatorname{cis} \left(-\frac{3\pi}{8} \right)$$

$$2. z_0 = 1, z_1 = \operatorname{cis} \left(\frac{2\pi}{5} \right), z_2 = \operatorname{cis} \left(\frac{4\pi}{5} \right), z_3 = \operatorname{cis} \left(-\frac{4\pi}{5} \right), z_4 = \operatorname{cis} \left(-\frac{2\pi}{5} \right)$$

$$3. \quad z_0 = 2, \quad z_1 = 2 \operatorname{cis}\left(\frac{2\pi}{5}\right), \quad z_2 = 2 \operatorname{cis}\left(\frac{4\pi}{5}\right), \quad z_3 = 2 \operatorname{cis}\left(-\frac{4\pi}{5}\right), \quad z_4 = 2 \operatorname{cis}\left(-\frac{2\pi}{5}\right)$$

$$4. \quad z_0 = 2 \operatorname{cis}\left(\frac{\pi}{12}\right), \quad z_1 = 2 \operatorname{cis}\left(\frac{7\pi}{12}\right), \quad z_2 = 2 \operatorname{cis}\left(-\frac{11\pi}{12}\right), \quad z_3 = 2 \operatorname{cis}\left(-\frac{5\pi}{12}\right)$$

$$5. \quad \sin \frac{2\pi}{5} = \frac{\sqrt{10+2\sqrt{5}}}{4}$$

Hint:

First, find $\cos \frac{2\pi}{5}$ and then use the relation

$$\sin \frac{2\pi}{5} = \sqrt{1 - \cos^2 \frac{2\pi}{5}}$$

Lesson 1.21. Graphical representation of n th roots of a complex number

Learning objectives

Given a complex number and using a ruler, learners should be able to represent n^{th} roots of that complex number in Argand plane correctly.

Prerequisites

⑦ Finding n^{th} roots of a complex number.

Teaching Aids

Exercise book, pencil, instruments of geometry and calculator

Activity 1.21 Learner's Book page 50



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ⑦ Critical thinking
- ⑦ Communication
- ⑦ Self confidence
- ⑦ Cooperation, interpersonal management and life skills
- ⑦ Peace and values education
- ⑦ Inclusive education

Answers


1. $z = 4, \arg(z) = 0$

$$z_k = \sqrt[5]{4} \operatorname{cis} \frac{2k\pi}{5} \quad z_2 = \sqrt[5]{4} \operatorname{cis} \frac{4\pi}{5}$$

$$z_0 = \sqrt[5]{4} \operatorname{cis} 0 = \sqrt[5]{4} \quad z_3 = \sqrt[5]{4} \operatorname{cis} \frac{6\pi}{5}$$

$$z_1 = \sqrt[5]{4} \operatorname{cis} \frac{2\pi}{5} \quad z_4 = \sqrt[5]{4} \operatorname{cis} \frac{8\pi}{5}$$

2. Representation of the obtained roots on Argand diagram and joining the obtained points.




3. See diagram above.

Synthesis

As conclusion, if the complex number for which we are computing the n^{th} roots is $z = r \operatorname{cis} \theta$, the radius of the circle will be $R = \sqrt[n]{r}$ and the first root z_0 corresponding to $k = 0$ will be at an amplitude of $\varphi = \frac{\theta}{n}$. This root will be followed by the $n-1$ remaining roots at equal distances apart.


Exercise 1.21 Learner's Book page 53

$$1. \quad z_0 = 3 \operatorname{cis}\left(\frac{\pi}{3}\right), \quad z_1 = 3 \operatorname{cis}(\pi), \quad z_2 = 3 \operatorname{cis}\left(-\frac{\pi}{3}\right)$$




$$2. \quad z_0 = \sqrt{2} \operatorname{cis}\left(\frac{\pi}{4}\right), \quad z_1 = \sqrt{2} \operatorname{cis}\left(\frac{3\pi}{4}\right), \quad z_2 = \sqrt{2} \operatorname{cis}\left(-\frac{3\pi}{4}\right),$$


$$z_3 = \sqrt{2} \operatorname{cis}\left(-\frac{\pi}{4}\right)$$



3. $z_0 = 2 \operatorname{cis}\left(\frac{\pi}{6}\right)$, $z_1 = 2 \operatorname{cis}\left(\frac{5\pi}{6}\right)$, $z_2 = 2 \operatorname{cis}\left(-\frac{\pi}{2}\right)$



4. $z_0 = \operatorname{cis}\left(\frac{\pi}{4}\right)$, $z_1 = \operatorname{cis}\left(\frac{3\pi}{4}\right)$, $z_2 = \operatorname{cis}\left(-\frac{3\pi}{4}\right)$, $z_3 = \operatorname{cis}\left(-\frac{\pi}{4}\right)$



Lesson 1.22. Construction of regular polygon

Learning objectives

Using n^{th} roots of unity and geometric instruments, learners should be able to construct a regular polygon in Argand plane accurately.

Prerequisites

- ✔ Finding n^{th} roots of unity
- ✔ Representation of n^{th} roots of unity in Argand diagram.

Teaching Aids

Exercise book, pencil, instrument of geometry and calculator

Activity 1.22 Learner's Book page 54



In this lesson, the following generic competence and cross-cutting issues are to be addressed:


- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $z_k = \text{cis} \frac{2k\pi}{3}, k = 0, 1, 2$

$$z_0 = \text{cis} 0 = 1, \quad z_1 = \text{cis} \frac{2\pi}{3}, \quad z_2 = \text{cis} \frac{4\pi}{3}$$

2. Graph



3. See the graph above.

4. The obtained figure is an equilateral triangle.


Synthesis

To draw a regular polygon with n sides follow the following steps:


- ④ Start by drawing a unit circle in Argand diagram. The radius and the centre of this circle will be the radius and centre of the regular polygon.
- ④ Around the circle, place the points with affixes $z_k = \text{cis} \frac{2k\pi}{n}$, $k = 0, 1, 2, \dots, n-1$. Those points are the vertices of the polygon.
- ④ Using a ruler, join the obtained points around the circle.
- ④ The obtained figure is the needed regular polygon.

Exercise 1.22 Learner's Book page 56


1. A regular hexagon (6 sides)




2. A regular heptagon (7 sides)



3. A regular octagon (8 sides)



4. A regular nonagon (9 sides)



Lesson 1.23. Exponential form of a complex number

Learning objectives

Given a complex number, learners should be able to express that complex number in exponential form accurately.

Prerequisites

- ✔ Finding modulus of a complex number.
- ✔ Finding argument of a complex number.

Teaching Aids

Exercise book, pen and calculator



Activity 1.23 Learner's Book page 56

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \frac{x^7}{7!} + \dots$$

Replacing x with $i\theta$ gives

$$\begin{aligned} e^{i\theta} &= 1 + i\theta + \frac{(i\theta)^2}{2!} + \frac{(i\theta)^3}{3!} + \frac{(i\theta)^4}{4!} + \frac{(i\theta)^5}{5!} + \frac{(i\theta)^6}{6!} + \frac{(i\theta)^7}{7!} + \dots \\ &= 1 + i\theta - \frac{\theta^2}{2!} - \frac{i\theta^3}{3!} + \frac{\theta^4}{4!} + \frac{i\theta^5}{5!} - \frac{\theta^6}{6!} - \frac{i\theta^7}{7!} + \dots \end{aligned}$$

$$\begin{aligned} 2. \quad e^{i\theta} &= 1 + i\theta - \frac{\theta^2}{2!} - \frac{i\theta^3}{3!} + \frac{\theta^4}{4!} + \frac{i\theta^5}{5!} - \frac{\theta^6}{6!} - \frac{i\theta^7}{7!} + \dots \\ &= 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} - \frac{\theta^6}{6!} + \dots + i \left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} - \frac{\theta^7}{7!} + \dots \right) \end{aligned}$$

Since $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} \dots$ and

$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} \dots$, we can write

$$e^{i\theta} = \cos \theta + i \sin \theta$$

3. The right hand side of the expression obtained in 2) is the polar form of complex number having modulus 1 and argument θ .

Synthesis

Exponential form of a complex number z , can be simply found from its polar form $z = r \operatorname{cis} \theta$.

For a complex number having modulus 1 and argument θ , we have the following equality; $e^{i\theta} = \cos \theta + i \sin \theta$, which leads to $r(\cos \theta + i \sin \theta) = r e^{i\theta}$.

Therefore, $z = r e^{i\theta}$ is exponential form of complex number $z = r(\cos \theta + i \sin \theta)$.

Exercise 1.23 Learner's Book page 58

- | | | |
|-----------------------------------|---------------------------|-----------------------------------|
| 1. $e^{\frac{i\pi}{2}}$ | 2. $2e^{\frac{2i\pi}{3}}$ | 3. $2\sqrt{2}e^{-\frac{i\pi}{4}}$ |
| 4. $2\sqrt{3}e^{-\frac{i\pi}{6}}$ | 5. $5e^{i\pi}$ | 6. $3\sqrt{2}e^{\frac{i\pi}{4}}$ |
| 7. $5e^{0.9i}$ | 8. $13e^{-1.9i}$ | |

Lesson 1.24. Trigonometric number of a multiple of an angle

Learning objectives

Given a multiple of an angle, learners should be able to find its trigonometric number accurately.

Prerequisites

- ① Binomial expansion.

Teaching Aids

Exercise book, pen and calculator



Activity 1.24 Learner's Book page 58

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Cooperation, interpersonal management and life skills
- ✔ Self confidence
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Newton binomial expansion gives

$$(\cos x + i \sin x)^n = {}^n C_0 \cos^n x + {}^n C_1 \cos^{n-1} x (i \sin x) + {}^n C_2 \cos^{n-2} x (i \sin x)^2 + \dots + {}^n C_n (i \sin x)^n \quad (1)$$

$$= {}^n C_0 \cos^n x + {}^n C_1 i \cos^{n-1} x \sin x - {}^n C_2 \cos^{n-2} x \sin^2 x + \dots + {}^n C_n i^n \sin^n x \quad (2)$$

2. Relations (1) and (2) are equivalent. Then,

$$\begin{aligned} \cos nx + i \sin nx &= {}^n C_0 \cos^n x - {}^n C_2 \cos^{n-2} x \sin^2 x + \dots \\ &+ i \left({}^n C_1 \cos^{n-1} x \sin x - {}^n C_3 \cos^{n-3} x \sin^3 x + \dots \right) \quad (3) \end{aligned}$$

3. Recall that two complex numbers are equal if they have the same real parts and same imaginary parts.

Thus, from (3), we have

$$\cos nx = {}^n C_0 \cos^n x - {}^n C_2 \cos^{n-2} x \sin^2 x + {}^n C_4 \cos^{n-4} x \sin^4 x + \dots$$

$$\sin nx = {}^n C_1 \cos^{n-1} x \sin x - {}^n C_3 \cos^{n-3} x \sin^3 x + {}^n C_5 \cos^{n-5} x \sin^5 x + \dots$$

Synthesis

Generally,

$$\cos nx = \sum_{0 \leq k \leq n} {}^n C_k i^k \cos^{n-k} x \sin^k x, \text{ with } k - \text{even}$$

$$i \sin nx = \sum_{1 \leq k \leq n} {}^n C_k i^k \cos^{n-k} x \sin^k x, \text{ with } k - \text{odd}$$

$${}^n C_k = \frac{n!}{k!(n-k)!}$$

Exercise 1.24 Learner's Book page 61

1. $2 \cos^2 x - 1$
2. $2 \cos x \sin x$
3. $\frac{3 \cot x - \cot^3 x}{1 - 3 \cot^2 x}$
4. $\frac{3 \cot x - \cot^3 x}{1 - 3 \cot^2 x}$
5. $\frac{\tan^5 x - 10 \tan^3 x + 5 \tan x}{5 \tan^4 x - 10 \tan^2 x + 1}$
6. $32 \cos^6 x - 48 \cos^4 x + 18 \cos^2 x - 1$
7. $-32 \cos^3 x \sin x + 32 \sin x \cos^5 x + 6 \sin x \cos x$

Lesson 1.25. Linearisation of trigonometric expressions**Learning objectives**

Given a trigonometric expression, learners should be able to linearise that trigonometric expression exactly.

Prerequisites

- ✔ Euler's formulae.

Teaching Aids

Exercise book, pen

Activity 1.25 Learner's Book page 61

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

Euler's formulae are

$$\cos x = \frac{1}{2}(e^{ix} + e^{-ix})$$

$$\sin x = \frac{1}{2i}(e^{ix} - e^{-ix})$$

From these formulae, we have

$$\begin{aligned}
 \sin^2 x \cos x &= \left(\frac{e^{ix} - e^{-ix}}{2i} \right)^2 \left(\frac{e^{ix} + e^{-ix}}{2} \right) \\
 &= \left(\frac{e^{2ix} + e^{-2ix} - 2}{-4} \right) \left(\frac{e^{ix} + e^{-ix}}{2} \right) \\
 &= \frac{e^{3ix} + e^{ix} + e^{-ix} + e^{-3ix} - 2e^{ix} - 2e^{-ix}}{-8} \\
 &= \frac{e^{3ix} + e^{-3ix} - e^{ix} - e^{-ix}}{-8} \\
 &= -\frac{1}{4} \left(\frac{e^{3ix} + e^{-3ix}}{2} - \frac{e^{ix} + e^{-ix}}{2} \right) \\
 &= -\frac{1}{4} (\cos 3x + \cos x)
 \end{aligned}$$

Synthesis

To linearise trigonometric expression (product in sum), we use Euler's formulae

$$\cos \theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta}) \quad \sin \theta = \frac{1}{2}(e^{i\theta} - e^{-i\theta})$$

Exercise 1.25 Learner's Book page 62

1. $\frac{1}{2} \cos(x-y) + \frac{1}{2} \cos(x+y)$
2. $\frac{1}{2} \cos(x-y) - \frac{1}{2} \cos(x+y)$
3. $\frac{1}{2} \sin 2x$
4. $\frac{1}{2} - \frac{1}{2} \cos 2x$
5. $\frac{1}{2} + \frac{1}{2} \cos 2x$
6. $\frac{3}{4} \sin x - \frac{1}{4} \sin 3x$
7. $\frac{3}{4} \cos x + \frac{1}{4} \cos 3x$
8. $\frac{1}{8} - \frac{1}{8} \cos 4x$

Lesson 1.26. Solving equation of the form

$$a \cos x + b \sin x = c, \quad a, b, c \in \mathbb{R} \text{ and } a \cdot b \neq 0$$

Learning objectives

Given equation of the form

$a \cos x + b \sin x = c$, $a, b, c \in \mathbb{R}$ and $a \cdot b \neq 0$, learners should be able to solve it using complex numbers precisely.

Prerequisites

- ✔ Putting a complex number in polar form.
- ✔ Solving simple trigonometric equation.

Teaching Aids

Exercise book, pen and scientific calculator.

Activity 1.26 Learner's Book page 63

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad z = 1 - i\sqrt{3} \qquad 2. \quad |z| = \sqrt{1+3} = 2$$

$$3. \quad \arg(z) = \arctan(-\sqrt{3}) = -\frac{\pi}{3}$$

$$4. \quad 2 \cos\left(x + \frac{\pi}{3}\right) = -1$$

$$\Leftrightarrow \cos\left(x + \frac{\pi}{3}\right) = -\frac{1}{2} \Leftrightarrow x + \frac{\pi}{3} = \begin{cases} \frac{2\pi}{3} + 2k\pi \\ -\frac{2\pi}{3} + 2k\pi \end{cases}$$

$$\Rightarrow x = \begin{cases} \frac{\pi}{3} + 2k\pi \\ -\pi + 2k\pi \end{cases}$$

Synthesis

To solve the equation of the form

$a \cos x + b \sin x = c$, $a, b, c \in \mathbb{R}$ and $a \cdot b \neq 0$, follow these steps:

1. Reduction of $a \cos x + b \sin x$ $a, b \in \mathbb{R}$




Figure 1.7: Reduction of a trigonometric expression

To get the expression equivalent to $a \cos x + b \sin x$, we use dot product expressed in terms of angle $\theta - x$ that is, between two vectors $\overrightarrow{OM} = (\cos x, \sin x)$ and $\overrightarrow{ON} = (a, b)$.

$$\text{Or } \cos(\theta - x) = \frac{\overrightarrow{OM} \cdot \overrightarrow{ON}}{|\overrightarrow{OM}| \cdot |\overrightarrow{ON}|}$$

$$\Leftrightarrow \cos(\theta - x) = \frac{a \cos x + b \sin x}{\sqrt{a^2 + b^2} \cdot \sqrt{\cos^2 x + \sin^2 x}}$$

$$\Leftrightarrow \cos(\theta - x) = \frac{a \cos x + b \sin x}{\sqrt{a^2 + b^2}}$$

$$\Rightarrow a \cos x + b \sin x = \sqrt{a^2 + b^2} \cos(\theta - x)$$

Therefore, $a \cos x + b \sin x = c$

$$\Leftrightarrow \sqrt{a^2 + b^2} \cos(\theta - x) = c$$

2. Solve reduction formula of $a \cos x + b \sin x = c$

$$a \cos x + b \sin x = c \Leftrightarrow \sqrt{a^2 + b^2} \cos(\theta - x) = c$$

$$\Leftrightarrow \cos(\theta - x) = \frac{c}{\sqrt{a^2 + b^2}}, \text{ as } \sqrt{a^2 + b^2} \neq 0$$

Since $\forall \alpha \in \mathbb{R}, -1 \leq \cos \alpha \leq 1 \Leftrightarrow |\cos \alpha| \leq 1$, thus,

$a \cos x + b \sin x = c$ has many solutions if and only if

$\left| \frac{c}{\sqrt{a^2 + b^2}} \right| \leq 1$ or $|c| \leq \sqrt{a^2 + b^2}$, otherwise, there is no solution.

Exercise 1.26 Learner's Book page 65

1. $\left\{ x = \frac{\pi}{6} + k\pi, x = \frac{\pi}{2}, k \in \mathbb{Z} \right\}$
2. $\left\{ x = \frac{\pi}{4} + k\pi, k \in \mathbb{Z} \right\}$
3. $\pm \frac{3\pi}{4} - \frac{\pi}{4} + 2k\pi, k \in \mathbb{Z}$
4. $\frac{\pi}{6} \pm \frac{\pi}{4} + 2k\pi, k \in \mathbb{Z}$
5. $\frac{\pi}{6} \pm \frac{\pi}{3} + 2k\pi, k \in \mathbb{Z}$
6. $-\frac{\pi}{4} + 2k\pi, k \in \mathbb{Z}$

Lesson 1.27. Alternating current problem

Learning objectives

By leading textbooks or accessing internet, learners should be able to solve alternating current problems that involve complex numbers accurately.

Prerequisites

- ✔ Converting a complex number to different forms.
- ✔ Make a research by reading textbooks or accessing internet.

Teaching Aids

Exercise book, pen, textbooks or internet if available.

Activity 1.27 Learner's Book page 65

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence



- ④ Cooperation, interpersonal management and life skills
- ④ Research
- ④ Peace and values education
- ④ Inclusive education

Answers

$$Z = R + j\omega L + \frac{1}{j\omega C}$$

when $R = 10, L = 5, C = 0.04$ and $\omega = 4$ we have

$$\begin{aligned} Z &= 10 + j(4)(5) + \frac{1}{j(4)(0.04)} \\ &= 10 + j20 + \frac{1}{0.16j} = 10 + j20 + \frac{-j0.16}{(j0.16)(-j0.16)} \\ &= 10 + j20 - \frac{j0.16}{0.0256} = 10 + j20 - j6.25 = 10 + j13.75 \end{aligned}$$

Thus, $Z = 10 + j13.75$ or $Z = 10 + 13.75j$

Synthesis



The voltage in an AC circuit can be represented as

$$\begin{aligned} V &= V_0 e^{j\omega t} \\ &= V_0 (\cos \omega t + j \sin \omega t) \end{aligned}$$

which denotes **Impedance**, V_0 is peak value of impedance and $\omega = 2\pi f$ where f is the frequency of supply. To obtain the measurable quantity, the real part is taken:

$\text{Re}(V) = V_0 \cos \omega t$ and is called **Resistance** while imaginary part denotes **Reactance** (inductive or capacitive).

Briefly, the current, I (cosine function) leads the applied potential difference (p.d.), V (sine function) by one quarter of a cycle i.e. $\frac{\pi}{2}$ radians or 90° .

Figure 1.8: $R-L$ series circuitFigure 1.9: $R-C$ series circuit

In the **Resistance and Inductance** ($R-L$) series circuit, as shown in figure 1.8,

$V_R + jV_L = V$ as $V_R = IR$ $V_L = IX_L$ (where X_L is the inductive reactance $2\pi fL$ ohms) and $V = IZ$ (where Z is the impedance), then, $R + jX_L = Z$.

In the **Resistance and Capacitance** ($R-C$) circuit, as shown in figure 1.9,

$V_R - jV_C = V$, from which $R - jX_C = Z$

(where X_C is the capacitive reactance, $X_C = \frac{1}{2\pi fC}$ Ω).

Exercise 1.27 Learner's Book page 69

1. a) $R = 3\Omega, L = 25.5mH$ b) $R = 2\Omega, L = 1061\mu F$
 c) $R = 0, L = 44.56mH$ d) $R = 4\Omega, L = 459.5\mu F$
2. $[15.76A, 23.20^\circ \text{ lagging}]$
3. $[27.25A, 3.37^\circ \text{ lagging}]$

- | | |
|-----------------------------------------------------------------------------------|--------------------------------|
| 4. a) $0.3A$ | b) V leads I by 52° |
| 5. $Z_0 = 390.2 \text{cis}(-10.43^\circ), \gamma = 0.1029 \text{cis} 61.92^\circ$ | |

Summary of the unit

1. Concepts of complex numbers

A **complex number** is a number that can be put in the form $a + bi$, where a and b are real numbers and $i = \sqrt{-1}$.

The set of all complex numbers is denoted by \mathbb{C} and is defined as $\mathbb{C} = \{z = a + bi : (a, b) \in \mathbb{R}^2 \text{ and } i^2 = -1\}$.

The real number a of the complex number $z = a + bi$ is called the **real part** of z , and the real number b is often called the **imaginary part**. A complex number whose real part is zero is said to be **purely imaginary**, whereas a complex number whose imaginary part zero is said to be a **real number** or simply **real**.

2. Algebraic form of a complex number

Powers of i : $i^{4k} = 1, i^{4k+1} = i, i^{4k+2} = -1, i^{4k+3} = -i$

$z = (a, b)$ is a **geometric** form of the complex number z .

$z = a + bi$ is the **algebraic** (or **standard** or **Cartesian** or **rectangular**) form of the complex number z .

If two complex numbers, say $a + bi$ and $c + di$ are equal then, both their real and imaginary parts are equal. That is,
 $a + bi = c + di \Leftrightarrow a = c \text{ and } b = d$.

The **addition and subtraction** of two complex numbers $a + bi$ and $c + di$ is defined by the formula:

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i$$

The **complex conjugate** of the complex number $z = x + yi$, denoted by \bar{z} or z^* , is defined to be $\bar{z} = x - yi$.

The complex number $-z = -x - yi$ is the **opposite** of $z = x + yi$, symmetric of z with respect to 0.

The **multiplication** of two complex numbers $c + di$ and $c + di$ is defined by the formula:

$$(a + bi)(c + di) = (ac - bd) + (bc + ad)i$$

The **inverse** of $z = a + bi$ is given by $\frac{1}{z} = z^{-1} = \frac{\bar{z}}{a^2 + b^2}$

If $z_1 = a + bi$ and $z_2 = c + di$ then,

$$\frac{z_1}{z_2} = \frac{a + bi}{c + di} = \left(\frac{ac + bd}{c^2 + d^2} \right) + \left(\frac{bc - ad}{c^2 + d^2} \right) i$$

If a complex number $x + yi$ is a square root of the complex

$$\text{number } a + bi, \text{ then, } \begin{cases} x = \pm \sqrt{\frac{1}{2} \left(a + \sqrt{a^2 + b^2} \right)} \\ y = \pm \sqrt{\frac{1}{2} \left(\sqrt{a^2 + b^2} - a \right)} \end{cases}$$

Let a , b and c be real numbers ($a \neq 0$), then the equation $az^2 + bz + c = 0$ has either two real roots, one double real root or two conjugate complex roots.

a) If $\Delta > 0$, there are two distinct real roots:

$$z_1 = \frac{-b + \sqrt{\Delta}}{2a} \text{ and } z_2 = \frac{-b - \sqrt{\Delta}}{2a}.$$

b) If $\Delta = 0$, there is a double real root:

$$z_1 = z_2 = -\frac{b}{2a}$$

c) If $\Delta < 0$, there is no real roots. In this case, there are two conjugate complex roots:

$$z_1 = \frac{-b + i\sqrt{-\Delta}}{2a} \text{ and } z_2 = \frac{-b - i\sqrt{-\Delta}}{2a}.$$

Where, $\Delta = b^2 - 4ac$

$$z_1 + z_2 = -\frac{b}{a}, \quad z_1 \cdot z_2 = \frac{c}{a}$$

Every polynomial of positive degree with coefficients in the system of complex numbers has a zero in the system of complex numbers. Moreover, every such polynomial can be factored linearly in the system of complex numbers.

3. Polar form of a complex number

The **absolute value (or modulus or magnitude)** of a complex number $z = x + yi$ is $r = |z| = \sqrt{x^2 + y^2}$

Principal argument of a complex number $z = x + yi$

$$\arg(z) = \begin{cases} \arctan \frac{y}{x}, & \text{if } x > 0 \\ \pi + \arctan \frac{y}{x}, & \text{if } x < 0, y \geq 0 \\ -\pi + \arctan \frac{y}{x}, & \text{if } x < 0, y < 0 \\ \frac{\pi}{2}, & \text{if } x = 0, y > 0 \\ -\frac{\pi}{2}, & \text{if } x = 0, y < 0 \\ \text{Undefined} & \text{if } x = 0 \text{ and } y = 0 \end{cases}$$

Polar (or modulus-argument) form is $z = r(\cos \theta + i \sin \theta)$ or $z = r \operatorname{cis} \theta$.

Given two complex numbers $z_1 = r_1(\cos \theta_1 + i \sin \theta_1)$ and $z_2 = r_2(\cos \theta_2 + i \sin \theta_2)$, the formulae for multiplication and division are $z_1 z_2 = r_1 r_2 (\cos (\theta_1 + \theta_2) + i \sin (\theta_1 + \theta_2))$ and $\frac{z_1}{z_2} = \frac{r_1}{r_2} (\cos (\theta_1 - \theta_2) + i \sin (\theta_1 - \theta_2))$ respectively.

Power of a complex number z is given by

$$z^n = (r(\cos \theta + i \sin \theta))^n = r^n (\cos n\theta + i \sin n\theta); n \in \mathbb{Z}_0$$

De Moivre's theorem: $(\cos \theta + i \sin \theta)^n = (\cos n\theta + i \sin n\theta)$

If $(z_k)^n = z$ for $z = r \operatorname{cis} \theta$, then

$$z_k = \sqrt[n]{r} \operatorname{cis} \left(\frac{\theta + 2k\pi}{n} \right) \quad k = 0, 1, 2, 3, \dots, n-1$$

To draw a regular polygon with n sides, the steps followed are:

- ▶ Start by drawing a unit circle in Argand diagram. The radius and the centre of this circle will be the radius and centre of the regular polygon.
- ▶ Around the circle, place the points with affixes $z_k = \text{cis} \frac{2k\pi}{n}$, $k = 0, 1, 2, \dots, n-1$. Those points are the vertices of the polygon.
- ▶ Using a ruler, join the obtained points around the circle.
- ▶ The obtained figure is the needed regular polygon.

4. Exponential form of a complex number

The exponential form of a complex number z whose modulus is r and argument is θ , is $z = re^{i\theta}$

Euler's formulae (these formulae are used to linearise trigonometric expressions):

$$\cos \theta = \frac{1}{2}(e^{i\theta} + e^{-i\theta})$$

$$\sin \theta = \frac{1}{2i}(e^{i\theta} - e^{-i\theta})$$

5. Applications

Formulae for trigonometric number of a multiple of an angle

$$\cos nx = \sum_{0 \leq k \leq n} C_k i^k \cos^{n-k} x \sin^k, \text{ with } k \text{ even}$$

$$i \sin nx = \sum_{1 \leq k \leq n} C_k i^k \cos^{n-k} x \sin^k, \text{ with } k \text{ odd}$$

$${}^n C_k = \frac{n!}{k!(n-k)!}$$


- ▶ To solve the equation $a \cos x + b \sin x = c$, solve the equation

$$\cos(x - \theta) = \frac{c}{\sqrt{a^2 + b^2}}, \quad \theta = \arg(a + bi)$$

Alternating current

Resistance and Capacitance (R-C)

Let a p.d. V be applied across a resistance R and a capacitance C in series. The same current I flows through each component and so the reference vector will be that representing I . The p.d. V_R across R is in phase with I , and V_C , that across C , lags on current I by 90° .



Phasor diagram




Figure showing Resistance and Capacitance in series


Vector sum of V_R and V_C is called Impedance and equals the applied p.d. V ;

$Z = V_R + jV_C$ where V_R and V_C are known as resistance and reactance respectively.

But $V_R = IR$ and $V_C = IX_C$ where X_C is the capacitive reactance of C and equals $\frac{1}{\omega C}$.

Resistance and inductance (R-L)

The analysis is similar but here, the p.d. V_L across L leads on current I and the p.d. V_R across R is again in phase with I .



Phasor diagram




Figure showing Resistance and Inductance in series

$Z = V_R + jV_L$ where V_R and V_L are known as resistance and reactance respectively.

But $V_R = IR$ and $V_L = IX_L$ where X_L is the inductive reactance of L and equals ωL

or $\omega = 2\pi f$.

For the n-branch parallel circuit, Impedance Z is given by:

$$\frac{1}{Z} = \sum_{k=1}^n \frac{1}{Z_k}$$

End of Unit Assessment answers Learner's Book page 75

1. a) $3-8i$ b) $5+5i$

c) $100+200i$ d) $\frac{1-7i}{5}$


2. a) $x = \pm\sqrt{-4} = \pm 2\sqrt{-1} = \pm 2i$

b) $x = \frac{-1 \pm i\sqrt{3}}{2} = -\frac{1}{2} \pm i\frac{\sqrt{3}}{2}$

c) $x = \frac{-6 \pm \sqrt{36-44}}{2} = \frac{-6 \pm i2\sqrt{2}}{2} = -3 \pm i\sqrt{2}$

d) $x = 1, -\frac{1}{2} \pm i\frac{\sqrt{3}}{2}$

3. Plot



a) $1 = cis\ 0$ b) $i = cis\ \frac{\pi}{2}$

c) $-3i = 3cis\left(-\frac{\pi}{2}\right)$ d) $1-i = \sqrt{2}cis\left(-\frac{\pi}{4}\right)$

e) $2+i = \sqrt{5}cis\left(\arctan\left(\frac{1}{2}\right)\right)$

f) $-3-2i = \sqrt{13}cis\left(\pi - \arctan\left(\frac{2}{3}\right)\right)$

g) $-3+2i = \sqrt{13}cis\left(\pi + \arctan\left(-\frac{2}{3}\right)\right)$

4. a) $2 \operatorname{cis} 0 = 2$ b) $3 \operatorname{cis} \pi = -3$
 c) $\operatorname{cis} \frac{\pi}{2} = i$ d) $3 \operatorname{cis} \frac{3\pi}{4} = -\frac{3\sqrt{2}}{2} + i \frac{3\sqrt{2}}{2}$
5. a) $i^7 = -i$ b) $(1+i)^5 = -4(1+i)$
 c) $(\sqrt{3}-i)^{-4} = 2^{-4} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)$
6. a) $z_0 = \operatorname{cis} \left(\frac{\pi}{6} \right) = \frac{\sqrt{3}}{2} + i \frac{1}{2}$, $z_1 = \operatorname{cis} \left(\frac{5\pi}{6} \right) = -\frac{\sqrt{3}}{2} + i \frac{1}{2}$,
 $z_2 = \operatorname{cis} \left(\frac{3\pi}{2} \right) = -i$
 b) $z_0 = \sqrt[8]{2} \operatorname{cis} \left(\frac{\pi}{16} \right)$, $z_1 = \sqrt[8]{2} \operatorname{cis} \left(\frac{9\pi}{16} \right)$, $z_2 = \sqrt[8]{2} \operatorname{cis} \left(\frac{17\pi}{16} \right)$,
 $z_3 = \sqrt[8]{2} \operatorname{cis} \left(\frac{25\pi}{16} \right)$
7. Polar form: $\frac{z_1}{z_2} = \operatorname{cis} \frac{7\pi}{12}$, Cartesian form:

$$\frac{z_1}{z_2} = \frac{\sqrt{2}}{4} - \frac{\sqrt{6}}{4} + i \left(\frac{\sqrt{2}}{4} + \frac{\sqrt{6}}{4} \right)$$

$$\begin{cases} \cos \frac{7\pi}{12} = \frac{\sqrt{2} - \sqrt{6}}{4} \\ \sin \frac{7\pi}{12} = \frac{\sqrt{2} + \sqrt{6}}{4} \end{cases}$$
8. $|e^{i\theta}| = |\cos \theta + i \sin \theta| = \sqrt{\cos^2 \theta + \sin^2 \theta} = 1$
9. $(e^{i\theta})^{-1} = (\cos \theta + i \sin \theta)^{-1}$

$$= \frac{1}{\cos \theta + i \sin \theta} = \frac{\cos \theta - i \sin \theta}{\cos^2 \theta + \sin^2 \theta} = \cos \theta - i \sin \theta = \cos(-\theta) + i \sin(-\theta) = e^{-i\theta}$$
10. The n^{th} roots of unit are given by:

$$z_k = \operatorname{cis} \frac{2k\pi}{n} \quad k = 0, 1, 2, 3, \dots, n-1$$

$$\begin{aligned}
 z_0 &= cis\ 0 = 1 & z_1 &= cis\left(\frac{2\pi}{n}\right) \\
 z_2 &= cis\left(\frac{4\pi}{n}\right) = \left(cis\left(\frac{2\pi}{n}\right)\right)^2 = z_1^2 \\
 z_3 &= cis\left(\frac{6\pi}{n}\right) = \left(cis\left(\frac{2\pi}{n}\right)\right)^3 = z_1^3 \\
 &\vdots \\
 z_{n-1} &= cis\left(\frac{2(n-1)\pi}{n}\right) = \left(cis\left(\frac{2\pi}{n}\right)\right)^{n-1} = z_1^{n-1}
 \end{aligned}$$

The sum is $s_n = 1 + z_1 + z_1^2 + z_1^3 + \dots + z_1^{n-1}$ (1)

Multiplying both sides by z_1 gives

$$\begin{aligned}
 z_1 s_n &= z_1 (1 + z_1 + z_1^2 + z_1^3 + \dots + z_1^{n-1}) \\
 \Leftrightarrow z_1 s_n &= z_1 + z_1^2 + z_1^3 + z_1^4 + \dots + z_1^n \quad (2)
 \end{aligned}$$

(1) – (2) gives

$$\begin{aligned}
 (s_n - z_1 s_n) &= 1 + z_1 + z_1^2 + z_1^3 + \dots + z_1^{n-1} \\
 &\quad \underline{- z_1 - z_1^2 - z_1^3 - z_1^4 - \dots - z_1^n}
 \end{aligned}$$

$$s_n (1 - z_1) = 1 - z_1^n$$

$$\Leftrightarrow s_n = \frac{1 - z_1^n}{1 - z_1}$$

But $z_1^n = cis\left(\frac{2n\pi}{n}\right) = cis(2\pi) = 1$, then

$$s_n = \frac{1 - z_1^n}{1 - z_1} = \frac{1 - 1}{1 - z_1} = \frac{0}{1 - z_1} = 0$$

Thus, the sum of n th roots of unit is zero.

11. $\cos\frac{\pi}{5} = \frac{1 + \sqrt{5}}{4}$
12. $z = -\frac{3}{2} + yi, \quad y \in \mathbb{R}$
13. $S = \{-3, -2i, 2i\}$

$$14. \left(\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)^{2001} = -1$$

15. a) M is a point on circle of diameter $[AB]$ if $\overline{BM} \perp \overline{AM}$.

We need to check if $\frac{z_M - z_B}{z_M - z_A}$ is pure imaginary.

$$\begin{aligned} \frac{z_M - z_B}{z_M - z_A} &= \frac{ie^{i\theta} - i}{ie^{i\theta} + i} = \frac{e^{i\theta} - 1}{e^{i\theta} + 1} \\ &= \frac{e^{i\frac{\theta}{2}} \left(e^{i\frac{\theta}{2}} - e^{-i\frac{\theta}{2}} \right)}{e^{i\frac{\theta}{2}} \left(e^{i\frac{\theta}{2}} + e^{-i\frac{\theta}{2}} \right)} = \frac{2i \sin \frac{\theta}{2}}{2 \cos \frac{\theta}{2}} = i \tan \frac{\theta}{2} \end{aligned}$$

which is a pure imaginary.

Thus, M is a point on circle of diameter $[AB]$.

b) Rotation of centre o and angle of $\frac{\pi}{2}$ is $z' = e^{i\frac{\pi}{2}} z = iz$

$$z_{M'} = i(1 + ie^{i\theta}) = i - e^{i\theta}$$

$$\frac{z_M - z_B}{z_{M'} - z_B} = \frac{ie^{i\theta} - i}{-e^{i\theta} - 1} = \frac{i(e^{i\theta} - 1)}{i^2(e^{i\theta} + 1)} = \frac{(e^{i\theta} - 1)}{i(e^{i\theta} + 1)} = \frac{2i \sin \frac{\theta}{2}}{2i \cos \frac{\theta}{2}} = \tan \frac{\theta}{2}$$

which is real.

Thus, points B, M and M' are collinear.

16. Values of x are 2 and -5

17. a) The locus is the mediator of the segment $[AB]$ such that $z_A = 2$ and $z_B = -1$.

b) The locus is the mediator of the segment $[AB]$ such that $z_A = 2i$ and $z_B = -2$.

c) The locus is the circle of centre $1 - 3i$ and radius 2.

d) The locus is the circle of centre $-1 + 0$ and radius 1.

e) The locus is the rectangular hyperbola.

f) The locus is the union of 2 bisectors of equations $y = -x$ and $y = x$ respectively.

18. The two complex numbers are $1+2i\sqrt{2}$ and $1-2i\sqrt{2}$
19. $z = cis\left(\frac{k\pi}{4}\right), k \in \mathbb{Z}$
20. $\operatorname{Re}(z) = 0, \operatorname{Im}(z) = -\frac{\sqrt{3}}{3}$
21. $|z| = \cot \frac{\theta}{2}, \arg(z) = \theta - \frac{\pi}{2}$
22. $z = \left(\frac{-1+\sqrt{3}}{2}\right) + \left(\frac{-1+\sqrt{3}}{2}\right)i$ or $z = \left(\frac{-1-\sqrt{3}}{2}\right) + \left(\frac{-1-\sqrt{3}}{2}\right)i$
23. $z = 6 + yi, y \in \mathbb{R}$

24. Isosceles triangle has two sides equal in length different from the third. We must check if there are two equal sides among $\|\overline{AB}\|, \|\overline{AC}\|$ and $\|\overline{BC}\|$

$$\|\overline{AB}\| = \|4 - 2i - 1 - 2i\| = \|3 - 4i\| = \sqrt{9+16} = 5$$

$$\|\overline{AC}\| = \|1 - 6i - 1 - 2i\| = \|-8i\| = \sqrt{64} = 8$$

$$\|\overline{BC}\| = \|1 - 6i - 4 + 2i\| = \|-3 - 4i\| = \sqrt{9+16} = 5$$

Then, $\|\overline{AB}\| = \|\overline{BC}\| \neq \|\overline{AC}\|$ and hence the triangle is isosceles.

25. a) (i) $\left(\frac{\sqrt{3}-i}{1+i\sqrt{3}}\right)^9 = -i$ (ii) $\left(\frac{\sqrt{3}+i}{\sqrt{3}-i} + \frac{\sqrt{3}-i}{\sqrt{3}+i} - 2\right)^{30} = 1$

b) (i) $n = 6k, k \in \{1, 2, 3, 4, 5, \dots\}$

(ii) $n = 3m, m \in \{1, 3, 5, 7, 9, \dots\}$

c) Let $z_1 = 1 + i$ $z_2 = 1 - i$

$|z_1| = \sqrt{2}$ $|z_2| = \sqrt{2}$

$\arg(z_1) = \frac{\pi}{4}$ $\arg(z_2) = -\frac{\pi}{4}$

$z_1 = \sqrt{2}cis\left(\frac{\pi}{4}\right)$ $z_2 = \sqrt{2}cis\left(-\frac{\pi}{4}\right)$

Then,


$$\begin{aligned}
 (1+i)^n + (1-i)^n &= z_1^n + z_2^n = \left[\sqrt{2} \operatorname{cis} \left(\frac{\pi}{4} \right) \right]^n + \left[\sqrt{2} \operatorname{cis} \left(-\frac{\pi}{4} \right) \right]^n \\
 &= (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) + i (\sqrt{2})^n \sin \left(\frac{n\pi}{4} \right) + (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) + i (\sqrt{2})^n \sin \left(-\frac{n\pi}{4} \right) \\
 &= (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) + i (\sqrt{2})^n \sin \left(\frac{n\pi}{4} \right) + (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) - i (\sqrt{2})^n \sin \left(\frac{n\pi}{4} \right) \\
 &= (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) + (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) = 2 (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) \\
 &= 2 (\sqrt{2})^n \cos \left(\frac{n\pi}{4} \right) = 2 (2)^{\frac{n}{2}} \cos \left(\frac{n\pi}{4} \right) = 2^{1+\frac{n}{2}} \cos \left(\frac{n\pi}{4} \right) \\
 &= 2^{\frac{n+2}{2}} \cos \left(\frac{n\pi}{4} \right) \text{ as required.}
 \end{aligned}$$

26. a) $E(-1) = (-1)^3 + 2(-1)^2 + 2(-1) + 1 = 0$. Thus, -1 is a root of E .

b) $a = 1, b = 1, c = 1$

c) $S = \left\{ -1, -\frac{1}{2} - i \frac{\sqrt{3}}{2}, -\frac{1}{2} + i \frac{\sqrt{3}}{2} \right\}$

27. a) Complex plane



b) $z_{\overline{AB}} = \frac{1}{2} - 2 + 3i = -\frac{3}{2} + 3i$, $z_{\overline{BC}} = 1 + 4i - \frac{1}{2} = \frac{1}{2} + 4i$

c) $E = \frac{5}{2} + i$

28. a) Polar form: $\frac{z_1}{z_2} = \frac{\sqrt{2}}{2} \text{cis}\left(\frac{5\pi}{12}\right)$

Algebraic form: $\frac{z_1}{z_2} = \frac{\sqrt{3}-1}{4} + i\frac{\sqrt{3}+1}{4}$

b)
$$\begin{cases} \cos \frac{5\pi}{12} = \frac{\sqrt{6}-\sqrt{2}}{4} \\ \sin \frac{5\pi}{12} = \frac{\sqrt{6}+\sqrt{2}}{4} \end{cases}$$

c) The lowest value of n is 12.

29. $11.86N, 146.77^\circ$ from force A

30. $8.394N, 208.68^\circ$ from force A

31. $(10 + j20)\Omega, 22.36\text{cis}63.43^\circ\Omega$

32. $\pm\left(\frac{mh}{2\pi}\right)$

33. $[14.42A, 43.85^\circ \text{lagging}]$

34. $14.58A, 2.51^\circ \text{leading}$

35. Current $I = \frac{V}{Z}$

Impedance Z for three branch parallel circuit is given by

$$\frac{1}{Z} = \frac{1}{Z_1} + \frac{1}{Z_2} + \frac{1}{Z_3}$$


For our case, $Z_1 = 4 + 3j, Z_2 = 10, Z_3 = 12 - 5j$.

And then,

$$\begin{aligned} \frac{1}{Z} &= \frac{1}{4+3j} + \frac{1}{10} + \frac{1}{12-5j} = \frac{2797}{8450} - \frac{764}{8450} = 0.331 - 0.0904j \\ &= 0.343 [\cos(-15^\circ 17') + j \sin(-15^\circ 17')] \end{aligned}$$

$$\begin{aligned} I = \frac{V}{Z} &= 240 \times 0.343 [\cos(-15.28^\circ) + j \sin(-15.28^\circ)] \\ &= 82.32 [\cos(-15.28^\circ) + j \sin(-15.28^\circ)] \end{aligned}$$

Thus, $I = 82.32A$, with $\theta = 15.28^\circ$ lagging.



Logarithmic and Exponential Functions

Learner's Book pages 81 - 141

Key unit competence

Extend the concepts of functions to investigate fully logarithmic and exponential functions, finding the domain of definition, the limits, asymptotes, variations, graphs, and model problems about interest rates, population growth or decay, magnitude of earthquake, etc

Vocabulary or key words concepts

- Depreciation:** A negative growth (diminishing in value over a period of time).
- Earthquake:** A sudden violent shaking of the ground as a result of movements within the earth's crust.
- Richter scale:** A logarithmic scale for expressing the magnitude of an earthquake on the basis of seismograph oscillations.

Guidance on the problem statement

The problem statement is "The population P of a city increases according to the formula $P = 500e^{at}$ where t is in years and $t = 0$ corresponds to 1980. In 1990, the population was 10,000. Find the value of the constant a ; correct your answer to 3 decimal places."

From this problem, if $t = 0$ corresponds to 1980, then 1990 corresponds to $t = 10$ and this gives the following equation:

$$500e^{at} = 1000 \text{ or } e^{at} = 2.$$

To find the value a , we take \ln on both sides and we get

$$\ln e^{10a} = \ln 2 \text{ or } 10a \ln e = \ln 2 \Rightarrow a = \frac{\ln 2}{10} = 0.069. \text{ Such kind of problems are solved using logarithms.}$$

List of lessons

No	Lesson title	Number of periods
1	Domain and range of natural logarithmic function	1
2	Limit and asymptotes for natural logarithmic function	1
3	Derivative of natural logarithmic function	1
4	Variation and curve of natural logarithmic function	2
5	Domain and range of logarithmic function with any base	1
6	Limit and asymptotes for logarithmic function with any base	1
7	Logarithmic differentiation	1
8	Further differentiation	1
9	Variation and curve of logarithmic function with any base	2
10	Domain and range of exponential function with base e	1
11	Limit and asymptotes for exponential function with base e	1
12	Derivative of exponential function with base e	1
13	Variation and curve of exponential function with base e	2
14	Domain and range of exponential function with any base	1

15	Limit and asymptotes for exponential function with any base	1
16	Derivative of exponential function with any base	1
17	Variation and curve of exponential function with any base	2
18	Compound interest problems	2
19	Mortgage amount problems	1
20	Population growth problems	1
21	Depreciation value problems	1
22	Earthquake problems	1
23	Carbon-14 dating	1
Total periods		28

Lesson development

Lesson 2.1. Domain and range of natural logarithmic functions

Learning objectives

Given any logarithmic function, learners should be able to find its domain and range accurately.

Prerequisites

- ✔ Finding domain of polynomial, rational/irrational and trigonometric functions.
- ✔ Finding range of polynomial, rational/irrational and trigonometric functions.

Teaching Aids

Exercise book, pen and calculator

Activity 2.1 Learner's Book page 82



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Self confidence
- ✔ Communication

- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education
- ④ Financial education

Answers

x	$\ln x$
-0.8	impossible
-0.6	impossible
-0.4	impossible
-0.2	impossible
0	impossible

x	$\ln x$
0.2	-1.61
0.4	-0.91
0.6	-0.51
0.8	-0.22
1	0

x	$\ln x$
1.5	0.40
2	0.69
2.5	0.91
3	1.09
3.5	1.25

1. (i) For negative x values and zero, \ln is impossible.
 (ii) For x values between 0 and 1, \ln is less than zero.
 (iii) For x values greater than 1, \ln is greater than zero.
2. Curve




Figure 2.1: Curve of $y = \ln x$

Synthesis

From figure 2.1, $\ln x$ is defined on positive real numbers, $]0, +\infty[$ and its range is all real numbers that is $\text{dom} f =]0, +\infty[$ and $\text{Im} f =]-\infty, +\infty[$.

Exercise 2.1 Learner's Book page 84

1. a) $]0, +\infty[$ b) $]0, 4[$
 c) $] -\infty, +\infty[$ d) $] -\infty, 3[$
 2. 316.2

Lesson 2.2. Limit and asymptotes for natural logarithmic functions**Learning objectives**

Given a natural logarithmic function, learners should be able to evaluate limits and deduce relative asymptotes accurately.

Prerequisites

- ✔ Evaluating limits.
- ✔ Finding relative asymptotes.

Teaching Aids

Exercise book, pen and calculator

Activity 2.2 Learner's Book page 84

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Self confidence
- ✔ Communication
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $\lim_{x \rightarrow 0^+} \ln x$ does not exist since at the left of zero \ln is impossible.

2. If x takes on values closer to 0 from the right, we have

x	$\ln x$
0.5	-0.69315
0.45	-0.79851
0.4	-0.91629
0.35	-1.04982
0.3	-1.20397
0.25	-1.38629
0.2	-1.60944
0.15	-1.89712
0.1	-2.30259
0.05	-2.99573

We see that if x takes on values closer to 0 from the right, $\ln x$ becomes smaller and smaller negative.

Then $\lim_{x \rightarrow 0^+} \ln x = -\infty$. There is a vertical asymptote $x = 0$.

3. If we give to x the values of the form 10^n ($n \in \mathbb{N}$), $\ln 10^n = n \ln 10 \approx 2.30n$ and let n take values 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, ..., we have;

n	$x = 10^n$	$\ln x$	$\frac{\ln x}{x}$
1	10	2.302585	0.230258509
2	100	4.60517	0.046051702
3	1000	6.907755	0.006907755
4	10000	9.21034	0.000921034
5	100000	11.51293	0.000115129
6	1000000	13.81551	0.000013816
7	10000000	16.1181	0.000001612
8	100000000	18.42068	0.000000184
9	1000000000	20.72327	0.000000021
10	10000000000	23.02585	0.000000002

We see that if x takes on values of the form 10^n ($n \in \mathbb{N}$),

$\ln x$ becomes larger and larger without bound and consequently approaches no fixed value. Then

$$\lim_{x \rightarrow +\infty} \ln(x) = +\infty. \text{ There is no horizontal asymptote.}$$

Also, that if x takes the values of the form

$$10^n \ (n \in \mathbb{N}), \ \frac{\ln x}{x} \text{ becomes closer to zero.}$$

$$\text{Then, } \lim_{x \rightarrow +\infty} \frac{\ln x}{x} = 0.$$

There is no oblique asymptote.

Synthesis

As conclusion, $\lim_{x \rightarrow +\infty} \ln x = +\infty$ and $\lim_{x \rightarrow 0^+} \ln x = -\infty$

There exists a vertical asymptote with equation $VA \equiv x = 0$

No horizontal asymptote.

Exercise 2.2 Learner's Book page 86

1) $-\infty$

2) 0

3) $+\infty$

4) $+\infty$

Lesson 2.3. Derivative of natural logarithmic functions

Learning objectives

Given a natural logarithmic function, learners should be able to differentiate it accurately.

Prerequisites

- ✔ Definition of derivative.
- ✔ Differentiating a polynomial, rational/ irrational and trigonometric functions.

Teaching Aids

Exercise book and pen



Activity 2.3 Learner's Book page 86

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. (\ln x)' = \lim_{h \rightarrow 0} \frac{\ln(x+h) - \ln x}{h} = \lim_{h \rightarrow 0} \frac{\ln\left(\frac{x+h}{x}\right)}{h}$$

$$= \lim_{h \rightarrow 0} \frac{1}{h} \ln\left(\frac{x+h}{x}\right) = \lim_{h \rightarrow 0} \ln\left(1 + \frac{h}{x}\right)^{\frac{1}{h}}$$

$$\text{Let } u = \frac{h}{x} \Rightarrow h = ux$$

$$\text{If } h \rightarrow 0, u \rightarrow 0$$

$$\begin{aligned} \lim_{h \rightarrow 0} \ln\left(1 + \frac{h}{x}\right)^{\frac{1}{h}} &= \lim_{u \rightarrow 0} \ln(1+u)^{\frac{1}{ux}} \\ &= \lim_{u \rightarrow 0} \frac{1}{x} \ln(1+u)^{\frac{1}{u}} \\ &= \frac{1}{x} \lim_{u \rightarrow 0} \ln(1+u)^{\frac{1}{u}} \\ &= \frac{1}{x} \ln \lim_{u \rightarrow 0} (1+u)^{\frac{1}{u}} \\ &= \frac{1}{x} \ln e \quad \text{since } \lim_{u \rightarrow 0} (1+u)^{\frac{1}{u}} = e \\ &= \frac{1}{x} \end{aligned}$$

$$\text{Thus, } (\ln x)' = \frac{1}{x}$$

2. $(\ln u)' = \frac{1}{u} u' = \frac{u'}{u}$. Thus, $(\ln u)' = \frac{u'}{u}$ where u is another differentiable function.

Synthesis

$(\ln x)' = \frac{1}{x}$; if u is another differentiable function of x then, $(\ln u)' = \frac{u'}{u}$.

Exercise 2.3 Learner's Book page 87

- | | | |
|--------------------------------------------|----------------------------------------------|----------------------------|
| 1. $\frac{2 \ln x}{x}$ | 2. $\frac{\tan^2 x + 1}{\tan x}$ | 3. $\frac{x}{x^2 - 1}$ |
| 4. $\frac{2}{x^2 - 1}$ | 5. $\frac{x \tan x - \ln(\sin x)}{x^2}$ | 6. $-\tan x + \frac{1}{x}$ |
| 7. $\tan^2 x - \frac{1 - \ln x}{3x^2} + 1$ | 8. $\frac{-2 \ln(\sqrt{x+1}) + 1}{2(x+1)^2}$ | |

Lesson 2.4. Variation and curve sketching of natural logarithmic functions

Learning objectives

Given a natural logarithmic function, learners should be able to study the variation and sketch its curve perfectly.

Prerequisites

- ✔ Finding domain and limits at the boundaries of the domain.
- ✔ Deducing relative asymptotes.
- ✔ Finding first and second derivative.
- ✔ Variation and concavity of a function.
- ✔ Sketch a curve in Cartesian plane given some points.

Teaching Aids

Exercise book, pencil, instrument of geometry and calculator



Activity 2.4 Learner's Book page 87

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $f(x) = \ln x$. The domain is $]0, +\infty[$.
 $\lim_{x \rightarrow 0^+} f(x) = -\infty$. There is a vertical asymptote $x = 0$
 $\lim_{x \rightarrow +\infty} f(x) = +\infty$. There is no horizontal asymptote
 $\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{\ln x}{x} = 0$. There is no oblique asymptote.
2. $f(x) = \ln x \Rightarrow f'(x) = \frac{1}{x}$. Since $x \in]0, +\infty[$, $f'(x)$ is always positive and hence $f(x) = \ln x$ increases on its domain. Since $f'(x) = \frac{1}{x} \neq 0$, there is no extrema (no maximum, no minimum).
3. $f'(x) = \frac{1}{x} \Rightarrow f''(x) = -\frac{1}{x^2}$. $f''(x)$ is always negative and hence the concavity of $f(x) = \ln x$ is turning down on its domain. Since $f''(x) = -\frac{1}{x^2} \neq 0$, there is no inflection points.

4. Completed table of variation

x	0	1		e	$+\infty$
Sign of $f'(x)$	+	+	+	+	+
Sign of $f''(x)$	-	-	-	-	-
Variation of $f(x)$					
Concavity of $f(x)$					

5. Intersection of $f(x)$ with axes of co-ordinates:

There is no intersection of $f(x)$ with y -axis since this axis is an asymptote.

Intersection with x -axis :

$$f(x) = 0 \Leftrightarrow \ln x = 0 \Rightarrow \ln x = \ln 1 \Rightarrow x = 1. \text{ Then,}$$


$$f(x) \cap ox = \{(1, 0)\}.$$

6. Additional points

x	0.1	0.3	0.5	0.7	0.9	1.1	1.3	1.5	1.7	1.9	2.1
y	-2.3	-1.2	-0.7	-0.4	-0.1	0.1	0.3	0.4	0.5	0.6	0.7

x	2.3	2.5	2.7	2.9	3.1	3.3	3.5	3.7	3.9
y	0.8	0.9	1.0	1.1	1.1	1.2	1.3	1.3	1.4

Graph



Synthesis

To sketch a function, follow the following steps:

- ① Find domain of definition.
- ② Evaluate limits at the boundary of domain and deduce relative asymptotes.
- ③ Find first derivative. Deduce maxima and draw variation table.
- ④ Find second derivative. Deduce inflection points and draw concavity table.
- ⑤ Find x and y intercepts.
- ⑥ Find additional points.
- ⑦ Sketch the curve.

For other function, you may need to study parity and periodicity. Also, you may need to find tangent lines at remarkable points (maxima, inflection points, x and y intercepts)


Exercise 2.4 Learner's Book page 91

1. Domain of definition: $]-\infty, 0[\cup]0, +\infty[$

Vertical asymptote $x = 0$

$f(x)$ decreases on interval $]-\infty, 0[$ and increases on interval $]0, +\infty[$.

Curve




2. Domain of definition: $]-1, +\infty[$

Vertical asymptote: $x = -1$

$g(x)$ increases on its domain

Curve




3. Domain of definition: $]0, +\infty[$

Vertical asymptote: $x = 0$

$h(x)$ increases on its domain

Curve




4. Domain of definition: $]-\infty, 1[\cup]2, +\infty[$

Vertical asymptote: $x = 1$ and $x = 2$

$k(x)$ decreases on interval $]-\infty, 1[$ and increases on interval $]2, +\infty[$

Curve



Lesson 2.5. Domain and range of logarithmic function with any base

Learning objectives

Given a logarithmic function with any base, learners should be able to find the domain and range accurately.

Prerequisites

- ✔ Domain of a natural logarithmic function.
- ✔ Range of a natural logarithmic function.

Teaching Aids

Exercise book, pen and calculator



Activity 2.5 Learner's Book page 91

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education
- ✔ Financial education

Answers

1. Existence condition: $x > 0$

Hence, $Domf =]0, +\infty[$

Limit on boundaries

$$\lim_{x \rightarrow 0^+} \frac{\ln x}{\ln 2} = -\infty \qquad \lim_{x \rightarrow \infty} \frac{\ln x}{\ln 2} = +\infty$$

From limits on boundaries, we get that range of $f(x)$ is $]-\infty, +\infty[$.

2. Existence condition: $x > 0$

Hence, $Domg =]0, +\infty[$

Limit on boundaries

$$\lim_{x \rightarrow 0^+} \frac{\ln x}{\ln \frac{1}{2}} = +\infty \qquad \lim_{x \rightarrow \infty} \frac{\ln x}{\ln \frac{1}{2}} = \lim_{x \rightarrow \infty} \frac{\ln x}{-\ln 2} = -\infty$$

From limits on boundaries, we get that range of $g(x)$ is $]-\infty, +\infty[$.

Synthesis

Logarithm of a real number x with base a is the number

denoted $\log_a x$ defined by $\log_a x = \frac{\ln x}{\ln a}$, $x \in \mathbb{R}_0^+$, $a \in \mathbb{R}_0^+ \setminus \{1\}$

By letting $a = 2$, the curve of $f(x) = \log_2 x$ is the following




Figure 2.2: Curve of $y = \log_2 x$

By letting $a = \frac{1}{2}$, you get the curve of $f(x) = \log_{\frac{1}{2}} x$ as illustrated in figure 2.2.




Figure 2.3: Curve of $y = \log_{\frac{1}{2}} x$

Exercise 2.5 Learner's Book page 94

- | | |
|--------------------------------------|--------------------------------------|
| 1. $]0, +\infty[$ | 2. $]-\infty, -1[\cup]1, +\infty[$ |
| 3. $]-\infty, -1[\cup]4, +\infty[$ | 4. $]-5, -2[\cup]-2, 0[$ |

Lesson 2.6. Limit of logarithmic function with any base

Learning objectives

Given a logarithmic function with any base, learners should be able to find limits and deduce its relative asymptotes accurately.

Prerequisites

- ✔ Limit of natural logarithmic function.

Teaching Aids

Exercise book, pen

Activity 2.6 Learner's Book page 94



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $Domf =]0, +\infty[$ and $\ln 3 > 0$,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{\ln x}{\ln 3} = \frac{\lim_{x \rightarrow 0^+} \ln x}{\ln 3} = \frac{-\infty}{\ln 3} = -\infty$$

There is a vertical asymptote $VA \equiv x = 0$

$$\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow +\infty} \frac{\ln x}{\ln 3} = \frac{\lim_{x \rightarrow +\infty} \ln x}{\ln 3} = \frac{+\infty}{\ln 3} = +\infty$$

There is no horizontal asymptote

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{\ln x}{x \ln 3} = \frac{\lim_{x \rightarrow +\infty} \frac{\ln x}{x}}{\ln 3} = \frac{0}{\ln 3} = 0$$

There is no oblique asymptote

2. For $0 < \frac{1}{3} < 1$, $\ln \frac{1}{3} < 0$,

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0^+} \frac{\ln x}{\ln \frac{1}{3}} = \frac{\lim_{x \rightarrow 0^+} \ln x}{\ln \frac{1}{3}} = \frac{-\infty}{\ln \frac{1}{3}} = +\infty$$

There is a vertical asymptote $VA \equiv x = 0$

$$\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow +\infty} \frac{\ln x}{\ln \frac{1}{3}} = \frac{\lim_{x \rightarrow +\infty} \ln x}{\ln \frac{1}{3}} = \frac{+\infty}{\ln \frac{1}{3}} = -\infty$$

There is no horizontal asymptote

$$\lim_{x \rightarrow +\infty} \frac{f(x)}{x} = \lim_{x \rightarrow +\infty} \frac{\ln x}{x \ln \frac{1}{3}} = \frac{\lim_{x \rightarrow +\infty} \frac{\ln x}{x}}{\ln \frac{1}{3}} = \frac{0}{\ln \frac{1}{3}} = 0$$

There is no oblique asymptote.

Synthesis

Figure 2.4 and figure 2.5, are helpful to note that

$$\lim_{x \rightarrow 0} \log_3 x = -\infty \text{ and } \lim_{x \rightarrow +\infty} \log_3 x = +\infty$$

$$\lim_{x \rightarrow 0} \log_{\frac{1}{3}} x = +\infty \text{ and } \lim_{x \rightarrow +\infty} \log_{\frac{1}{3}} x = -\infty.$$




Figure 2.4: Curve of $y = \log_3 x$




Figure 2.5: Curve of $y = \log_{\frac{1}{3}} x$

Generally, calculating limit of logarithmic function with any base, for example $\log_a x$, from definition

$\log_a x = \frac{\ln x}{\ln a}$, you get the following results:

$$\textcircled{\heartsuit} \quad \lim_{x \rightarrow 0^+} f(x) = \begin{cases} -\infty & \text{if } a > 1 \\ +\infty & \text{if } 0 < a < 1 \end{cases}$$

Thus, there is a vertical asymptote $VA \equiv x = 0$

$$\textcircled{\heartsuit} \quad \lim_{x \rightarrow +\infty} f(x) = \begin{cases} +\infty & \text{if } a > 1 \\ -\infty & \text{if } 0 < a < 1 \end{cases}$$

Then, there is no horizontal asymptote. In addition, no oblique asymptote.

Exercise 2.6 Learner's Book page 95

1. $+\infty$ 2. $+\infty$ 3. $-\infty$ 4. $+\infty$

Lesson 2.7. Logarithmic differentiation

Learning objectives

Given a logarithmic function with any base, learners should be able to find its derivative accurately.

Prerequisites

- ✔ Differentiation of natural logarithmic functions.

Teaching Aids

Exercise book and pen



Activity 2.7 Learner's Book page 95

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. f(x) = \frac{\ln x}{\ln 2} \Rightarrow f'(x) = \frac{\frac{1}{x}}{\ln 2} = \frac{1}{x \ln 2}$$

$$2. \left(\frac{\ln x^2}{\ln 2} \right)' = \frac{\frac{1}{x^2} (x^2)'}{\ln 2} = \frac{2x}{x^2 \ln 2} = \frac{2}{x \ln 2}$$

Synthesis

As conclusion, $(\log_a x)' = \frac{1}{x \ln a}$. Also, if u is another differentiable function of x , then

$$(\log_a u)' = \frac{u'}{u \ln a}$$

Exercise 2.7 Learner's Book page 96

1.
$$\frac{2x+2}{(x^2+2x+1)\ln 10}$$

2.
$$-\frac{6}{(x^2-4x-5)\ln 2}$$

3.
$$\frac{-3x^2-2}{(2x^3+4x-16)\ln 2}$$

4.
$$-\frac{\sqrt{x} \sin \sqrt{x}}{2x \ln 3 \cos \sqrt{x}}$$

Lesson 2.8. Further logarithmic differentiation**Learning objectives**

Given a function containing more complicated products and quotients, learners should be able to differentiate it moderately.

Prerequisites

- ✔ The laws of logarithms,
- ✔ The derivative of logarithmic functions, and
- ✔ The differentiation of implicit functions.

Teaching Aids

Exercise book and pen.

Activity 2.8 Learner's Book page 96

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad y = \frac{x+1}{x-3}$$

Taking \ln on both sides gives $\ln y = \ln \frac{x+1}{x-3}$

Applying laws of logarithms, we get

$$\ln y = \ln(x+1) - \ln(x-3)$$

2. Differentiating with respect to x yields

$$\frac{1}{y} \frac{dy}{dx} = \frac{1}{x+1} - \frac{1}{x-3}$$

$$\text{Rearranging gives } \frac{dy}{dx} = y \left(\frac{1}{x+1} - \frac{1}{x-3} \right)$$

$$\text{Substituting for } y \text{ gives } \frac{dy}{dx} = \left(\frac{x+1}{x-3} \right) \left(\frac{1}{x+1} - \frac{1}{x-3} \right)$$

Synthesis

For functions containing more complicated products and quotients, differentiation is often made easier if the logarithm of the function is taken before differentiating. And then apply the property of differentiation of implicit functions.

Exercise 2.8 Learner's Book page 98

$$1. \quad \frac{dy}{dx} = \frac{(x-2)(x+1)}{(x-1)(x+3)} \left(\frac{1}{x-2} + \frac{1}{x+1} - \frac{1}{x-1} - \frac{1}{x+3} \right)$$

$$2. \quad \frac{dy}{dx} = \frac{(2x-1)\sqrt{x+2}}{(x-3)\sqrt{(x+1)^3}} \left(\frac{2}{2x-1} + \frac{1}{2(x+1)} - \frac{1}{x-3} - \frac{3}{2(x+1)} \right)$$

$$3. \quad \frac{dy}{d\theta} = 3\theta \sin \theta \cos \theta \left(\frac{1}{\theta} + \tan \theta - \cot \theta \right)$$

$$4. \quad \frac{dy}{dx} = \frac{x^3 \ln 2x}{e^x \sin x} \left(\frac{3}{x} + \frac{1}{x \ln 2x} - 1 - \cot x \right)$$

$$5. \quad \frac{dy}{dx} = \frac{2x^4 \tan x}{e^{2x} \ln 2x} \left(\frac{4}{x} + \frac{1}{\sin x \cos x} - 2 - \frac{1}{x \ln 2x} \right)$$

Lesson 2.9. Variation and curves of logarithmic functions with any base

Learning objectives

Given a logarithmic function with any base, learners should be able to study the variation and sketch its curve accurately.

Prerequisites

- ✔ Finding domain and limits at the boundaries of the domain.
- ✔ Deducing relative asymptotes.
- ✔ Finding first and second derivative.
- ✔ Variation and concavity of a function.
- ✔ Sketch a curve in Cartesian plane given some points.

Teaching Aids

Exercise book, pencil, instrument of geometry and calculator

Activity 2.9 Learner's Book page 98



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $f(x) = \log_2 x$

a) From Activity 2.6

$$\lim_{x \rightarrow 0^+} f(x) = -\infty$$

There is a vertical asymptote $VA \equiv x = 0$

$$\lim_{x \rightarrow +\infty} f(x) = +\infty$$

There is no horizontal asymptote. In addition, no oblique asymptote.

b) $f(x) = \left(\frac{\ln}{\ln 2} \right) = \frac{1}{\ln 2}$

For $\ln 2 > 0$, $f'(x) = \frac{1}{x \ln 2} > 0$ since $x > 0$

The function $f(x) = \log_2 x$ increases on its

domain $f'(x) = \frac{1}{x \ln 2} \neq 0$, for $\forall x > 0$, no extrema.

c) $f''(x) = \left(\frac{1}{x \ln 2} \right)' = -\frac{1}{x^2 \ln 2}$

For $\ln 2 > 0$, $f''(x) = -\frac{1}{x^2 \ln 2} < 0$ since $x^2 > 0$

The concavity of function $f(x) = \log_2 x$ turns downward on domain of $f(x)$.

$f''(x) = -\frac{1}{x^2 \ln 2} \neq 0$, for $\forall x > 0$, no inflection points.

d) Intersection of $f(x)$ with axes of co-ordinates:

No intersection with y -axis since this axis is a vertical asymptote.

Intersection with x -axis :

$$\log_2 x = 0 \Leftrightarrow \frac{\ln x}{\ln 2} = 0 \Leftrightarrow \ln x = 0 \Rightarrow x = 1.$$


Hence, $f(x) \cap ox = \{(1, 0)\}$

e) Additional points for $f(x) = \log_2 x$

x	0.1	0.3	0.5	0.7	0.9	1.1	1.3	1.5	1.7	1.9
y	-3.3	-1.7	-1.0	-0.5	-0.2	0.1	0.4	0.6	0.8	0.9

x	2.1	2.3	2.5	2.7	2.9	3.1	3.3	3.5	3.7	3.9
y	1.1	1.2	1.3	1.4	1.5	1.6	1.7	1.8	1.9	2.0

Curve



2. $g(x) = \log_{\frac{1}{2}} x$

a) From Activity 2.6

$$\lim_{x \rightarrow 0^+} g(x) = +\infty$$

There is a vertical asymptote $VA \equiv x = 0$

$$\lim_{x \rightarrow +\infty} g(x) = -\infty$$

There is no horizontal asymptote. In addition, no oblique asymptote.

b)
$$g'(x) = \left(\frac{\ln x}{\ln \frac{1}{2}} \right)' = \frac{\frac{1}{x}}{\ln \frac{1}{2}} = \frac{1}{x \ln \frac{1}{2}} = \frac{2}{-x \ln 2},$$

since $\ln \frac{1}{2} = -\ln 2$

For $\ln 2 > 0$, $g'(x) = \frac{1}{-x \ln 2} < 0$ since $x > 0$

The function $g(x) = \log_{\frac{1}{2}} x$ decreases on its domain

$$g'(x) = \frac{1}{-x \ln 2} \neq 0, \text{ no extrema.}$$

c)
$$g''(x) = \left(\frac{1}{-x \ln 2} \right)' = \frac{1}{x^2 \ln 2}$$

For $\ln 2 > 0$, $g''(x) = \frac{1}{x^2 \ln 2} > 0$ since $x^2 > 0$

The concavity of function $g(x) = \log_{\frac{1}{2}} x$ turns upward on domain of $g(x)$.

$$g''(x) = \frac{1}{x^2 \ln 2} \neq 0, \text{ no inflection points.}$$

d) Intersection of $f(x)$ with axes of co-ordinates

No intersection with y -axis since this axis is a vertical asymptote.

Intersection with x -axis :


$$\log_{\frac{1}{2}} x = 0 \Leftrightarrow \frac{\ln x}{\ln 2} = 0 \Leftrightarrow \ln x = 0 \Rightarrow x = 1.$$

Hence, $f(x) \cap ox = \{(1, 0)\}$.

e) Additional points for $g(x) = \log_{\frac{1}{2}} x$

x	0.1	0.3	0.5	0.7	0.9	1.1	1.3	1.5	1.7	1.9
y	3.3	1.7	1.0	0.5	0.2	-0.1	-0.4	-0.6	-0.8	-0.9

x	2.1	2.3	2.5	2.7	2.9	3.1	3.3	3.5	3.7	3.9
y	-1.1	-1.2	-1.3	-1.4	-1.5	-1.6	-1.7	-1.8	-1.9	-2.0

Curve**Synthesis**

To sketch a function, follow the following steps:

- ④ Find domain of definition.
- ④ Evaluate limits at the boundary of domain and deduce relative asymptotes.
- ④ Find first derivative. Deduce maxima and draw variation table.
- ④ Find second derivative. Deduce inflection points and draw concavity table.
- ④ Find x and y intercepts.
- ④ Find additional points.
- ④ Sketch the curve.

For other function, you may need to study parity and periodicity. Also, you may need to find tangent lines at remarkable points (maxima, inflection points, x and y intercepts).


Exercise 2.9 Learner's Book page 102

1. Domain: $]-1, +\infty[$

Vertical asymptote: $x = -1$

$f(x)$ increases on its domain

Curve




2. Domain: $]2, +\infty[$

Vertical asymptote: $x = 2$

$g(x)$ increases on its domain

Curve




3. Domain: $]-\infty, 0[\cup]0, +\infty[$

Vertical asymptote: $x = 0$

$h(x)$ increases on interval $]-\infty, 0[$ and decreases on interval $]0, +\infty[$

Curve




4. Domain: $]0, +\infty[$

Vertical asymptote: $x = 0$

$k(x)$ decreases on its domain

Curve



Lesson 2.10. Domain and range of exponential functions with base "e"

Learning objectives

Given an exponential function with base "e", learners should be able to find domain and range accurately.

Prerequisites

- ✔ Domain of natural logarithmic function.
- ✔ Range of natural logarithmic function.

Teaching Aids

Exercise book and pen



Activity 2.10 Learner's Book page 102

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education
- ✔ Financial education

Answers

We saw that the domain of $f(x) = \ln x$ is $]0, +\infty[$ and its range is \mathbb{R} . Since $g(x)$ is the inverse of $f(x)$, the domain of $g(x)$ is \mathbb{R} and its range is $]0, +\infty[$.

Synthesis

The domain of definition of $y = e^x$ is $]-\infty, +\infty[$ and its range is $]0, +\infty[$ as illustrated in figure 2.6.




Figure 2.6: Curve of $y = e^x$

Exercise 2.10 Learner's Book page 103

- 1) $\mathbb{R} \setminus \{2, 5\}$ 2) \mathbb{R} 3) $]0, +\infty[$ 4) $[4, +\infty[$

Lesson 2.11. Limit of exponential functions with base "e"

Learning objectives

Given an exponential function with base "e", learners should be able to find limit and deduce relative asymptote accurately.

Prerequisites

- ⑦ Finding limits using table of values.
- ⑦ Deduction of relative asymptotes.

Teaching Aids

Exercise book, pen, calculator and instruments of geometry



Activity 2.11 Learner's Book page 104

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Completed table

x	e^x	x	e^x
-1	0.36787944117144	1	2.7182818
-2	0.13533528323661	2	7.3890561
-5	0.00673794699909	5	148.4131591
-15	0.00000030590232	15	3269017.3724721
-30	0.00000000000009	30	10686474581524.5

2. From table in 1), when x takes values approaching to $-\infty$, e^x takes value closed to zero. Hence,


$$\lim_{x \rightarrow -\infty} e^x = 0.$$

There exists a horizontal asymptote $y = 0$, no oblique asymptote.

Also, when x takes value approaching to $+\infty$, e^x increases without bound. Hence, $\lim_{x \rightarrow +\infty} e^x = +\infty$.

There is no horizontal asymptote.

3. Graph

**Synthesis**

From the above figure, it is clear that

$$\lim_{x \rightarrow -\infty} e^x = 0 \text{ and } \lim_{x \rightarrow +\infty} e^x = +\infty.$$

There exists horizontal asymptote: $H.A \equiv y = 0$.

$$\lim_{x \rightarrow -\infty} \frac{e^x}{x} = 0, \lim_{x \rightarrow +\infty} \frac{e^x}{x} = +\infty$$

There is no oblique asymptote.

Exercise 2.11 Learner's Book page 105

- | | | |
|------------------|------|--------------|
| 1. $\sqrt[3]{e}$ | 2. 0 | 3. $+\infty$ |
| 4. 0 | 5. 0 | |

Lesson 2.12. Derivative of exponential functions with base “e”

Learning objectives

Given an exponential functions with base "e", learners should be able to differentiate it correctly.

Prerequisites

- ✔ Use the derivative of natural logarithmic function.
- ✔ Rule of differentiating inverse functions.

Teaching Aids

Exercise book and pen



Activity 2.12 Learner's Book page 105

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. f'(x) = \frac{1}{(f^{-1})'(y)}$$

$$f(x) = e^x \Rightarrow f^{-1}(x) = \ln x \text{ but } (\ln x)' = \frac{1}{x}$$

$$f'(x) = \frac{1}{\frac{1}{e^x}} = e^x$$

$$\text{Thus, } (e^x)' = e^x$$

$$2. (f \circ g)' = f'(g)g'. \text{ Then, } (e^u)' = u'e^u$$

Synthesis

$$(e^x)' = e^x \text{ and if } u \text{ is another differentiable function of } x, \\ (e^u)' = u'e^u.$$

Or from the definition of differentiation,

$$\begin{aligned}\frac{df(x)}{dx} &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}, \text{ thus,} \\ \frac{d(e^x)}{dx} &= \lim_{h \rightarrow 0} \frac{e^{x+h} - e^x}{h} \\ &= \lim_{h \rightarrow 0} \frac{e^x(e^h - 1)}{h} = e^x \lim_{h \rightarrow 0} \left(\frac{e^h - 1}{h} \right) \\ &= e^x \ln e \left[\text{as } \lim_{h \rightarrow 0} \left(\frac{e^h - 1}{h} \right) = \ln e \right] \\ &= e^x \ln e = e^x\end{aligned}$$

Therefore, $(e^x)' = \frac{d(e^x)}{dx} = e^x$

Exercise 2.12 Learner's Book page 106

- | | |
|-------------------------------|----------------------------------|
| 1. $2e^{2x-1}$ | 2. $2(e^{2x} + e^{-2x})$ |
| 3. $(1 + \tan^2 x)e^{\tan x}$ | 4. $\frac{(x-2)e^x}{(x-1) x-1 }$ |

Lesson 2.13. Variation and curve of exponential functions with base "e"

Learning objectives

For an exponential function with base "e", learners should be able to study the variation and sketch its curve accurately.

Prerequisites

- ✔ Reflecting a curve about the first bisector.
- ✔ Properties of inverse functions.

Teaching Aids

Exercise book, pencil, calculator and instruments of geometry.



Activity 2.13 Learner's Book page 106

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

When reflecting the curve of $f(x) = \ln x$ about the first bisector, we obtain




Figure 2.6: Reflection of $y = \ln x$ about first bisector

Synthesis

Since e^x is the inverse of $\ln x$, the curve of $g(x) = e^x$ is the image of the curve of $f(x) = \ln x$ with respect to the first bisector, $y = x$. Then, the coordinates of the points for $f(x) = \ln x$ are reversed to obtain the coordinates of the points for $g(x) = e^x$.


Exercise 2.13 Learner's Book page 109

1. Domain: $]-\infty, +\infty[$

Horizontal asymptote: $y = 0$

$f(x)$ increases on its domain

Curve




2. Domain: $]-\infty, 0[\cup]0, +\infty[$

Vertical asymptote: $x = 0$ and horizontal asymptote: $y = 0$

$g(x)$ increases on interval $]1, +\infty[$

$g(x)$ decreases on intervals: $]-\infty, 0[$ and $]0, 1[$

Curve




3. Domain: $]-\infty, +\infty[$

Horizontal asymptote: $y = 0$

$h(x)$ increases on its domain

Curve




4. Domain: $]-\infty, -2[\cup]-2, +\infty[$

Vertical asymptote: $x = -2$ and horizontal asymptote: $y = 0$

$k(x)$ decreases on interval $]-\infty, -2[\cup]-2, -1[$ and increases on interval $]-1, +\infty[$

Curve



Lesson 2.14. Domain and range of exponential functions with any base

Learning objectives

Given an exponential functions with any base, learners should be able to find the domain and range accurately.

Prerequisites

- ✔ Domain of logarithmic function with any base.
- ✔ Range of logarithmic function with any base.

Teaching Aids

Exercise book and pen

Activity 2.14 Learner's Book page 110



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education
- ✔ Financial education

Answers

We know that the domain of $f(x) = \log_a x$ is $]0, +\infty[$ and its range is \mathbb{R} . Since $g(x)$ is the inverse of $f(x)$, the domain of $g(x)$ is \mathbb{R} and its range is $]0, +\infty[$.

Synthesis

The domain of $f(x) = a^x$ with $a > 0$ and $a \neq 1$, is the set of real numbers and its image is the positive real numbers.

Exercise 2.14 Learner's Book page 111

- | | |
|--------------------------------------|---------------------------------------|
| 1. $\mathbb{R} \setminus \{-5, -2\}$ | 2. \mathbb{R} |
| 3. $]-\infty, -1] \cup]3, +\infty[$ | 4. $]-\infty, -3[\cup]-2, +\infty[$ |

Lesson 2.15. Limit of exponential functions with any base

Learning objectives

Given an exponential functions with any base, learners should be able to evaluate limit and deduce relative asymptotes accurately.

Prerequisites

- ✔ Finding limits using table of values.
- ✔ Deduction of relative asymptotes.

Teaching Aids

Exercise book, pen and calculator



Activity 2.15 Learner's Book page 111

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. a) Table of values

x	2^x
-1	0.5
-2	0.25
-5	0.03125
-15	0.0000305176
-30	0.0000000009

x	2^x
1	2
2	4
5	32
15	32768
30	1073741824

b) From table in a), when x takes values approaching to $-\infty$, 2^x takes values closed to zero.

Hence, $\lim_{x \rightarrow -\infty} 2^x = 0$.

There exists a horizontal asymptote $y = 0$, no oblique asymptote.

Also, when x takes values approaching to $+\infty$, 2^x increases without bound. Hence, $\lim_{x \rightarrow +\infty} 2^x = +\infty$.

There is no horizontal asymptote.

2. a) Table of values

x	$\left(\frac{1}{2}\right)^x$
-1	2
-2	4
-5	32
-15	32768
-30	1073741824

x	$\left(\frac{1}{2}\right)^x$
1	0.5
2	0.25
5	0.03125
15	0.0000305176
30	0.0000000009

b) From table in a), when x takes values approaching

to $-\infty$, $\left(\frac{1}{2}\right)^x$ increases without bound. Hence,

$\lim_{x \rightarrow -\infty} \left(\frac{1}{2}\right)^x = +\infty$. There is no horizontal asymptote.

Also, when x takes values approaching to $+\infty$, $\left(\frac{1}{2}\right)^x$

takes values closed to zero. Hence, $\lim_{x \rightarrow +\infty} \left(\frac{1}{2}\right)^x = 0$.

There exists a horizontal asymptote $y = 0$, no oblique asymptote.

Synthesis

If $a > 1$, $\lim_{x \rightarrow -\infty} a^x = 0$ and $\lim_{x \rightarrow +\infty} a^x = +\infty$

If $0 < a < 1$, $\lim_{x \rightarrow -\infty} a^x = +\infty$ and $\lim_{x \rightarrow +\infty} a^x = 0$

There is horizontal asymptote $y = 0$.

No vertical asymptote since the domain is the set of real numbers. In addition there is no oblique asymptote.

Exercise 2.15 Learner's Book page 115

- | | | |
|--------|----------|----------|
| 1. 1 | 2. e^4 | 3. e^2 |
| 4. e | 5. e^k | 6. e^k |

Lesson 2.16. Derivative of exponential functions with any base**Learning objectives**

Given an exponential functions with any base, learners should be able to differentiate it accurately.

Prerequisites

- ✔ Derivative of logarithmic function with any base.
- ✔ Rule of differentiating inverses functions.

Teaching Aids

Exercise book and pen

**Activity 2.16 Learner's Book page 115**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. f'(x) = \frac{1}{(f^{-1})'(y)}$$

$$f(x) = 3^x \Rightarrow f^{-1}(x) = \log_3 x \text{ but } (\log_3 x)' = \frac{1}{x \ln 3}$$

$$f'(x) = \frac{1}{\frac{1}{3^x \ln 3}} = 3^x \ln 3$$

$$\text{Thus, } (3^x)' = 3^x \ln 3$$

$$2. (f \circ g)' = f'(g)g'. \text{ Then, } (3^{\cos x})' = (\cos x)'(3^{\cos x})(\ln 3)$$

$$\text{Or } (3^{\cos x})' = -\sin x(3^{\cos x})(\ln 3)$$

Synthesis

As conclusion, $(a^x)' = a^x \ln a$. Also, if u is another differentiable function of x , we have $(a^u)' = u' a^u \ln a$

Exercise 2.16 Learner's Book page 116

1. a) $-2(0.3)^x \ln(0.3)$ b) $10^x \left(\frac{1}{x} + \ln x \ln 10 \right)$
 c) $\sin x(\sin x + 2x \cos x)$ d) $x(4)^{\ln x} (2 + \ln 4)$
2. a) $\ln 2$ b) $e(e^e + 1)$ c) $\frac{1}{2}$ d) $\frac{1}{2}$

Lesson 2.17. Variation and curve of exponential functions with any base

Learning objectives

Given an exponential function with base any base, learners should be able to study the variation and sketch its curve accurately.

Prerequisites

- ✔ Reflecting a curve about the first bisector.
- ✔ Properties of inverse functions.

Teaching Aids

Exercise book, pencil, calculator and instruments of geometry



Activity 2.17 Learner's Book page 117

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. When reflecting the curve of $f(x) = \log_2 x$ about the first bisector, we obtain




Figure 2.8: Reflection of $y = \log_2 x$ about first bisector

2. When reflecting the curve of $f(x) = \log_{\frac{1}{2}} x$ about the first bisector, we obtain




Figure 2.9: Reflection of $\log_{\frac{1}{2}} x$ about first bisector

Synthesis

As a^x is the inverse of $\log_a x$, we can obtain a curve of a^x by symmetry with respect to the first bisector $y = x$.

If $a = 2$, we have $f(x) = 2^x$.

x	-4	-3.6	-3.2	-2.8	-2.4	-2	-1.6	-1.2	-0.8	-0.4
y	0.06	0.08	0.11	0.14	0.19	0.25	0.33	0.44	0.57	0.76

x	0	0.4	0.8	1.2	1.6	2	2.4	2.8	3.2	3.6	4
y	1.00	1.32	1.74	2.30	3.03	4.00	5.28	6.96	9.19	12.13	16.00

Curve:




Figure 2.10: Curve of 2^x

If $a = \frac{1}{2}$, we have $f(x) = \left(\frac{1}{2}\right)^x$

x	-4	-3.6	-3.2	-2.8	-2.4	-2	-1.6	-1.2	-0.8	-0.4	0
y	16	12.13	9.19	6.96	5.28	4.00	3.03	2.30	1.74	1.32	1.00

x	0.4	0.8	1.2	1.6	2	2.4	2.8	3.2	3.6	4
y	0.76	0.57	0.44	0.33	0.25	0.19	0.14	0.11	0.08	0.06

Curve:




Figure 2.11: Curve of $\left(\frac{1}{2}\right)^x$


Exercise 2.17 Learner's Book page 119

1. Domain: $]-\infty, +\infty[$

Horizontal asymptote: $y = 0$

$f(x)$ increases on its domain

Curve




2. Domain: $]-\infty, +\infty[$

Horizontal asymptote: $y = 0$

$g(x)$ increases on interval $]0, +\infty[$ and decreases on interval: $]0, +\infty[$

Curve




3. Domain: $]-\infty, +\infty[$

Horizontal asymptote: $y = 0$

$h(x)$ decreases on its domain

Curve




4. Domain: $]-\infty, +\infty[$

Horizontal asymptote: $y = 0$

$k(x)$ increases on interval $]-\infty, -1[$ and increases on interval $]-1, +\infty[$

Curve



Lesson 2.18. Compound interest problems

Learning objectives

By reading text books or accessing internet, learners should be able to use logarithmic and exponential functions to solve compound interest problems accurately.

Prerequisites

- ④ Use of logarithmic and exponential functions.

Teaching Aids

Exercise book, pen and textbooks or internet if available.



Activity 2.18 Learner's Book page 120

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Research
- ④ Peace and values education
- ④ Inclusive education
- ④ Financial education

Answers

If P is the principal, n is the number of years, r is the interest rate per period, k is the number of periods per year, and A the total amount at the end of periods, then

$$A = P \left(1 + \frac{r}{k} \right)^{kn}$$

Here, $P = 4000$, $r = 0.06$, $k = 4$, $n = 5$

Then,

$$A = 4000 \left(1 + \frac{0.06}{4} \right)^{4 \times 5} = 4000(1.015)^{20} = 5387.42$$

After 5 years there, will be 5,387.42 *FRW* on the account.

Synthesis

If P is the principal, n is the number of years, k is the interest rate per period, k is the number of periods per year, and A the total amount at the end of periods, then

$$A = P \left(1 + \frac{r}{k} \right)^{kn}$$

Exercise 2.18 Learner's Book page 121

- 1) 11,358.24 FRW
- 2) 7,007.08 FRW
- 3) Approximately 7.9 years
- 4) Approximately 6.3 years
- 5) Approximately 23.1 years

Lesson 2.19. Mortgage amount problems

Learning objectives

By reading text books or accessing internet, learners should be able to use logarithmic and exponential functions to solve mortgage amount problems accurately.

Prerequisites

- ✔ Use of logarithmic and exponential functions.

Teaching Aids

Exercise book, pen and textbooks or internet if available.



Activity 2.19 Learner's Book page 121

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Research
- ✔ Peace and values education
- ✔ Inclusive education
- ✔ Financial education

Answers

The following formula illustrates the relationship:

$$P = \frac{\frac{rM}{n}}{1 - \left(1 + \frac{r}{n}\right)^{-nt}}$$

Where

$P =$ the payment, r the annual rate,

$M =$ the mortgage amount, $t =$ the number of years and

$n =$ the number of payments per year

Here, $P = 800$, $M = 100000$, $n = 12$, $r = 0.09$ and we need t .

Now,

$$800 = \frac{0.09 \times 100000}{12} \frac{1}{1 - \left(1 + \frac{0.09}{12}\right)^{-12t}}$$

$$\Leftrightarrow 800 = \frac{750}{1 - (1.0075)^{-12t}} \quad \Leftrightarrow (1.0075)^{-12t} = -\frac{750}{800} + 1$$

$$\Leftrightarrow 1 - (1.0075)^{-12t} = \frac{750}{800} \quad \Leftrightarrow (1.0075)^{-12t} = \frac{-750 + 800}{800}$$

$$\Leftrightarrow -(1.0075)^{-12t} = \frac{750}{800} - 1 \quad \Leftrightarrow (1.0075)^{-12t} = \frac{1}{16}$$

Take natural logarithm both sides

$$\Leftrightarrow \ln(1.0075)^{-12t} = \ln \frac{1}{16} \quad \Leftrightarrow -12t \ln(1.0075) = \ln \frac{1}{16}$$

$$\Leftrightarrow -12t = \frac{\ln(0.0625)}{\ln(1.0075)} \quad \Leftrightarrow -12t = -371.06 \Rightarrow t = 30.92$$

Then, you have to make payments to pay off the mortgage in approximately 30 years and 11 months. You would have 370 payments of 800 FRW and the last payment would be 850.40 FRW. The interest paid over the term of the mortgage would be 216,850.40 FRW.

Synthesis

There is a relationship between the mortgage amount M , the number of payments per year n , the amount of the payment P , how often the payment is made t , and the interest rate r . The following formula illustrates the

relationship:
$$P = \frac{\frac{rM}{n}}{1 - \left(1 + \frac{r}{n}\right)^{-nt}}$$

Exercise 2.19 Learner's Book page 123

- | | |
|------------------|------------------|
| 1. 2,400,000 FRW | 2. 2,400,000 FRW |
| 3. 12,719.89 FRW | 4. 8.42% |

Lesson 2.20. Population growth problems

Learning objectives

By reading text books or accessing internet, learners should be able to use logarithmic and exponential functions to solve population growth problems accurately.

Prerequisites

- ✔ Use of logarithmic and exponential functions.

Teaching Aids

Exercise book, pen and textbooks or internet if available.



Activity 2.20 Learner's Book page 124

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Research
- ✔ Peace and values education
- ✔ Inclusive education

Answers

If P_0 is the population at the beginning of a certain period and $r\%$ is the constant rate of growth per period, the population n periods will be $P_n = P_0 (1+r)^n$

Here, $P_0 = 1,000$, $r = 0.5$, $n = 5$

$$P_5 = P_0 (1+r)^5 = 1,000(1+0.5)^5 = 1,000(1.5)^5 = 7,593.75$$

Thus, the population of bacteria in flask at the start of day 5 is 7,593.75.

Synthesis

If P_0 is the population at the beginning of a certain period and $r\%$ is the constant rate of growth per period, the population for n periods will be $P_n = P_0(1+r)^n$.

Exercise 2.20 Learner's Book page 125

1. a) 4200 b) 4% c) 5109
2. a) 1,726,458.24 b) 2020

Lesson 2.21. Depreciation value problems

Learning objectives

By reading text books or accessing internet, learners should be able to use logarithmic and exponential functions to solve depreciation value problems moderately.

Prerequisites

- ✔ Use of logarithmic and exponential functions

Teaching Aids

Exercise book, pen and textbooks or internet if available.

Activity 2.21 Learner's Book page 126



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Research
- ✔ Peace and values education
- ✔ Inclusive education

Answers

If V_0 is the value at a certain time, and $r\%$ is the rate of depreciation per period, the value V_t at the end of t periods is $V_t = V_0(1-r)^t$.

Here, $V_0 = 2.3 \times 10^{30}$, $t = 5$, $r = 0.5$ since the number of bacteria halves every second.

Then,

$$\begin{aligned} V_5 &= 2.3 \times 10^{30} (1-0.5)^5 \\ &= 2.3 \times 10^{30} (0.5)^5 = 2.3 \times 10^{30} \times 0.03125 \\ &= 0.071875 \times 10^{30} = 7.2 \times 10^{28} \end{aligned}$$

Thus, 7.2×10^{28} bacteria were left after 5 seconds.

Synthesis

Depreciation (or decay) is negative growth. If V_0 is the value at a certain time, and $r\%$ is the rate of depreciation per period, the value V_t at the end of t periods is $V_t = V_0(1-r)^t$.

Exercise 2.21 Learner's Book page 127

1. $V = x(0.75)^t$
2. 19 years

Lesson 2.22. Earthquake problems

Learning objectives

By reading text books or accessing internet, learners should be able to use logarithmic and exponential functions to solve earthquake problems.

Prerequisites

- ⑦ Use of logarithmic and exponential functions.

Teaching Aids

Exercise book, pen and textbooks or internet if available.



Activity 2.22 Learner's Book page 127

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Research
- ✔ Peace and values education
- ✔ Inclusive education

Answers

The formula $M = \log \frac{I}{S}$ determines the magnitude of an earthquake, where I is the intensity of the earthquake and S is the intensity of a "standard earthquake."

Here,

$$8 = \log \frac{I_1}{S}, \quad 6 = \log \frac{I_2}{S} \qquad \frac{10^8}{10^6} = \frac{\frac{I_1}{S}}{\frac{I_2}{S}} \Rightarrow 100 = \frac{I_1}{I_2}$$

$$10^8 = \frac{I_1}{S}, \quad 10^6 = \frac{I_2}{S}$$

So, the earthquake will be a hundred times stronger.

Synthesis

The magnitude of an earthquake is given by $M = \log \frac{I}{S}$ where I is the intensity of the earthquake and S is the intensity of a "standard earthquake"

Exercise 2.22 Learner's Book page 129

1. 5
2. 2.6
3. a) 39.8 times more intense b) 7.2
4. 1.26 times more intense

Lesson 2.23. Carbon-14 dating problems

Learning objectives

By reading text books or accessing internet, learners should be able to use logarithmic and exponential functions to solve carbon-14 dating problems accurately.

Prerequisites

- ✔ Use of logarithmic and exponential functions

Teaching Aids

Exercise book, pen and textbooks or internet if available.



Activity 2.23 Learner's Book page 130

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Research
- ✔ Peace and values education
- ✔ Inclusive education

Answers

A formula used to calculate how old a sample is by

$$\text{carbon-14 dating is: } t = \frac{\ln\left(\frac{N_f}{N_0}\right)}{-0.693} \times t_{\frac{1}{2}}$$

where $\frac{N_f}{N_0}$ is the percent of carbon-14 in the sample compared to the amount in living tissue, and $t_{\frac{1}{2}}$ is the half-life of carbon-14 ($5,730 \pm 30$ years).

$$\text{Then, } t = \frac{\ln(0.10)}{-0.693} \times 5700 = 18,940 \text{ years old}$$

Synthesis

Carbon dating is used to work out the age of organic material — in effect, any living thing. The technique hinges on carbon-14, a radioactive isotope of the element that, unlike other more stable forms of carbon, decays away at a steady rate. The half-life of a substance is the amount of time it takes for half of that substance to decay.

A formula to calculate how old a sample is by carbon-14 dating is:

$$t = \frac{\ln\left(\frac{N_f}{N_0}\right)}{-0.693} \times t_{\frac{1}{2}}$$

where $\frac{N_f}{N_0}$ is the percent of carbon-14 in the sample compared to the amount in living tissue, and $t_{\frac{1}{2}}$ is the half-life of carbon-14 ($5,730 \pm 30$ years).

Exercise 2.23 Learner's Book page 132

1. 8,260 years
2. 9,953 years
3. 0.239 mg
4. 3.2 per minutes per gram
5. 3,870 years
6. a) A common rule of thumb is that a radioactive dating method is good out to about 10 half-lives. Given a Carbon-14 half-life of 5730 years, you can see that Carbon-14 dating is (theoretically) good out to around 60,000 years (more-or-less). In fact, due to fluctuations in the carbon amount in the atmosphere, modern Carbon-14 dating needs to be correlated to dates determined by analysis of tree-ring records (dendrochronology).

- b) A skull does not have very much (if any) carbon in it after 73 million years. It would not be dated using Carbon-14 dating. In fact, the value of 73 million years is not arrived at by directly testing the skull. Minerals containing radioactive elements are dated and the age of the skull would be assumed to be of the same age as the strata in which it was discovered.

Summary of the unit

1. Logarithmic functions

- Domain of definition and range:

The Natural logarithm of x is denoted as $\ln x$ or $\log_e x$ and defined on positive real numbers, $]0, +\infty[$, its range is all real numbers.

$$\forall x \in]1, +\infty[, \ln x > 0 \text{ and } \forall x \in]0, 1[, \ln x < 0$$

The equation $\ln x = 1$ has, in interval $]0, +\infty[$, a unique solution, a rational number

2.718281828459045235360.... . This number is denoted by e .

$$\text{Hence } \ln x = 1 \Leftrightarrow x = e.$$

$$\text{Generally } e = \lim_{x \rightarrow +\infty} \left(1 + \frac{1}{x} \right)^x$$

- Limits on boundaries:

Logarithmic function $f(x) = \ln x$ being defined on $]0, +\infty[$, $\lim_{x \rightarrow +\infty} \ln x = +\infty$ and $\lim_{x \rightarrow 0^+} \ln x = -\infty$.

From $\lim_{x \rightarrow +\infty} \ln x = +\infty$, we deduce that there is no horizontal asymptote.

From $\lim_{x \rightarrow 0^+} \ln x = -\infty$, we deduce that there exists a vertical asymptote with equation $VA \equiv x = 0$

- Derivative of natural logarithmic functions or logarithmic derivative:

$$x \in \mathbb{R}_0^+, (\ln x)' = \frac{1}{x} \text{ and } (\ln x)' > 0$$

Also, if u is differentiable function at x then,

$$(\ln u)' = \frac{u'}{u}$$

With certain functions containing more complicated products and quotients, differentiation is often made easier if the logarithm of the function is taken before differentiating.

- Domain and limits on boundaries of a logarithmic function with any base:

Logarithm function of a real number x with base a is a function f denoted $f(x) = \log_a x$ and defined by

$$\log_a x = \frac{\ln x}{\ln a}, x \in \mathbb{R}_0^+, a \in \mathbb{R}_0^+ \setminus \{1\}$$

$$\forall x \in \mathbb{R}_0^+, \log_a x = y \Leftrightarrow x = a^y$$

$$\lim_{x \rightarrow 0^+} f(x) = \begin{cases} -\infty & \text{if } a > 1 \\ +\infty & \text{if } 0 < a < 1 \end{cases}$$

There is a vertical asymptote $VA \equiv x = 0$

$$\lim_{x \rightarrow +\infty} f(x) = \begin{cases} +\infty & \text{if } a > 1 \\ -\infty & \text{if } 0 < a < 1 \end{cases}$$

There is no horizontal asymptote nor oblique asymptote.

- Logarithmic Differentiation:

$$\text{If } f(x) = \log_a x, \text{ then } f'(x) = \frac{1}{x \ln a}$$

Also, if u is another differentiable function in x , then

$$(\log_a u)' = \frac{u'}{u \ln a}$$

2. Exponential functions

Exponential function with base “ e ”

⑦ Domain and range of exponential functions with base “ e ”

The domain of definition of $y = e^x$ is $]-\infty, +\infty[$ and its range is $]0, +\infty[$.

Then, $\forall x \in]0, +\infty[, y \in]-\infty, +\infty[: y = \ln x \Leftrightarrow x = e^y$.

⑦ Limit of exponential functions with base “ e ”

$$\lim_{x \rightarrow -\infty} e^x = 0 \text{ and } \lim_{x \rightarrow +\infty} e^x = +\infty$$

There exists horizontal asymptote: $H.A \equiv y = 0$

⑦ Derivative of exponential functions with base “ e ”

$$\forall x \in \mathbb{R}, (e^x)' = e^x$$

If u is another differentiable function at x ,

$$(e^u)' = u' e^u$$

Remarks

1. $\forall y > 0, y = e^{\ln y}$

In particular, $a^x = e^{\ln a^x}$ means $a^x = e^{x \ln a}$.

Hence, to study the function $y = u^v$ is the same as to study the function $y = e^{v \ln u}$ where u and v are two other functions.

2. Whenever an expression to be differentiated contains a term raised to a power which is itself a function of the variable, then logarithmic differentiation must be used. For example, the differentiation of expressions such

as $|x|^x$, $(1-x)^{1-x^2}$, $\sqrt{x+2}$, $(x)^{\sin x}$ and so on can only be achieved using logarithmic differentiation.

3. Applications

a) Compound interest problems

If P is the principal, n is the number of years, r is the interest rate per period, k in the number of periods per year, and A the total amount at the end of periods, then

$$A = P \left(1 + \frac{r}{k} \right)^{kn}.$$

b) Population growth problems

If P_0 is the population at the beginning of a certain period and $r\%$ is the constant rate of growth per period, the population after n periods will be $P_n = P_0(1+r)^n$.

c) Depreciation value problems

Depreciation (or decay) is negative growth. If V_0 is the value at a certain time, and $r\%$ is the rate of depreciation per period, the value V_t at the end of t periods is $V_t = V_0(1-r)^t$.

d) Earthquake problems

Charles Richter defined the magnitude of an earthquake to be $M = \log \frac{I}{S}$ where I is the intensity of the earthquake (measured by the amplitude of a seismograph reading taken 100 km from the epicentre of the earthquake) and S is the intensity of a "standard earthquake" (whose amplitude is $1 \text{ micron} = 10^{-4} \text{ cm}$).

e) Carbon-14 dating problems

Carbon dating is used to work out the age of organic material — in effect, any living thing. By measuring the ratio of the radio isotope to non-radioactive carbon, the

amount of carbon-14 decay can be worked out, thereby giving an age for the specimen in question.

Through research, scientists have agreed that the half-life of C^{14} is approximately 5700 years.

A formula to calculate how old a sample is by carbon-14 dating is:

$$t = \frac{\ln\left(\frac{N_f}{N_0}\right)}{-0.693} \times t_{\frac{1}{2}}$$

where $\frac{N_f}{N_0}$ is the percent of carbon-14 in the sample compared to the amount in living tissue, and $t_{\frac{1}{2}}$ is the half-life of carbon-14 ($5,730 \pm 30$ years).

End of Unit Assessment answers Learner's Book page 138

- | | |
|---------------------------------|----------------------------------------------------------|
| 1. $]1, +\infty[$ | 2. $]0, 1[\cup]1, +\infty[$ |
| 3. $] -4, 1[\cup]2, +\infty[$ | 4. $\mathbb{R} \setminus \{2 - \sqrt{3}, 2 + \sqrt{3}\}$ |
| 5. $]0, 1[\cup]1, +\infty[$ | 6. $\mathbb{R} \setminus \{-2, -1, 0\}$ |
| 7. \mathbb{R} | 8. $\mathbb{R} \setminus \{1\}$ |
| 9. 0 | 10. $+\infty$ |
| 11. 0 | 12. $+\infty$ |
| 13. 1 | 14. 0 |
| 15. $+\infty$ | 16. 0 |
| 17. $+\infty$ | 18. $+\infty$ |
| 19. $\frac{2}{3}$ | 20. e |
| 21. $\frac{1}{\sqrt[3]{e^2}}$ | 22. 0 |

23. $x^2 e^x$

24. $\frac{1}{2} \left(\tan \frac{x}{2} + \cot \frac{x}{2} \right)$

25. $\frac{1}{x \ln x}$

26. $\frac{\sqrt{x^2 + a^2}}{x^2 + a^2}$

27. $\frac{2e^x}{e^{2x} + 2e^x + 1}$

28. $\frac{-2\sqrt{x^2 + 1}}{x^2 + 1}$

29. $\frac{x^x \ln x}{e^x}$


30. $(\cos x)^x [\ln(\cos x) - x \tan x]$

31. Domain: $]-\infty, +\infty[$

Horizontal asymptote: $y = 0$

$f(x)$ increases on intervals $]-\infty, -2[$ and $]0, +\infty[$, it decreases on interval $]-2, 0[$

Curve




32. Domain: $]-\infty, +\infty[$

No asymptote

$f(x)$ decreases on interval $]-\infty, 0[$, it increases on interval $]0, +\infty[$.

Curve




33. Domain: $]-\infty, +\infty[$

Vertical asymptote $x = 1$ and horizontal asymptote

$y = 0$

$f(x)$ increases on its domain

Curve




34. Domain: $]0, +\infty[$

Vertical asymptote $x = 0$ and horizontal asymptote

$y = 0$


$f(x)$ increases on interval $]0, \sqrt{e}[$ and decreases on interval $]\sqrt{e}, +\infty[$.

Curve



35. a) $f(t) = 1,000,000(0.9)^t$ b) $t = 109.27$ min
 36. a) $f(t) = 75,000 \times e^{0.98083t}$ b) 2,942,490
 37. \$3,315.53
 38. Monthly payment is \$550.32. Interest is \$123,115.20
 39. \$72,537.23
 40. a) $f(t) = 3^t$ b) 4.239×10^{28}
 41. a) $f(t) = 100,000 \left(\frac{1}{2}\right)^t$ b) 97.65625
 42. a) 7.3 b) 125,892,451 as greater as A_0
 43. 8.43
 44. a) $3.16 \times 10^{-3} \text{ mol/l}$ b) 12,589 times more acidic
 45. 1,000,000 times more intense
 46. 70dB





Taylor and Maclaurin's Expansions

Learner's Book pages 143 - 186

Key unit competence

Use Taylor and Maclaurin's expansion to solve problems about approximations, limits, ...

Extend the Maclaurin's expansion to Taylor series.

Vocabulary or key words concepts

Power series: Infinite series of the form $\sum_{n=0}^{\infty} a_n (x-c)^n$.

Taylor series of function $f(x)$ at point x_0 : The infinite series of the form $\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x-x_0)^n$.

Maclaurin series: The special case of the Taylor series when $x_0 = 0$.

Lagrange remainder: The remainder function in Taylor series.

Guidance on the problem statement

The problem statement is

"Suppose that we need to complete the table below.

Angle, x	0^0	1^0	2^0	3^0	4^0	5^0
$\sin x$						

For $x = 0^\circ$ is very easy since this angle is a remarkable angle. But, what about other angles, $1^\circ, 2^\circ, 3^\circ, 4^\circ, 5^\circ$? How can we find their sine without using sine button on scientific calculator?

To solve this problem, we need the Maclaurin series of $\sin x$ and then x will be replaced by its value, remembering that all angles must be expressed in radian.

List of lessons

No	Lesson title	Number of periods
1	Finite series	1
2	Infinite series	1
3	Test for convergence of series	1
4	Power series	1
5	Taylor and Maclaurin series	2
6	Taylor series by using Maclaurin series	1
7	Calculation of limits	1
8	Estimation of the number e	1
9	Estimation of the number π	1
10	Estimation of trigonometric number of an angle	1
11	Estimation of an irrational number	1
12	Estimation of a natural logarithm number	1
13	Estimation of roots of equations	1
Total periods		14

Lesson development

Lesson 3.1. Finite series

Learning objectives

Given a finite series, learners should be able to sum that series accurately.

Prerequisites

- ✔ Terms of a series.
- ✔ General term of a series.
- ✔ Sigma notation.

Teaching Aids

Exercise book and pen

Activity 3.1 Learner's Book page 144



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $u_k = f(k) - f(k+1)$
 For $k = 1$, $u_1 = f(1) - f(2)$
 For $k = 2$, $u_2 = f(2) - f(3)$
 For $k = 3$, $u_3 = f(3) - f(4)$
 For $k = 4$, $u_4 = f(4) - f(5)$
 For $k = 5$, $u_5 = f(5) - f(6)$
 ⋮
 For $k = n-1$, $u_{n-1} = f(n-1) - f(n)$
 For $k = n$, $u_n = f(n) - f(n+1)$

2. Adding obtained terms we have

$$\begin{aligned} u_1 + u_2 + u_3 + u_4 + u_5 + \dots + u_{n-1} + u_n &= \\ = f(1) - f(2) + f(2) - f(3) + f(3) - f(4) + f(4) & \\ - f(5) + f(5) - f(6) + \dots + f(n-1) - f(n) + f(n) - f(n+1) & \\ = f(1) - f(n+1) & \end{aligned}$$

Thus, adding these terms, on the right hand side, nearly all the terms cancel out leaving just $f(1) - f(n+1)$ and on the left hand side, is the required sum of the

series. Thus, $\sum_{k=1}^n u_n = f(1) - f(n+1)$.

Synthesis

As conclusion, the sum of the series $u_1, u_2, u_3, \dots, u_n$ is given by $\sum_{k=1}^n u_n = f(1) - f(n+1)$ where $f(k)$ is a function of k .

Exercise 3.1 Learner's Book page 148

1. $1 - \frac{1}{n+1}$

2. $\frac{1}{2} - \frac{1}{4n+2}$

3. $\frac{1}{4}n^4 + \frac{3}{2}n^3 + \frac{11}{4}n^2 + \frac{3}{2}n$

4. $\frac{1}{2} \left(-\frac{1}{n+1} - \frac{1}{n+2} + \frac{3}{2} \right)$

Lesson 3.2. Infinite series

Learning objectives

Given an infinite series or a repeating decimal, learners should be able to find the sum of infinite series or find a rational number represented by the repeating decimal accurately.

Prerequisites

- ④ Evaluating limits

Teaching Aids

Exercise book and pen



Activity 3.2 Learner's Book page 148

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. S_n = \frac{7}{10} + \frac{7}{10^2} + \frac{7}{10^3} + \frac{7}{10^4} + \dots + \frac{7}{10^n} + \dots$$

$$\frac{1}{10} S_n = \frac{1}{10} \left(\frac{7}{10} + \frac{7}{10^2} + \frac{7}{10^3} + \frac{7}{10^4} + \dots + \frac{7}{10^n} + \dots \right)$$

$$\Rightarrow \frac{1}{10} S_n = \frac{7}{10^2} + \frac{7}{10^3} + \frac{7}{10^4} + \frac{7}{10^5} + \dots + \frac{7}{10^{n+1}} + \dots$$

2. Subtracting, we have

$$S_n = \frac{7}{10} + \frac{7}{10^2} + \frac{7}{10^3} + \frac{7}{10^4} + \dots + \frac{7}{10^n} + \dots$$

$$-\frac{1}{10} S_n = -\frac{7}{10^2} - \frac{7}{10^3} - \frac{7}{10^4} - \frac{7}{10^5} - \dots - \frac{7}{10^{n+1}} - \dots$$

$$S_n - \frac{1}{10} S_n = \frac{7}{10} - \frac{7}{10^{n+1}}$$

$$\Rightarrow \frac{9}{10} S_n = \frac{7}{10} \left(1 - \frac{7}{10^n} \right)$$

$$\Rightarrow S_n = \frac{7}{10} \times \frac{10}{9} \left(1 - \frac{7}{10^n} \right) \Rightarrow S_n = \frac{7}{9} \left(1 - \frac{7}{10^n} \right)$$

3. Taking limit as $n \rightarrow +\infty$.

$$\lim_{n \rightarrow \infty} S_n = \lim_{n \rightarrow \infty} \frac{7}{9} \left(1 - \frac{7}{10^n} \right) = \frac{7}{9}$$

Synthesis

It is impossible to add up infinitely many numbers, thus, we will deal with infinite sums by limiting process involving sequences.

An **infinite series** is an expression of the form $u_1 + u_2 + u_3 + \dots + u_k + \dots$ or in sigma notation $\sum_{k=1}^{+\infty} u_k$. The terms u_1, u_2, u_3, \dots are called **terms** of the series.

To carry out this summation process, we proceed as follows:

Let s_n denote the sum of the first n terms of the series. Thus,

$$s_1 = u_1$$

$$s_2 = u_1 + u_2$$

$$s_3 = u_1 + u_2 + u_3$$

$$\vdots$$

$$s_n = u_1 + u_2 + u_3 + \dots + u_n = \sum_{k=1}^n u_k$$

The number s_n is called the n^{th} **partial sum** of the series and the sequence $\{s_n\}_{n=1}^{+\infty}$ is called the **sequence of partial sums**.

Exercise 3.2 Learner's Book page 154

1. a) $\frac{1}{2}n^2 + \frac{1}{2}n$

b) $\frac{1}{3}n^3 + \frac{1}{2}n^2 + \frac{1}{6}n$

c) $\frac{1}{4}n^4 + \frac{1}{2}n^3 + \frac{1}{4}n^2$

d) $\frac{1}{3}n^3 + n^2 + \frac{2}{3}n$

2. a) $\frac{3}{11}$

b) $\frac{5}{6}$

c) $\frac{49}{396}$

Lesson 3.3. Tests for convergence of series

Learning objectives

Given a series and by using comparison test, limit comparison test, the ratio test or the n^{th} root test, learners should be able to test for convergence accurately.

Prerequisites

- ✔ Evaluating limits.
- ✔ Compare two expressions.
- ✔ Compare real numbers.

Teaching Aids

Exercise book and pen

Activity 3.3 Learner's Book page 154



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$\begin{aligned}
 1. \quad \text{a) } \lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} &= \lim_{n \rightarrow \infty} \frac{3^{n+1} + 1}{\frac{5^{n+1}}{3^n + 1}} = \lim_{n \rightarrow \infty} \frac{3^{n+1} + 1}{5^n \times 5} \times \frac{5^n}{3^n + 1} = \lim_{n \rightarrow \infty} \frac{3^n \times 3 + 1}{5(3^n + 1)} \\
 &= \lim_{n \rightarrow \infty} \frac{3^n \left(3 + \frac{1}{3^n}\right)}{5 \times 3^n \left(1 + \frac{1}{3^n}\right)} = \lim_{n \rightarrow \infty} \frac{3 + \frac{1}{3^n}}{5 \times \left(1 + \frac{1}{3^n}\right)} = \frac{3 + 0}{5(1 + 0)} = \frac{3}{5} \\
 \text{b) } \lim_{n \rightarrow \infty} \sqrt[n]{u_n} &= \lim_{n \rightarrow \infty} \frac{\sqrt[n]{n}}{3} = \lim_{n \rightarrow \infty} \frac{n^{\frac{1}{n}}}{3} = \frac{n^0}{3} = \frac{1}{3}
 \end{aligned}$$

2. Taking $\frac{1}{2n-1}$, if we add 1 to the denominator, we get $\frac{1}{2n}$ and then by comparison methods for rational numbers, $\frac{1}{2n-1} > \frac{1}{2n}$ since the numerators are the same and denominator $2n > 2n-1$

Synthesis

Comparison test

Let $\sum_{n=1}^{\infty} a_n$ be a series with positive terms;

- a) $\sum_{n=1}^{\infty} a_n$ converges if there exists a convergent series

$\sum_{n=1}^{\infty} b_n$ such that $a_n \leq b_n$ for all $n > N$, where N is some positive integer.

- b) $\sum_{n=1}^{\infty} a_n$ diverges if there exists a divergent series

$\sum_{n=1}^{\infty} c_n$ such that $a_n \geq c_n$ for all $n > N$, where N is some positive integer.

Limit comparison test

If the series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are two series with positive terms, and $\lim_{n \rightarrow \infty} \frac{a_n}{b_n}$ is finite, both series converge or diverge.

The ratio test

Let $\sum_{n=1}^{\infty} u_n$ be a series with positive terms and let

$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = L$, then;

- the series converges if $L < 1$.
- the series diverges if $L > 1$.
- the series may or may not converge if $L = 1$ (i.e. the test is inconclusive).

The n^{th} root test

Let $\sum_{n=1}^{\infty} u_n$ be a series with positive terms and let $\lim_{n \rightarrow \infty} \sqrt[n]{u_n} = L$, then;

- the series converges if $L < 1$.
- the series diverges if $L > 1$.
- the test is inconclusive $L = 1$.

Exercise 3.3 Learner's Book page 157

- | | | |
|--------------|--------------|--------------|
| 1. Converges | 2. Converges | 3. Converges |
| 4. Diverges | 5. Converges | 6. Converges |

Lesson 3.4. Power series**Learning objectives**

Through examples, learners should be able to define a power series and to find radius of convergence accurately.

Prerequisites

- ✔ Test for convergence of series.

Teaching Aids

Exercise book and pen

Activity 3.4 Learner's Book page 157

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $u_n = (-1)^{n+1} \frac{x^n}{n}$

n^{th} root test:

$$\lim_{n \rightarrow \infty} \sqrt[n]{u_n} = \lim_{n \rightarrow \infty} \sqrt[n]{\left|(-1)^{n+1} \frac{x^n}{n}\right|} = \lim_{n \rightarrow \infty} \sqrt[n]{\frac{|x^n|}{n}} = \lim_{n \rightarrow \infty} \frac{\sqrt[n]{|x^n|}}{\sqrt[n]{n}} = \lim_{n \rightarrow \infty} \frac{|x|}{\frac{1}{n}} = |x|$$

The series is convergence for $|x| < 1$ (and divergence for $|x| > 1$)

2. $u_n = \frac{x^n}{n!}$ and

$$\lim_{n \rightarrow \infty} \left| \frac{u_{n+1}}{u_n} \right| = \lim_{n \rightarrow \infty} \left| \frac{x^{n+1}}{(n+1)!} \times \frac{n!}{x^n} \right| = \lim_{n \rightarrow \infty} \frac{|x|}{n+1} = 0 < 1$$

for all x . Therefore, the series is absolutely convergence.

Synthesis

Power series is like an infinite polynomial. It has the form

$$\sum_{n=0}^{\infty} a_n (x-c)^n = a_0 + a_1 (x-c) + a_2 (x-c)^2 + \dots + a_n (x-c)^n + \dots$$

- ④ The power series converges at $x = c$. Here, the radius of convergence is zero.
- ④ The power series converges for all x , i.e $]-\infty, +\infty[$. Here, the radius of convergence is infinity.
- ④ There is a number R called the **radius of convergence** such that the series converges for all $c - R < x < c + R$ and the series diverges outside this interval.

Exercise 3.4 Learner's Book page 159

- | | |
|-----------------------------------|--------------------------------------------|
| 1. $-3 < x < -1, R = 1$ | 2. $-\frac{2}{3} < x < 0, R = \frac{1}{3}$ |
| 3. $-2 < x < -1, R = \frac{1}{2}$ | 4. $-1 < x < 3, R = 2$ |
| 5. All $x, R \rightarrow \infty$ | 6. $-1 < x < 1, R = 1$ |
| 7. All $x, R \rightarrow \infty$ | 8. $-8 < x < -2, R = 3$ |
| 9. $x = 3, R = 0$ | 10. All $x, R \rightarrow \infty$ |

Lesson 3.5. Taylor and Maclaurin series

Learning objectives

Using power series, learners should be able to give general form of a Taylor and Maclaurin series without errors.

Prerequisites

- ✔ Power series

Teaching Aids

Exercise book and pen



Activity 3.5 Learner's Book page 159

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad f(x) = c_0 + c_1(x-a) + c_2(x-a)^2 + c_3(x-a)^3 + c_4(x-a)^4 + \dots + c_n(x-a)^n + \dots$$

$$f(a) = c_0$$

$$2. \quad f'(x) = c_1 + 2c_2(x-a) + 3c_3(x-a)^2 + 4c_4(x-a)^3 + \dots + nc_n(x-a)^{n-1} + \dots$$

$$f'(a) = c_1$$

$$3. \quad f''(x) = 2 \times c_2 + 3 \times 2 \times c_3(x-a) + 4 \times 3 \times c_4(x-a)^2 + \dots + n(n-1)c_n(x-a)^{n-2} + \dots$$

$$f''(a) = 2!c_2 \Rightarrow c_2 = \frac{f''(a)}{2!}$$

$$4. \quad f'''(x) = 3 \times 2 \times c_3 + 4 \times 3 \times 2 \times c_4(x-a) + \dots + n(n-1)(n-2)c_n(x-a)^{n-3} + \dots$$

$$f'''(a) = 3 \times 2 \times c_3 = 3!c_3 \Rightarrow c_3 = \frac{f'''(a)}{3!}$$

5. $f^{(iv)}(x) = 4 \times 3 \times 2 \times c_4 + \dots + n(n-1)(n-2)(n-3)c_n(x-a)^{n-4} + \dots$

$$f^{(iv)}(a) = 4 \times 3 \times 2 \times c_4 = 4!c_4 \Rightarrow c_2 = \frac{f^{(iv)}(a)}{4!}$$

6. Now, we can see the pattern. If we continue to differentiate and substitute $x = a$, we obtain

$$f^{(n)}(a) = n(n-1)(n-2)(n-3)\dots 1 \times c_n \text{ or using factorial notation; } f^{(n)}(a) = n!c_n$$

Solving we get $c_n = \frac{f^{(n)}(a)}{n!}$

7. Now,

$$f(x) = f(a) + \frac{f'(a)}{1!}(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \frac{f^{(iv)}(a)}{4!}(x-a)^4 + \dots + \frac{f^{(n)}(a)}{n!}(x-a)^n + \dots$$

Using sigma notation, we can write,

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x-a)^n$$

Synthesis

As conclusion, the Taylor series for $f(x)$ is given by

$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!}(x-x_0)^n$ and the Maclaurin series is given by

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}x^n = f(0) + \frac{f'(0)}{1!}x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \dots + \frac{f^{(n)}(0)}{n!}x^n + \dots$$

Exercise 3.5 Learner's Book page 164

a) $6 - 11(x+2) + 6(x+2)^2 - (x+2)^3 + \dots$

b) $\sum_{n=0}^{\infty} \frac{(-1)^n}{2^{k+1}}(x-2)^n$ c) $\sum_{n=0}^{\infty} \frac{(-1)^n}{k!e}(2x-1)^n$

d) $\sum_{n=0}^{\infty} \frac{(-1)^n}{(2n)! \sqrt{2}} \left(x - \frac{\pi}{4}\right)^{2n} + \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)! \sqrt{2}} \left(x - \frac{\pi}{4}\right)^{2n+1}$

Lesson 3.6. Taylor series by using Maclaurin series

Learning objectives

By using Maclaurin series ($x_0 = 0$) without necessarily using Taylor's formula, learners should be able to find the Taylor series for other functions accurately.

Prerequisites

- ☑ Maclaurin series

Teaching Aids

Exercise book and pen

Activity 3.6 Learner's Book page 164



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ☑ Critical thinking
- ☑ Communication
- ☑ Self confidence
- ☑ Cooperation, interpersonal management and life skills
- ☑ Peace and values education
- ☑ Inclusive education

Answers

1. a) $\sin x$

$$\begin{aligned} f(x) &= \sin x & f(0) &= 0 \\ f'(x) &= \cos x & f'(0) &= 1 \\ f''(x) &= -\sin x & f''(0) &= 0 \\ f'''(x) &= -\cos x & f'''(0) &= -1 \\ f^{(4)}(x) &= \sin x & f^{(4)}(0) &= 0 \\ &\vdots & & \end{aligned}$$

Since the derivatives repeat in a cycle of four, we can write the Maclaurin series as follows:

$$\begin{aligned} \sin x &= 0 + \frac{1}{1!}x + \frac{0}{2!}x^2 + \frac{-1}{3!}x^3 + \frac{0}{4!}x^4 + \dots \\ &= x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!} \end{aligned}$$

b) $\cos x$

$$f(x) = \cos x \quad f(0) = 1$$

$$f'(x) = -\sin x \quad f'(0) = 0$$

$$f''(x) = -\cos x \quad f''(0) = -1$$

$$f'''(x) = \sin x \quad f'''(0) = 0$$

$$f^{(4)}(x) = \cos x \quad f^{(4)}(0) = 1$$

$$\vdots$$

Since the derivatives repeat in a cycle of four, we can write the Maclaurin series as follows:

$$\begin{aligned} \cos x &= 1 + \frac{0}{1!}x + \frac{-1}{2!}x^2 + \frac{0}{3!}x^3 + \frac{1}{4!}x^4 + \dots \\ &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \end{aligned}$$

Alternative method

Since $\cos x = \frac{d}{dx}(\sin x)$, we can differentiate the Maclaurin series for $\sin x$ obtained in a) to get one for $\cos x$. That, is,

$$\begin{aligned} \cos x &= \frac{d}{dx}(\sin x) = \frac{d}{dx} \left(x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \right) \\ &= 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \\ &= \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!} \end{aligned}$$

c) $\ln(1+x)$

$$f(x) = \ln(1+x) \quad f(0) = 0$$

$$f'(x) = \frac{1}{1+x} \quad f'(0) = 1$$

$$f''(x) = -\frac{1}{(1+x)^2} \quad f''(0) = -1$$

$$f'''(x) = \frac{2}{(1+x)^3} \quad f'''(0) = 2$$

$$f^{(4)}(x) = -\frac{6}{(1+x)^4} \quad f^{(4)}(0) = -6$$

$$\vdots$$

Now,

$$\begin{aligned}\ln(1+x) &= 0 + \frac{1}{1!}x + \frac{-1}{2!}x^2 + \frac{2}{3!}x^3 + \frac{-6}{4!}x^4 + \dots \\ &= x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \\ &= \sum_{n=1}^{\infty} (-1)^{n-1} \frac{x^n}{n}\end{aligned}$$

2. From results in 1),

$$\begin{aligned}\sin 2x &= 2x - \frac{(2x)^3}{3!} + \frac{(2x)^5}{5!} - \frac{(2x)^7}{7!} + \dots \\ &= 2x - \frac{4}{3}x^3 + \frac{4}{15}x^5 - \frac{8}{315}x^7 + \dots\end{aligned}$$

$$\begin{aligned}\cos 2x &= 1 - \frac{(2x)^2}{2!} + \frac{(2x)^4}{4!} - \frac{(2x)^6}{6!} + \dots \\ &= 1 - 2x^2 + \frac{2}{3}x^4 - \frac{4}{45}x^6 + \dots\end{aligned}$$

$$\begin{aligned}\ln(1+2x) &= 2x - \frac{(2x)^2}{2} + \frac{(2x)^3}{3} - \frac{(2x)^4}{4} + \dots \\ &= 2x - 2x^2 + \frac{8}{3}x^3 - 4x^4 + \dots\end{aligned}$$

Synthesis

As conclusion, in calculating the limit of some functions, find the Maclaurin series for the transcendental functions contained in the given function, simplify and then evaluate the limit.

Exercise 3.6 Learner's Book page 166

$$1. \text{ a) } \frac{1}{3} - \frac{(x-3)}{3^2} + \frac{(x-3)^2}{3^3} + \dots + (-1)^n \frac{(x-3)^n}{3^{n+1}} + \dots$$

$$\text{b) } 1 + \frac{x}{2!} + \frac{x^2}{3!} + \dots + \frac{x^{n-1}}{n!} + \dots$$

$$\text{c) } 1 - \frac{1}{3!}x^2 + \frac{1}{5!}x^4 - \dots + (-1)^{n+1} \frac{1}{(2n-1)!}x^{2n-2} + \dots$$

$$\text{d) } 1 - \frac{\pi^2}{4^2 2!}(x-2)^2 + \frac{\pi^4}{4^4 4!}(x-2)^4 - \dots + (-1)^n \frac{\pi^{2n}}{4^{2n} (2n)!}(x-2)^{2n} + \dots$$

$$2. \quad \begin{array}{ll} \text{a) } 1 + \sum_{n=1}^{\infty} (-1)^n \frac{2^{2n-1}}{(2n)!} x^{2n} & \text{b) } \sum_{n=0}^{\infty} \frac{x^{n+2}}{k!} \\ \text{c) } 1 - \frac{1}{2}x^3 - \frac{1}{8}x^6 - \dots & \text{d) } 1 + 2x + \frac{5}{2}x^2 + \dots \end{array}$$

Lesson 3.7. Calculation of limits

Learning objectives

Given a function involving transcendental functions and by using Maclaurin series, learners should be able to evaluate its limit at a given point correctly.

Prerequisites

- ✔ Maclaurin series of some functions like e^x , $\sin x$, $\cos x$, $\tan x$, $\ln x$,...
- ✔ Limits concepts.

Teaching Aids

Exercise book, pen and calculator



Activity 3.7 Learner's Book page 166

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad \cos 4x = 1 - 8x^2 \quad \text{and} \quad \sin 3x = 3x - \frac{9}{2}x^3$$

$$2. \quad \frac{1 - \cos 4x + x \sin 3x}{x^2} = \frac{1 - (1 - 8x^2) + x \left(3x - \frac{9}{2}x^3 \right)}{x^2}$$

$$= \frac{1 - 1 + 8x^2 + 3x^2 - \frac{9}{2}x^4}{x^2} = \frac{11x^2 - \frac{9}{2}x^4}{x^2} = 11 - \frac{9}{2}x^2$$

$$\text{Then } \lim_{x \rightarrow 0} \frac{1 - \cos 4x + x \sin 3x}{x^2} = \lim_{x \rightarrow 0} \left(11 - \frac{9}{2}x^2 \right) = 11$$

Synthesis

As conclusion, find the Maclaurin series for the transcendental functions contained in the given function, simplify and then evaluate the limit.

Exercise 3.7 Learner's Book page 168

- 1) $-\frac{1}{2}$ 2) 2 3) $\frac{1}{2}$ 4) 0

Lesson 3.8. Estimation of the number e

Learning objectives

Given number e and by using Maclaurin series of e^x , learners should be able to estimate this number to some decimal places perfectly.

Prerequisites

Maclaurin series of e^x

Teaching Aids

Exercise book, pen and calculator

Activity 3.8 Learner's Book page 168



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \frac{x^6}{6!} + \frac{x^7}{7!} + \frac{x^8}{8!} + \frac{x^9}{9!} + \frac{x^{10}}{10!} + \frac{x^{11}}{11!} + \frac{x^{12}}{12!} + \dots + \frac{x^n}{n!} + \dots$$

Putting $x = 1$, we have

$$e = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \frac{1}{5!} + \frac{1}{6!} + \frac{1}{7!} + \frac{1}{8!} + \frac{1}{9!} + \frac{1}{10!} + \frac{1}{11!} + \frac{1}{12!} + \dots$$

Since we need this number to 8 decimal places, we will stop when we reach the decimal term less than 10^{-8} .

Here, $\frac{1}{12!} = 2 \times 10^{-9} < 10^{-8}$, so we will stop at $\frac{1}{12!}$

Then,

$$\begin{aligned} e \approx & 2 + 0.5 + 0.1666667 + 0.04166667 + 0.00833333 + 0.00138889 \\ & + 0.00019841 + 0.00002480 + 0.00000275 + 0.00000027 \\ & + 0.00000003 + 0.00000000 \approx 2.71828182 \end{aligned}$$

Synthesis

By putting $x = 1$ in the development of e , we can easily estimate the value of the number e to desired decimal places.

Exercise 3.8 Learner's Book page 169

- | | |
|-------------------------|-----------------------------|
| 1. $e \approx 2.71$ | 2. $e \approx 2.7182$ |
| 3. $e \approx 2.718281$ | 4. $e \approx 2.7182818284$ |

Lesson 3.9. Estimation of the number π

Learning objectives

Given number π and by using Maclaurin series of $\arctan x$, learners should be able to estimate this number to some decimal places perfectly.

Prerequisites

- ✔ Maclaurin series of $\arctan x$.
- ✔ Change degrees to radians.
- ✔ Find trigonometric number of an angle.

Teaching Aids

Exercise book, pen and calculator



Activity 3.9 Learner's Book page 170

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Let $f(x) = \arctan x$

$$f'(x) = \frac{1}{x^2+1} \Rightarrow f'(0) = 1$$

$$f''(x) = \frac{-2x}{(x^2+1)^2} \Rightarrow f''(0) = 0$$

$$f'''(x) = \frac{6x^2-2}{(x^2+1)^3} \Rightarrow f'''(0) = -2$$

$$f^{(4)}(x) = \frac{-24x^3+24x}{(x^2+1)^4} \Rightarrow f^{(4)}(0) = 0$$

$$f^{(5)}(x) = \frac{120x^4-240x^2+24}{(x^2+1)^5} \Rightarrow f^{(5)}(0) = 24$$

$$f^{(6)}(x) = \frac{-720x^5+2400x^3-720x}{(x^2+1)^6} \Rightarrow f^{(6)}(0) = 0$$

$$f^{(7)}(x) = \frac{5040x^6-25200x^4+15120x^2-720}{(x^2+1)^7} \Rightarrow f^{(7)}(0) = -720$$

$$\text{Then, } \arctan x = x - \frac{2}{3!}x^3 + \frac{24}{5!}x^5 - \frac{720}{7!}x^7 + \dots$$

Or

$$\arctan x = x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \dots + (-1)^n \frac{x^{2n+1}}{2n+1} + \dots$$

The general term is $(-1)^n \frac{x^{2n+1}}{2n+1}$

$$2. \tan x = \frac{\sqrt{3}}{3} \Rightarrow x = \arctan \frac{\sqrt{3}}{3} \Rightarrow x = \frac{\pi}{6}$$

Synthesis

By using the series

$$\arctan x = x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \dots + (-1)^n \frac{x^{2n+1}}{2n+1} + \dots,$$

and $x = \frac{\pi}{6} \Rightarrow \pi = 6x$ we get that

$$\pi = 6 \left[\frac{\sqrt{3}}{3} - \frac{\left(\frac{\sqrt{3}}{3}\right)^3}{3} + \frac{\left(\frac{\sqrt{3}}{3}\right)^5}{5} - \frac{\left(\frac{\sqrt{3}}{3}\right)^7}{7} + \dots + (-1)^n \frac{\left(\frac{\sqrt{3}}{3}\right)^{2n+1}}{2n+1} + \dots \right]$$

or

$$\pi = 6 \frac{\sqrt{3}}{3} - 6 \frac{\left(\frac{\sqrt{3}}{3}\right)^3}{3} + 6 \frac{\left(\frac{\sqrt{3}}{3}\right)^5}{5} - 6 \frac{\left(\frac{\sqrt{3}}{3}\right)^7}{7} + \dots + (-1)^n 6 \frac{\left(\frac{\sqrt{3}}{3}\right)^{2n+1}}{2n+1} + \dots$$

we can easily estimate the number π .

Exercise 3.9 Learner's Book page 171

1. $\pi \approx 3.141$

2. $\pi \approx 3.14159$

3. $\pi \approx 3.1415926$

4. $\pi \approx 3.141592653$

Lesson 3.10. Estimation of trigonometric number of an angle

Learning objectives

Given an angle and by using Maclaurin series of trigonometric functions, learners should be able to estimate the trigonometric number of that angle accurately.

Prerequisites

- ✔ Maclaurin series of trigonometric functions

Teaching Aids

Exercise book, pen and calculator

Activity 3.10 Learner's Book page 171



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots + (-1)^n \frac{x^{2n+1}}{(2n+1)!} + \dots$$

$$2. \text{ If } x = \frac{\pi}{4}, \text{ we get}$$

$$\sin \frac{\pi}{4} = \frac{\pi}{4} - \frac{\left(\frac{\pi}{4}\right)^3}{3!} + \frac{\left(\frac{\pi}{4}\right)^5}{5!} - \frac{\left(\frac{\pi}{4}\right)^7}{7!} + \dots + (-1)^n \frac{\left(\frac{\pi}{4}\right)^{2n+1}}{(2n+1)!} + \dots$$

Since we need $\sin \frac{\pi}{4}$ to 4 decimal places, we will stop when we reach the decimal term less than 10^{-4} .

Here, $\frac{\left(\frac{\pi}{4}\right)^7}{7!} = 3 \times 10^{-5} < 10^{-4}$, so we will stop at $\frac{\left(\frac{\pi}{4}\right)^7}{7!}$

Then,

$$\begin{aligned}\sin \frac{\pi}{4} &= \frac{\pi}{4} - \frac{\left(\frac{\pi}{4}\right)^3}{3!} + \frac{\left(\frac{\pi}{4}\right)^5}{5!} - \frac{\left(\frac{\pi}{4}\right)^7}{7!} \\ &= 0.7854 - 0.0807 + 0.0024 - 0.0000 \\ &= 0.7071\end{aligned}$$

Remember that on the right hand side π is replaced by 3.1415... not 180^0

Synthesis

x being expressed in radian, we can approximate the value of any trigonometric number using the series of trigonometric functions.

Exercise 3.10 Learner's Book page 173

- | | |
|-----------|-------------|
| 1. 0.866 | 2. 0.017452 |
| 3. 0.4226 | 4. -0.70711 |

Lesson 3.11. Estimation of an irrational number

Learning objectives

Given an irrational number and by using Maclaurin series of $(1+x)^m$, learners should be able to estimate correctly that irrational number to some decimal places accurately.

Prerequisites

- ① Maclaurin series of $(1+x)^m$.

Teaching Aids

Exercise book, pen and calculator



Activity 3.1 | Learner's Book page 173

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. 1, 4, 9, 16, 25, 36, 49, 64, 81, 100, 121, ...
2. 2, 8, 18, 32, 50, 72, 98, 128, 162, 200, 242, ...
3. Take 49 and 50. Their ratio is 0.98, closed to 1. We can also take 121 and 128. If we extend the sequence, we can get other two numbers, 289 and 288. Their ratio is 1.003.
4. Now take 289 and 288, try to transform $\sqrt{2}$. Knowing that $288 = 2 \times 144$, we have

$$\sqrt{2} = \sqrt{\frac{289 \times 2 \times 144}{289 \times 144}} = \frac{17}{12} \sqrt{\frac{2 \times 144}{289}} = \frac{17}{12} \sqrt{\frac{288}{289}} = \frac{17}{12} \sqrt{1 - \frac{1}{289}}$$

Synthesis

Using the Maclaurin series of $(1+x)^m$ for $|x| < 1$, we can estimate any irrational number like $\sqrt{2}$, $\sqrt{3}$, $\sqrt[3]{5}$, ...

Procedure:

Suppose that we need to estimate the value of $\sqrt[n]{a}$ to 6 decimal places.

1. Write down a sequence of natural numbers to the power n (as we need n th root).

2. Multiply each term in obtained sequence from 1) by the radicand (here radicand is a).
3. Take two numbers from sequence in 1) and another from sequence in 2) such that their ratio is closed to 1.

Using the obtained numbers from 3), transform the radicand so that it differs little from 1, then use expansion of $(1+x)^n$ to get $\sqrt[n]{a}$.

Exercise 3.11 Learner's Book page 176

- | | |
|-----------------------------|------------------------------|
| 1. $\sqrt{3} \approx 1.732$ | 2. $\sqrt{5} \approx 2.2361$ |
| 3. $\sqrt[3]{2} = 1.259921$ | 4. $\sqrt[3]{4} = 1.587401$ |

Lesson 3.12. Estimation of natural logarithm of a number

Learning objectives

Given a positive real number and by using Maclaurin series of $\ln(1+x)$, learners should be able to estimate a natural logarithm of that number accurately.

Prerequisites

- ☑ Maclaurin series of $\ln(1+x)$ and $\ln(1-x)$.

Teaching Aids

Exercise book, pen and calculator



Activity 3.12 Learner's Book page 176

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ☑ Critical thinking
- ☑ Communication
- ☑ Self confidence
- ☑ Cooperation, interpersonal management and life skills
- ☑ Peace and values education
- ☑ Inclusive education

Answers

$$1. \ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n-1} \frac{x^n}{n} + \dots$$

2. Replacing x with $-x$ in result obtained in 1), we get

$$\ln(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots - \frac{x^n}{n} - \dots$$

3. Subtracting result obtained in 2) from result obtained in 1), we get

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots + (-1)^{n-1} \frac{x^n}{n} + \dots$$

$$-\ln(1-x) = x + \frac{x^2}{2} + \frac{x^3}{3} + \frac{x^4}{4} + \dots + \frac{x^n}{n} + \dots$$

$$\ln(1+x) - \ln(1-x) = 2x + 2\frac{x^3}{3} + 2\frac{x^5}{5} + \dots + 2\frac{x^{2n+1}}{2n+1} + \dots$$

$$\text{Then, } \ln \frac{1+x}{1-x} = 2 \left(x + \frac{x^3}{3} + \frac{x^5}{5} + \dots + \frac{x^{2n+1}}{2n+1} + \dots \right)$$

Synthesis

As conclusion, the relation

$$\begin{aligned} \ln \frac{1+x}{1-x} &= 2 \left(x + \frac{x^3}{3} + \frac{x^5}{5} + \dots + \frac{x^{2n+1}}{2n+1} + \dots \right) \\ &= 2 \sum_{n=0}^{\infty} \frac{x^{2n+1}}{2n+1} \end{aligned}$$

can help us to estimate \ln of any positive number where

$\ln \frac{1+x}{1-x}$ is equated to that number for finding the value of x in the series.

Exercise 3.12 Learner's Book page 178

$$1. \ln 3 \approx 1.0986$$

$$2. \ln 0.8 \approx -0.223$$

$$3. \ln 7 \approx 1.94591$$

$$4. \ln 0.2 \approx -1.61$$

Lesson 3.13. Estimation of roots of equations

Learning objectives

Given an equation and by using Maclaurin series, learners should be able to estimate the roots of that equations accurately.

Prerequisites

- ✔ Maclaurin series of transcendental functions

Teaching Aids

Exercise book, pen and calculator



Activity 3.13 Learner's Book page 178

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \ln(1+x) = x - \frac{x^2}{2}$$

$$2. \text{The equation } \ln(1+x) + x = 0 \text{ becomes } x - \frac{x^2}{2} + x = 0$$

$$\text{or } 2x - \frac{x^2}{2} = 0$$

$$3. 2x - \frac{x^2}{2} = 0 \Rightarrow x \left(2 - \frac{x}{2} \right) = 0$$

$$x = 0 \text{ or } 2 - \frac{x}{2} = 0 \Rightarrow x = 0 \text{ or } x = 4$$

$$4. \text{If } x = 0, \ln(1+0) + 0 = 0 \Leftrightarrow 0 = 0 \text{ TRUE}$$

$$\text{If } x = 4, \ln(1+4) + 0 = 0 \Leftrightarrow \ln 5 = 0 \text{ FALSE}$$

$$\text{Hence, } S = \{0\}$$

Synthesis

The n^{th} order Maclaurin polynomial is helpful to estimate the roots of a given equation involving transcendental functions.

Exercise 3.13 Learner's Book page 179

$$1. \quad S = \left\{ -\frac{\sqrt{10}}{5}, \frac{\sqrt{10}}{5} \right\} \qquad 2. \quad S = \left\{ \frac{1}{2} \right\}$$

$$3. \quad S = \left\{ 0, \frac{4}{7} \right\}$$

Summary of the unit

1. Generalities on series**① Definitions**

A finite series is an expression of the form

$$u_1 + u_2 + u_3 + \dots + u_n \text{ or in sigma notation } \sum_{k=1}^n u_k,$$

where the index of summation, k , takes consecutive integer values from the lower limit, 1, to the upper limit, n . The terms $u_1, u_2, u_3, \dots, u_n$ are called terms of the series and the term u_n is the general term.

To obtain $\sum_{k=1}^n u_k$, the method of difference is usually used i.e.

$$\sum_{k=1}^n u_k = f(1) - f(n+1) \text{ where } u_k = f(k) - f(k+1), \text{ with } f(k) \text{ a function of } k.$$

② Convergence and divergence of a series

Let $\{s_n\}$ be the sequence of partial sums of the series

$$\sum_{k=1}^{+\infty} u_k. \text{ If the sequence } \{s_n\} \text{ converges to a limit } S, \text{ then}$$

the series is said to converge and S is called the sum of the series. We denote this by writing $S = \sum_{k=1}^{+\infty} u_k$.

If the sequence of partial sums of a series diverges, then the series is said to diverge. A divergent series has no sum.

Comparison test

Let $\sum_{n=1}^{\infty} a_n$ be a series with positive terms.

a) $\sum_{n=1}^{\infty} a_n$ converges if there exists a convergent series $\sum_{n=1}^{\infty} b_n$ such that $a_n \leq b_n$ for all $n > N$, where N is some positive integer.

b) $\sum_{n=1}^{\infty} a_n$ diverges if there exists a divergent series $\sum_{n=1}^{\infty} c_n$ such that $a_n \geq c_n$ for all $n > N$, where N is some positive integer.

Limit comparison test

If the series $\sum_{n=1}^{\infty} a_n$ and $\sum_{n=1}^{\infty} b_n$ are two series with positive terms, and $\lim_{n \rightarrow \infty} \frac{a_n}{b_n}$ is finite, both series converge or diverge.

The ratio test

Let $\sum_{n=1}^{\infty} u_n$ be a series with positive terms and let

$\lim_{n \rightarrow \infty} \frac{u_{n+1}}{u_n} = L$, then,

- the series converges if $L < 1$,
- the series diverges if $L > 1$,
- the series may or may not converge if $L = 1$ (i.e., the test is inconclusive).

The n^{th} root test

Let $\sum_{n=1}^{\infty} u_n$ be a series with positive terms and let

$$\lim_{n \rightarrow \infty} \sqrt[n]{u_n} = L, \text{ then,}$$

- the series converges if $L < 1$,
- the series diverges if $L > 1$,
- the test is inconclusive $L = 1$.

2. Power series

Power series is like an infinite polynomial. It has the form

$$\sum_{n=0}^{\infty} a_n (x-c)^n = a_0 + a_1 (x-c) + a_2 (x-c)^2 + \dots + a_n (x-c)^n + \dots$$

Here, c is any real number and a series of this form is called a **power series** centred at c .

Let $f(x) = \sum_{n=0}^{\infty} a_n (x-c)^n$ be the function defined by this power series. $f(x)$ is only defined if the power series converges, so we will consider the domain of the function f to be the set of x values for which the series converges. There are **three** possible cases:

- ⑦ The power series converges at $x = c$. Here, the radius of convergence is zero.
- ⑦ The power series converges for all x , i.e. $]-\infty, +\infty[$. Here, the radius of convergence is infinity.
- ⑦ There is a number R called the radius of convergence such that the series converges for all $c - R < x < c + R$ and the series diverges outside this interval.

3. Taylor and Maclaurin series

If $f(x)$ is a function defined on the open interval (a, b) , and which can be differentiated $(n+1)$ times on (a, b) , then the equality

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x-x_0)^n + R_{n+1}(x)$$

for any values of x and x_0 in (a, b) is called **Taylor's formula** $R_{n+1}(x)$ is called the **remainder function**.

The resulting function (without $R_{n+1}(x)$) is called the Taylor expansion of $f(x)$ with respect to about the point $x = x_0$ of order n .

One of the most common forms of the remainder function is the Lagrange form:

$$R_{n+1}(x) = \frac{(x-x_0)^{n+1}}{(n+1)!} f^{(n+1)}(x_0 + \theta(x-x_0)) \text{ where } 0 < \theta < 10.$$

If $\lim_{n \rightarrow \infty} R_{n+1}(x) = 0$ for some terms in

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x-x_0)^n + R_{n+1}(x), \text{ then the infinite}$$

series

$$f(x) = f(x_0) + \sum_{n=1}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x-x_0)^n$$

is called the Taylor series for $f(x)$.

A **Maclaurin series** is a Taylor series with $x_0 = 0$

Note that if $f(x)$ is a polynomial of degree, then it will have utmost only n non-zero derivatives; all other higher-order derivatives will be identically equal to zero.

The following series are very important. All of them are Maclaurin series ($x_0 = 0$) and, it is possible to find the Taylor series for other functions by using these formulae without necessarily using Taylor's formula.

$$\text{a) } e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$

$$\text{b) } \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} + \dots + (-1)^{n-1} \frac{x^{2n-1}}{(2n-1)!} + \dots$$

$$\text{c) } \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} + \dots + (-1)^n \frac{x^{2n}}{(2n)!}$$

d) If $-1 < x < 1$, then

$$(1+x)^m = 1 + mx + \frac{m(m-1)}{2!}x^2 + \frac{m(m-1)(m-2)}{3!}x^3 + \dots$$

$$+ \frac{m(m-1)(m-2)\dots(m-n+1)}{n!}x^n + \dots$$

Particularly, if $|x| < 1$, then

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + \dots + (-1)^n x^n + \dots$$

If $-1 < x \leq 1$, then

$$\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} + \dots + \frac{(-1)^{n-1} x^n}{n} + \dots$$

End of Unit Assessment answers Learner's Book page 184

1. a) $\frac{1}{3}n^3 + \frac{5}{2}n^2 + \frac{13}{6}n$ b) $\frac{1}{4}n^4 + \frac{5}{2}n^3 + \frac{37}{4}n^2 + 15n$
- c) $-\frac{1}{3}\left(\frac{1}{n+4} + \frac{1}{n+5} + \frac{1}{n+6} - \frac{37}{60}\right)$
- d) $\frac{1}{3}n^3 + \frac{3}{2}n^2 + \frac{7}{6}n$ e) $-\frac{1}{2}\left(\frac{1}{n+1} - \frac{1}{n+2} - \frac{1}{2}\right)$
2. a) $\frac{233}{990}$ b) $\frac{50}{99}$ c) $\frac{11}{999}$
3. a) $-\frac{7}{2} < x < \frac{1}{2}, r = 2$ b) $-1 < x < 1, r = 1$
- c) $-1 \leq x \leq 1, r = 1$ d) $-2 < x < 2, r = 2$
- e) $-2 \leq x < 2, r = 2$ f) $-\frac{3}{2} < x < \frac{3}{2}, r = \frac{3}{2}$

4. a) $(x-1) - \frac{(x-1)^2}{2} + \frac{(x-1)^3}{3} - \frac{(x-1)^4}{2} + \dots$

b) $1 - (x-1) + (x-1)^2 - (x-1)^3 + \dots$

c) $\frac{\sqrt{2}}{2} \left[1 + \left(x - \frac{\pi}{4}\right) - \frac{\left(x - \frac{\pi}{4}\right)^2}{2} - \frac{\left(x - \frac{\pi}{4}\right)^3}{6} - \dots \right]$

5. $1 - \pi^2 \frac{\left(x - \frac{1}{2}\right)^2}{2!} + \pi^4 \frac{\left(x - \frac{1}{2}\right)^4}{4!} + \dots$

6. $5(x-1) + 6(x-1)^2 + 4(x-1)^3 + (x-1)^4$

7. $(x-1)e + (x-1)^2 e + \frac{(x-1)^3}{2} e + \frac{(x-1)^4}{6} e$

8. a) $-\frac{x^4}{3} + \frac{x^6}{45}$ b) $x + \frac{x^3}{2} + \frac{3x^5}{8}$

c) $x - x^2 + \frac{x^3}{2}$ d) $x - x^3 + x^6$

9. $x - \frac{x^2}{2} + \frac{x^3}{3} - \dots; 2\left(x + \frac{x^3}{3} + \frac{x^5}{5} + \dots\right) = 2\sum_{n=1}^{\infty} \frac{x^{2n-1}}{2n-1}$

10. $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3}$. The absolute value

of the remainder term in Lagrange form is

$$\frac{6x^4}{(1+c)^4 4!} = \frac{x^4}{4(1+c)^4} \text{ where } 0 < c < x. \text{ The maximum}$$

value of the remainder term is obtained where $c = 0$

and so, equals $\frac{x^4}{4}$. We must then have $\frac{x^4}{4} < 5 \times 10^{-4}$

and so $x < 0.211$.

11. a) $\sin \pi = 0$ b) cose

12. $x^2 - \frac{x^4}{3!} + \frac{x^6}{5!} - \dots$

$$13. \quad \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}} = 1 + \frac{1}{2} \frac{v^2}{c^2} + \frac{3}{8} \left(\frac{v^2}{c^2}\right)^2 + \dots$$

And so

$$K = \left(1 + \frac{1}{2} \frac{v^2}{c^2} + \frac{3}{8} \left(\frac{v^2}{c^2}\right)^2 + \dots - 1\right) mc^2 = \frac{1}{2} mv^2 + \frac{3}{8} mv^2 \left(\frac{v^2}{c^2}\right)^2 + \dots$$

This is approximately $K = \frac{1}{2} mv^2$ if $v \ll c$ since the neglected terms are small.

$$14. \quad 1 - \frac{x^2}{2} + \frac{5x^4}{24} + \dots; \frac{1}{2}$$

$$15. \quad x - \frac{x^3}{3} + \frac{x^5}{10} + \dots; \frac{1}{3}$$

$$16. \quad -1 + \sqrt{2}$$

$$17. \quad x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots \text{ and } 1 - \frac{x}{2} + \frac{x^2}{3} - \frac{x^3}{4} + \dots$$

$$18. \quad 1 - \frac{x}{2} + \frac{3x^2}{8} + \dots$$

$$19. \quad 1 + x^2 + x^4 + \dots$$

$$20. \quad \frac{1}{6} + \frac{11x}{36} + \frac{49x^2}{216} + \frac{179x^3}{1296} + \dots$$

$$21. \quad x - \frac{7x^3}{6} + \frac{27x^5}{40} + \dots, \text{ limit is } -\frac{7}{6}$$

$$22. \quad 1 + x + \frac{x^2}{2} - \frac{x^4}{8} + \dots$$

$$23. \quad e^{i\theta} = 1 + i\theta - \frac{\theta^2}{2!} - i\frac{\theta^3}{3!} + \frac{\theta^4}{4!} + i\frac{\theta^5}{5!} + \dots,$$

$$e^{i\theta} = 1 - \frac{\theta^2}{2!} + \frac{\theta^4}{4!} + \dots + i \left(\theta - \frac{\theta^3}{3!} + \frac{\theta^5}{5!} + \dots \right)$$

Substituting $\theta = \pi$ gives

$$e^{i\pi} = \cos \pi + i \sin \pi = -1 \Rightarrow e^{i\pi} + 1 = 0$$

24. a) $1 + \frac{x}{3} - \frac{2x^2}{9} + \dots$

b) $\sqrt[3]{n^3 + 1} = n\sqrt[3]{1 + \frac{1}{n^3}}$. In a) put $x = \frac{1}{n^3}$,

$$\begin{aligned}\sqrt[3]{n^3 + 1} &= n\sqrt[3]{1 + \frac{1}{n^3}} = n\left(1 + \frac{1}{3n^3} - \frac{2}{9n^6} + \dots\right) \\ &= n + \frac{1}{3n^2} - \frac{2}{9n^5} + \dots\end{aligned}$$

and,

$$\begin{aligned}\sqrt[3]{n^3 + 1} - n &= n + \frac{1}{3n^2} - \frac{2}{9n^5} + \dots - n \\ &= \frac{1}{3n^2} - \frac{2}{9n^5} + \dots \\ &\approx \frac{1}{3n^2} \quad \text{when } n \text{ is large}\end{aligned}$$

c) Use the limit comparison test with the series $\frac{1}{3n^2}$



Integration

Learner's Book pages 187 - 288

Key unit competence

Use integration as the inverse of differentiation and as the limit of a sum then apply it to find area of plane surfaces, volumes of solid of revolution, lengths of curved lines.

Vocabulary or key words concepts

Primitive function of function $f(x)$: Is a function $F(x)$ such that $F'(x) = f(x)$.

Integration: Process of finding primitive functions (or anti derivative functions).

Indefinite integrals: Primitive functions.

Definite integrals: Primitive functions evaluated at a given closed interval.

Improper integrals: Definite integrals involving infinity limits or a discontinuous point in the interval of integration.

Volume of revolution: Volume obtained when a curve of a function or a surface between two curves is revolved around an axis.

Guidance on the problem statement

The problem statement is; "Suppose a student differentiated a function $f(x)$ and found $3x^2 + 6x + 10$. The question are what is $f(x)$ and how can we obtain function $f(x)$?" Here, we

need $f(x)$ such that $f'(x) = 3x^2 + 6x + 10$, we can separate the terms and say $(x^3)' = 3x^2$, $(3x^2)' = 6x$, $(10x)' = 10$ and then $f(x) = x^3 + 3x^2 + 10x$. Since derivative of a constant is zero, we can write $f(x) = x^3 + 3x^2 + 10x + c$, $c \in \mathbb{R}$.

But this is a simple problem. That is why, in general, the answer for this problem will be found by integration where we write $f(x) = \int (3x^2 + 6x + 10) dx$.

List of lessons

No	Lesson title	Number of periods
1	Differentials	2
2	Definition of indefinite integrals	1
3	Properties of integrals	1
4	Integration by substitution	1
5	Integration of rational function where numerator is expressed in terms of derivative of denominator	2
6	Integration of rational function where degree of numerator is greater or equal to the degree of denominator	2
7	Integration of rational function where denominator is factorised into linear factors	2
8	Integration of rational function where denominator is a quadratic factor	2
9	Integral of the form $\int \sin mx \cos nx dx$, $\int \cos mx \cos nx dx$, $\int \sin mx \sin nx dx$	2
10	Integral of the form $\int \sin^m x \cos^n x dx$	2
11	Integral of the form $\int \tan^m x \sec^n x dx$	2
12	Integral containing $\sin x$, $\cos x$, $\tan x$ on denominator	2

13	Integral containing $\sin^2 x, \cos^2 x$ on denominator	2
14	Integral containing $\sqrt[n]{ax+b}$	2
15	Integral containing $\sqrt{ax^2+bx+c}$	2
16	Integration by parts	2
17	Integration by reduction formulae	2
18	Integration by Maclaurin series	1
19	Definition of definite integrals	1
20	Properties of definite integrals	1
21	Improper integrals: Infinite limits of integration	1
22	Discontinuous integrand	1
23	Calculation of area of plane surface	2
24	Calculation of volume of solid of revolution	2
25	Calculation of arc length of curved lines	2
Total periods		42

Lesson development

Lesson 4.1. Differentials

Learning objectives

Given a function, learners should be able to find differential of that function and the percentage error perfectly.

Prerequisites

- ⑦ Differentiation of a function

Teaching Aids

Exercise book and pen



Activity 4.1 Learner's Book page 188

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Self confidence
- ④ Communication
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

a) If $y = \sin x$, $x = \frac{\pi}{3}$, $\Delta x = 0.006$ then,

$$dy = \cos x dx = \cos\left(\frac{\pi}{3}\right) dx = \frac{1}{2}(0.006) = 0.003$$

Thus, the change in the value of $\sin x$ is approximately 0.003.

b) $\sin\left(\frac{\pi}{3} + 0.006\right) \approx \sin\frac{\pi}{3} + 0.003 = \frac{\sqrt{3}}{2} + 0.003 = 0.869$

Synthesis

As conclusion,

Differential dy is given by $dy = f'(x)dx$ for $y = f(x)$

Whenever one makes an approximation, it is wise to try and estimate how big the error might be. Relative change

in x is $\frac{\Delta x}{x}$ and percentage change in x is $100 \times \frac{\Delta x}{x}$.

Exercise 4.1 Learner's Book page 190

1. a) $df = (2x - 3)dx$ b) $df = -\frac{4}{x^2 + 4x + 4}dx$

c) $df = -\frac{3}{8\sqrt{2-x}}dx$

2. +2

3. 1.75%

4. ± 10

Lesson 4.2. Definition of indefinite integrals

Learning objectives

Through examples, learners should be able to define indefinite integrals rightfully.

Prerequisites

- ✔ Derivative of a function

Teaching Aids

Exercise book, calculator and pen

Activity 4.2 Learner's Book page 191



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Self confidence
- ✔ Communication
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

- | | |
|----------------------------------|------------------------------------------|
| a) $x^3 + c, c \in \mathbb{R}$ | b) $\frac{x^2}{2} + c, c \in \mathbb{R}$ |
| c) $2\sqrt{x}, c \in \mathbb{R}$ | d) $\frac{1}{x} + c, c \in \mathbb{R}$ |

Synthesis

As conclusion, the function $F(x)$ is an indefinite integral of $f(x)$ if $F'(x) = f(x)$.

Exercise 4.2 Learner's Book page 193

- | | |
|-----------------------------------------------------------|--------------------------------------|
| 1. $2x^2 - 5x + c$ | 2. $2x^3 + 2x^2 + 3x + c$ |
| 3. $\frac{1}{4}x^4 + \frac{1}{3}x^3 + \frac{1}{2}x^2 + c$ | 4. $3x^3 - 12x^2 + 16x + c$ |
| 5. $5x + c$ | 6. $-\frac{8}{3}x^3 + 9x^2 - 9x + c$ |

7. $\frac{2}{5}x^5 - x^3 - 5x + c$

8. $\frac{4}{5}x^5 - \frac{4}{3}x^3 + x + c$

9. $\frac{1}{4}x^4 - 2x^3 + 6x^2 - 8x + c$

Lesson 4.3. Properties of integrals

Learning objectives

Through examples, learners should be able to use properties of indefinite integrals accurately.

Prerequisites

Integrals of simple functions.

Teaching Aids

Exercise book, calculator and pen



Activity 4.3 Learner's Book page 193

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Self confidence
- ✔ Communication
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \int f(x) dx = \int \cos x dx = \sin(x) + c$$

$$\frac{d \int f(x) dx}{dx} = \frac{d[\sin(x) + c]}{dx} = \cos x$$

$$\text{Observation: } \frac{d}{dx} \int f(x) dx = f(x)$$

2. Differential of $f(x)$ is $df = \cos x dx$

$$\int df = \int \cos x dx = \sin x + c$$

Observation: $\int df(x) = f(x) + c$

3. $\int 3x dx = \frac{3x^2}{2} + c$ and

$$3 \int x dx = 3 \left(\frac{x^2}{2} + k \right) = \frac{3x^2}{2} + 3k = \frac{3x^2}{2} + c$$

Observation: $\int kf(x) dx = k \int f(x) dx, k \in \mathbb{R}$

4. $\int f(x) dx + \int g(x) dx = \int (x^3 + 3x - 1) dx + \int (x^2 + 2x + 2)$

$$= \frac{x^4}{4} + \frac{3x^2}{2} - x + \frac{x^3}{3} + x^2 + 2x + c$$

$$= \frac{x^4}{4} + \frac{x^3}{3} + \frac{5x^2}{2} + x + c$$

$$\int [f(x) + g(x)] dx = \int (x^3 + 3x - 1 + x^2 + 2x + 2) dx$$

$$= \int (x^3 + x^2 + 5x + 1) dx$$

$$= \frac{x^4}{4} + \frac{x^3}{3} + \frac{5x^2}{2} + x + c$$

Observation:

$$\int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx$$

5. $\frac{d}{dx} \cos(2x+3) = -(2x+3)' \sin(2x+3) = -2 \sin(2x+3)$

$$\int [-\sin(2x+3)] dx = \frac{1}{2} \cos(2x+3) + c$$

6. Hence;

$$\int f(ax+b) dx = \frac{1}{a} F(ax+b) + c \quad a, b, c \in \mathbb{R}, a \neq 0$$

Synthesis

1. The derivative of the indefinite integral is equal to the function to be integrated.

$$\frac{d}{dx} \int f(x) dx = f(x)$$

2. The integral of differential of a function is equal to the sum of that function and an arbitrary constant.

$$\int df(x) = f(x) + c$$

3. Each constant function may be pulled out of integral sign.

$$\int k f(x) dx = k \int f(x) dx, \quad k \in \mathbb{R}$$

4. The indefinite integral of the algebraic sum of two functions is equal to the algebraic sum of the indefinite integrals of those functions.

$$\int [f(x) \pm g(x)] dx = \int f(x) dx \pm \int g(x) dx$$

5. If $F(x)$ is a primitive function of $f(x)$, then, the integral

$$\int f(ax+b) dx = \frac{1}{a} F(ax+b) + c \quad a, b, c \in \mathbb{R}, a \neq 0$$

Exercise 4.3 Learner's Book page 194

1. $4 \int f(x) dx = 4(x^2 + 2x + c) = 4x^2 + 8x + k$
2. $\frac{2}{5} \int [g(x) - 6] dx = \frac{2}{5} \left[\int g(x) dx - \int 6 dx \right]$
 $= \frac{2}{5} \left[\int g(x) dx - 6 \int dx \right]$

$$\begin{aligned}
 &= \frac{2}{5}(x^3 - 3x^2 - 4x + k - 6x) \\
 &= \frac{2}{5}(x^3 - 3x^2 - 10x + k) \\
 &= \frac{2}{5}x^3 - \frac{6}{5}x^2 - 4x + \frac{2}{5}k \\
 &= \frac{2}{5}x^3 - \frac{6}{5}x^2 - 4x + c
 \end{aligned}$$

$$\begin{aligned}
 3. \quad \int [f(x) + 3g(x)] dx &= \int f(x) dx + 3 \int g(x) dx \\
 &= x^2 + 2x + c + 3(x^3 - 3x^2 - 4x + k) \\
 &= x^2 + 2x + c + 3x^3 - 9x^2 - 12x + 3k \\
 &= 3x^3 - 8x^2 - 10x + d
 \end{aligned}$$

$$\begin{aligned}
 4. \quad \frac{d}{dx} \int [2f(x) - 3g(x)] dx &= 2f(x) - 3g(x) \\
 &= 2(2x + 2) - 3(3x^2 - 6x - 4) \\
 &= 4x + 4 - 9x^2 + 18x + 12 \\
 &= -9x^2 + 22x + 16
 \end{aligned}$$

Lesson 4.4. Integration by substitution

Learning objectives

Through examples, learners should be able to find integrals by substitution method correctly.

Prerequisites

- ⑦ Differentiation of a function.

Teaching Aids

Exercise book and pen



Activity 4.4 Learner's Book page 195

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$u = 5x + 2 \Rightarrow du = 5dx$$

$$\begin{aligned} \int e^{5x+2} dx &= \int e^u \frac{du}{5} &&= \frac{1}{5} e^u + c \\ &= \frac{1}{5} \int e^u du &&= \frac{1}{5} e^{5x+2} + c \end{aligned}$$

Synthesis

Integration by substitution is based on rule for differentiating composite functions. Substitution means to let $f(x)$ be a function of another function.

In $\int f(x) dx$, let x be $x(t)$; thus, $dx = x'(t) dt$ and then we get $\int f(x) dx = \int f(x(t)) x'(t) dt$ that is a formula of integration by substitution.

Exercise 4.4 Learner's Book page 196

- | | |
|---------------------------------------|------------------------------------------|
| 1. a) $e^x - \frac{x^{e+1}}{e+1} + c$ | b) $\frac{1}{3}e^{x^3} + c$ |
| c) $\frac{1}{2}e^{2x} + 2e^x + x + c$ | d) $-\frac{1}{2}e^{\frac{1}{x^2}} + c$ |
| e) $\sin(e^x) + c$ | f) $\frac{1}{3}e^{3\cos 2x} \sin 2x + c$ |
| g) $\sin(\ln x) + c$ | h) $\frac{1}{36}(4x^3 - 12)^3 + c$ |
2. $100m$

Lesson 4.5. Integration of rational functions where numerator is expressed in terms of derivative of denominator

Learning objectives

Given a rational function where numerator is expressed in terms of derivative of denominator, learners should be able to find primitive function moderately.

Prerequisites

- ✔ Derivative of $\ln[g(x)]$.
- ✔ Derivative of $\arctan[g(x)]$.

Teaching Aids

Exercise book, calculator and pen

Activity 4.5 Learner's Book page 197



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. We see that $(1-x^2)' = -2x \Rightarrow x = -\frac{1}{2}(1-x^2)'$. So, we can write

$$\begin{aligned} \int \frac{x}{(1-x^2)^2} dx &= \int \frac{-\frac{1}{2}(1-x^2)'}{(1-x^2)^2} dx \\ &= -\frac{1}{2} \int \frac{(1-x^2)'}{(1-x^2)^2} dx \\ &= \frac{1}{2(1-x^2)} + c \quad \text{since } \left(\frac{1}{g}\right)' = -\frac{g'}{g^2} \end{aligned}$$

2. We see that

$$(3x^2 - 3x + 1)' = 6x - 3 = 3(2x - 1) \Rightarrow 2x - 1 = \frac{1}{3}(3x^2 - 3x + 1)'$$

So, we can write

$$\begin{aligned} \int \frac{2x-1}{3x^2-3x+1} dx &= \int \frac{\frac{1}{3}(3x^2-3x+1)'}{3x^2-3x+1} dx \\ &= \frac{1}{3} \int \frac{(3x^2-3x+1)'}{3x^2-3x+1} dx \\ &= \frac{1}{3} \ln|3x^2-3x+1| + c \quad \text{since } (\ln u)' = \frac{u'}{u} \end{aligned}$$

Synthesis

The following basic integration formulae are most helpful:

$$\int \frac{u'}{u} dx = \ln|u| + c, \quad \int \frac{u'}{u^2} dx = -\frac{1}{u} + c \quad \text{and} \quad \int \frac{u'}{u^2+1} dx = \arctan u + c$$

Exercise 4.5 Learner's Book page 199

- | | |
|---------------------------------|-----------------------------------|
| 1. $-\frac{1}{2(x^2+2x+3)} + c$ | 2. $\frac{1}{2(1-x^2)} + c$ |
| 3. $-\frac{1}{6(2x^3+5)} + c$ | 4. $-\frac{1}{4(x^2+2x+5)^2} + c$ |

Lesson 4.6. Integration of rational functions where degree of numerator is greater or equal to the degree of denominator

Learning objectives

Given an irrational function where degree of numerator is greater or equal to the degree of denominator, learners should be able to find primitive function accurately.

Prerequisites

- ✔ Long division of polynomials.

Teaching Aids

Exercise book, calculator and pen

Activity 4.6 Learner's Book page 199



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $\frac{2x+4}{5x-3} = \frac{2}{5} + \frac{26}{25x-15}; \frac{2}{5}x + \frac{26}{25} \ln|5x-3| + c$
2. $\frac{x^2-3x+2}{x^2+1} = 1 + \frac{1-3x}{x^2+1}; x - \frac{3}{2} \ln(x^2+1) - \arctan x + 1 + c$
3. $\frac{x^2+1}{x-1} = x+1 + \frac{2}{x-1}; \frac{x^2}{2} + x + 2 \ln|x-1| + c$
4. $\frac{x^3+2x-4}{x^2+2} = x - \frac{4}{x^2+2}; \frac{x^2}{2} - 2\sqrt{2} \arctan \frac{\sqrt{2}x}{2} + c$

Synthesis

If we want to find $\int \frac{f(x)}{g(x)} dx$ when the degree of $f(x)$ is greater than the degree of $g(x)$, we proceed by long division to find $\int \frac{f(x)}{g(x)} dx = \int q(x) dx + \int \frac{r(x)}{g(x)} dx$ where $q(x)$ is the quotient, $r(x)$ the remainder and then integrate the new expression on the right hand side.

Exercise 4.6 Learner's Book page 200

1. $\frac{1}{2}x^2 - \frac{1}{2}\ln(x^2 + 1) - 2 \arctan x + c$
2. $-\frac{1}{3}\ln|x-1| - \frac{2}{3}\ln|x+2| + x + c$
3. $\frac{1}{6}\ln|x| - \frac{13}{54}\ln|3x-2| - \frac{1}{9}x + c$
4. $\frac{1}{3}(x^3 + a^3 \ln|x^3 - a^3|) + c$
5. $\frac{1}{2}x^2 - 26\ln|x+3| + 63\ln|x+4| - 7x + c$

Lesson 4.7. Integration of rational functions where denominator is factorised into linear factors

Learning objectives

Given an irrational function where denominator is factorised into linear factors, learners should be able to find primitive function accurately.

Prerequisites

- ✔ Factorise completely a polynomial.

Teaching Aids

Exercise book, calculator and pen



Activity 4.7 Learner's Book page 201

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ▼ Critical thinking
- ▼ Communication
- ▼ Self confidence
- ▼ Cooperation, interpersonal management and life skills
- ▼ Peace and values education
- ▼ Inclusive education

Answers

1. $x^2 + 2x = x(x+2)$

$$\frac{x-2}{x^2+2x} = \frac{A}{x} + \frac{B}{x+2} = \frac{A(x+2)+Bx}{x^2+2x}$$

$$x-2 = A(x+2) + Bx$$

Solving, we get

$$\begin{cases} A = -1 \\ B = 2 \end{cases}$$

Then,

$$\frac{x-2}{x^2+2x} = -\frac{1}{x} + \frac{2}{x+2}$$

$$\text{And } \int \frac{x-2}{x^2+2x} dx = -\int \frac{dx}{x} + \int \frac{2}{x+2} dx$$

$$= -\ln|x| + 2\ln|x+2| + c$$

$$= \ln|x| + \ln(x+2)^2 + c = \ln \frac{(x+2)^2}{|x|} + c$$

2. $x^2 + 3x + 2 = (x+1)(x+2)$

$$\frac{x}{x^2+3x+2} = \frac{A}{x+1} + \frac{B}{x+2} = \frac{A(x+2)+B(x+1)}{x^2+3x+2}$$

$$x = A(x+2) + B(x+1)$$

Solving we get,

$$\begin{cases} A = -1 \\ B = 2 \end{cases}$$

Then,

$$\frac{x}{x^2 + 3x + 2} = -\frac{1}{x+1} + \frac{2}{x+2}$$

Therefore;

$$\begin{aligned} \int \frac{x}{x^2 + 3x + 2} dx &= -\int \frac{dx}{x+1} + \int \frac{2}{x+2} dx \\ &= -\ln|x+1| + 2\ln|x+2| + c \\ &= \ln|x+1| + \ln(x+2)^2 + c = \ln \frac{(x+2)^2}{|x+1|} + c \end{aligned}$$

$$\begin{aligned} 3. \quad \frac{2}{x^2 - 1} &= \frac{A}{x-1} + \frac{B}{x+1} \\ \frac{2}{x^2 - 1} &= \frac{A(x+1) + B(x-1)}{x^2 - 1} \\ 2 &= A(x+1) + B(x-1) \end{aligned}$$

Solving, we get

$$\begin{cases} A = 1 \\ B = -1 \end{cases} \quad \frac{2}{x^2 - 1} = \frac{1}{x-1} - \frac{1}{x+1}$$

$$\begin{aligned} \text{Then, } \int \frac{2}{x^2 - 1} dx &= \int \frac{dx}{x-1} - \int \frac{dx}{x+1} \\ &= \ln|x-1| - \ln|x+1| + c = \ln \frac{x-1}{x+1} + c \end{aligned}$$

$$\begin{aligned} 4. \quad \frac{2x-3}{x^2 - x - 2} \\ x^2 - x - 2 &= (x-2)(x+1) \\ \frac{2x-3}{x^2 - x - 2} &= \frac{A}{x-2} + \frac{B}{x+1} = \frac{A(x+1) + B(x-2)}{x^2 - x - 2} \\ 2x-3 &= A(x+1) + B(x-2) \end{aligned}$$

Solving, we get

$$\begin{cases} A = \frac{1}{3} \\ B = \frac{5}{3} \end{cases} \quad \frac{2x-3}{x^2-x-2} = \frac{1}{3(x-2)} + \frac{5}{3(x+1)}$$

Finally,

$$\begin{aligned} \int \frac{2x-3}{x^2-x-2} dx &= \int \frac{dx}{3(x-2)} + \int \frac{5}{3(x+1)} dx \\ &= \frac{1}{3} \ln|x-2| + \frac{5}{3} \ln|x+1| + c = \frac{1}{3} \ln(x-2)(x+1)^5 + c \end{aligned}$$

Synthesis

For integration of rational function where denominator is factorised into linear factors, before integrating, note that to each factor $ax+b$ occurring once in the denominator of a proper rational fraction, there corresponds a single partial fraction of the form $\frac{A}{ax+b}$ where A is a constant to be found, but to each factor $ax+b$ occurring n times in the denominator of a proper rational fraction, there corresponds a sum of n partial fractions $\frac{A_1}{ax+b} + \frac{A_2}{(ax+b)^2} + \dots + \frac{A_n}{(ax+b)^n}$ where A_n are constants to be found, and then integrate the new expression.

Exercise 4.7 Learner's Book page 205

1. $\ln \left| \frac{x-1}{x+1} \right| + c$

2. $\ln \left| \frac{(x+2)^2}{x+1} \right| + c$

3. $\ln \sqrt{\frac{x-2}{x^3}} + c$

4. $\ln|x+1| - \frac{x}{x+1} + c$

5. $3 \ln|x-2| - \frac{3x}{x-2} + c$

6. $\ln|(x+1)^3(x-2)^5| + \frac{12-5x}{x-2} + c$

Lesson 4.8. Integration of rational functions where denominator is a quadratic factor

Learning objectives

Given an irrational function where denominator is a quadratic factor, learners should be able to find primitive function correctly.

Prerequisites

Use of the relation $ax^2 + bx + c = a \left[\left(x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2} \right]$.

Teaching Aids

Exercise book, calculator and pen



Activity 4.8 Learner's Book page 205

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Relation to be used;

$$ax^2 + bx + c = a \left[\left(x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2} \right]$$

$$\int \frac{dx}{x^2 + 3x + 2}, \quad a = 1, \quad b = 3, \quad c = 2$$

$$\int \frac{dx}{x^2 + 3x + 2} = \int \frac{dx}{\left(x + \frac{3}{2} \right)^2 - \frac{1}{4}}$$

$$\int \frac{dx}{x^2 + 3x + 2} = \int \frac{dx}{\left(x + \frac{3}{2} \right)^2 - \frac{1}{4}}$$

$$\text{Let } u = x + \frac{3}{2} \Rightarrow du = dx$$

$$\int \frac{dx}{x^2+3x+2} = \int \frac{du}{u^2 - \frac{1}{4}} = \int \frac{du}{u^2 - \left(\frac{1}{2}\right)^2}$$

Using the formula $\int \frac{dx}{x^2 - k^2} = \frac{1}{2k} \ln \left| \frac{x-k}{x+k} \right| + d$, we have

$$\begin{aligned} \int \frac{dx}{x^2+3x+2} &= \int \frac{du}{u^2 - \left(\frac{1}{2}\right)^2} = \frac{1}{2 \times \frac{1}{2}} \ln \left| \frac{u - \frac{1}{2}}{u + \frac{1}{2}} \right| + d \\ &= \ln \left| \frac{x + \frac{3}{2} - \frac{1}{2}}{x + \frac{3}{2} + \frac{1}{2}} \right| + d = \ln \left| \frac{x+1}{x+2} \right| + d \end{aligned}$$

2. $\int \frac{dx}{x^2 - 4x + 4}$, $a=1$, $b=-4$, $c=4$

$$x^2 - 4x + 4 = \left(x - \frac{4}{2}\right)^2 - \frac{(-4)^2 - 4 \times 4}{4} = (x-2)^2$$

$$\int \frac{dx}{x^2 - 4x + 4} = \int \frac{dx}{(x-2)^2}$$

Let $u = x - 2 \Rightarrow du = dx$

$$\int \frac{dx}{x^2 - 4x + 4} = \int \frac{du}{u^2}$$

Using the formula $\int \frac{u'}{u^2} du = -\frac{1}{u} + d$, we have

$$\int \frac{dx}{x^2 - 4x + 4} = \int \frac{du}{u^2} = -\frac{1}{u} + d = -\frac{1}{x-2} + d$$

3. $\int \frac{dx}{x^2 - 6x + 18}$, $a=1$, $b=-6$, $c=18$

$$x^2 - 6x + 18 = \left(x - \frac{6}{2}\right)^2 - \frac{(-6)^2 - 4 \times 18}{4} = (x-3)^2 + 9$$

$$\int \frac{dx}{x^2 - 6x + 18} = \int \frac{dx}{(x-3)^2 + 9}$$

Let $u = x - 3 \Rightarrow du = dx$

$$\int \frac{dx}{x^2 - 6x + 18} = \int \frac{du}{u^2 + 9} = \int \frac{du}{u^2 + 3^2}$$

Using the formula $\int \frac{dx}{x^2+k^2} = \frac{1}{k} \arctan \frac{x}{k} + d$, we have

$$\int \frac{dx}{x^2+3x+2} = \int \frac{du}{u^2+3^2} = \frac{1}{3} \arctan \frac{u}{3} + d = \frac{1}{3} \arctan \frac{x-3}{3} + d$$

Synthesis

For the integral of the form $\int \frac{dx}{ax^2+bx+c}$,

⊙ If $b^2 - 4ac = 0$, then,

$$\int \frac{dx}{ax^2+bx+c} = \frac{1}{a} \int \frac{dx}{\left(x + \frac{b}{2a}\right)^2} \text{ and we let } u = x + \frac{b}{2a}$$

⊙ If $b^2 - 4ac > 0$, then,

$$\int \frac{dx}{ax^2+bx+c} = \frac{1}{a} \int \frac{dx}{\left(x + \frac{b}{2a}\right)^2 + \frac{b^2-4ac}{4a^2}}. \text{ We let}$$

$$u = x + \frac{b}{2a}, \quad k^2 = \frac{b^2-4ac}{4a^2}$$

and use the standard integral

$$\int \frac{dx}{x^2-k^2} = \frac{1}{2k} \ln \left| \frac{x-k}{x+k} \right| + d$$

⊙ If $b^2 - 4ac < 0$, then,

$$\int \frac{dx}{ax^2+bx+c} = \frac{1}{a} \int \frac{dx}{\left(x + \frac{b}{2a}\right)^2 - \frac{b^2-4ac}{4a^2}}. \text{ We let}$$

$$u = x + \frac{b}{2a}, \quad -k^2 = \frac{b^2-4ac}{4a^2} \text{ and use the standard}$$

$$\text{integral } \int \frac{dx}{x^2+k^2} = \frac{1}{k} \arctan \frac{x}{k} + d$$

Exercise 4.8 Learner's Book page 210

1. $\frac{2}{\sqrt{7}} \arctan \left(\frac{2x+1}{\sqrt{7}} \right) + c$

2. $-\frac{1}{9} \arctan(3x+1) + \frac{1}{18} \ln(9x^2+6x+2) + c$

$$3. -\frac{\sqrt{2}}{2} \arctan(\sqrt{2}x) + 3 \ln|x-2| + c$$

$$4. \frac{1}{2} \ln(x^2 + 2) - \ln|2x+1| + c$$

Lesson 4.9. Integrals of the form

$$\int \sin mx \cos nx \, dx, \int \cos mx \cos nx \, dx, \int \sin mx \sin nx \, dx$$

Learning objectives

Given integrals of the form

$$\int \sin mx \cos nx \, dx, \int \cos mx \cos nx \, dx, \int \sin mx \sin nx \, dx,$$

learners should be able to find primitive function accurately.

Prerequisites

☑ Identities:

$$\sin A \cos B = \frac{1}{2} [\sin(A-B) + \sin(A+B)]$$

$$\sin A \sin B = \frac{1}{2} [\cos(A-B) - \cos(A+B)]$$

$$\cos A \cos B = \frac{1}{2} [\cos(A-B) + \cos(A+B)]$$

Teaching Aids

Exercise book and pen

Activity 4.9 Learner's Book page 211



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ☑ Critical thinking
- ☑ Communication
- ☑ Self confidence
- ☑ Cooperation, interpersonal management and life skills
- ☑ Peace and values education
- ☑ Inclusive education

Answers

$$1. \quad \sin 2x \cos x = \frac{1}{2}(\sin x + \sin 3x) = \frac{1}{2} \sin x + \frac{1}{2} \sin 3x$$

$$\Rightarrow \int \sin 2x \cos x dx = \int \left(\frac{1}{2} \sin x + \frac{1}{2} \sin 3x \right) dx = -\frac{1}{2} \cos x - \frac{1}{6} \cos 3x + c$$

$$2. \quad \sin x \sin 5x = \frac{1}{2}(\cos 4x - \cos 6x) = \frac{1}{2} \cos 4x - \frac{1}{2} \cos 6x$$

$$\Rightarrow \int \sin x \sin 5x dx = \int \left(\frac{1}{2} \cos 4x - \frac{1}{2} \cos 6x \right) dx = \frac{1}{8} \sin 4x - \frac{1}{12} \sin 6x + c$$

$$3. \quad \cos 2x \cos 3x = \frac{1}{2}(\cos x + \cos 5x) = \frac{1}{2} \cos x + \frac{1}{2} \cos 5x$$

$$\Rightarrow \int \cos 2x \cos 3x dx = \int \left(\frac{1}{2} \cos x + \frac{1}{2} \cos 5x \right) dx = \frac{1}{2} \sin x + \frac{1}{10} \sin 5x + c$$

$$4. \quad \sin x \sin 3x = \frac{1}{2}[\cos(-2x) - \cos 4x] = \frac{1}{2} \cos 2x - \frac{1}{2} \cos 4x$$

$$\begin{aligned} \sin x \sin 3x \sin 4x &= \frac{1}{2} \cos 2x \sin 4x - \frac{1}{2} \cos 4x \sin 4x \\ &= \frac{1}{2} \left[\frac{1}{2}(\sin 2x + \sin 6x) \right] - \frac{1}{2} \left[\frac{1}{2}(\sin 0 + \sin 8x) \right] \\ &= \frac{1}{4} \sin 2x + \frac{1}{4} \sin 6x - \frac{1}{4} \sin 8x \end{aligned}$$

$$\begin{aligned} \Rightarrow \int \sin x \sin 3x \sin 4x dx &= \int \left(\frac{1}{4} \sin 2x + \frac{1}{4} \sin 6x - \frac{1}{4} \sin 8x \right) dx \\ &= -\frac{1}{8} \cos 2x + \frac{1}{32} \cos 8x - \frac{1}{24} \cos 6x + c \end{aligned}$$

Synthesis

To evaluate the integral of the form $\int \sin mx \cos nxdx$ or $\int \cos mx \cos nxdx$ or $\int \sin mx \sin nxdx$, we express the product into sum by using the corresponding identities:

$$\sin A \cos B = \frac{1}{2}[\sin(A - B) + \sin(A + B)]$$

$$\sin A \sin B = \frac{1}{2}[\cos(A - B) - \cos(A + B)]$$

$$\cos A \cos B = \frac{1}{2}[\cos(A - B) + \cos(A + B)]$$

And then integrate the new expression.

Exercise 4.9 Learner's Book page 213

- | | |
|---------------------------------------------------|----------------------------------------------------|
| 1. $-\frac{1}{2}\cos x - \frac{1}{10}\cos 5x + c$ | 2. $\frac{1}{2}\cos x - \frac{1}{10}\cos 5x + c$ |
| 3. $-\frac{1}{12}\sin 6x + \frac{1}{2}x + c$ | 4. $\frac{1}{2}\sin^2 x + c$ |
| 5. $\frac{1}{12}\sin 6x + \frac{1}{2}x + c$ | 6. $\frac{1}{12}\sin 6x + \frac{1}{16}\sin 8x + c$ |

Lesson 4.10. Integrals of the form $\int \sin^m x \cos^n x dx$ **Learning objectives**

Given an integral of the form $\int \sin^m x \cos^n x dx$, learners should be able to find primitive function accurately.

Prerequisites

- ✔ Derivative of $\cos x$ and $\sin x$.
- ✔ Identity $\cos^2 x + \sin^2 x = 1$.
- ✔ Identity $\cos 2x = \cos^2 x - \sin^2 x$.
- ✔ Identity $\sin 2x = 2 \sin x \cos x$.

Teaching Aids

Exercise book and pen

Activity 4.10 Learner's Book page 213

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $u = \cos x \Rightarrow du = -\sin x dx$

$$\int \sin x \cos^2 x dx = \int \cos^2 x \sin x dx = -\int u^2 du = -\frac{u^3}{3} + c = -\frac{\cos^3 x}{3} + c$$

$$\begin{aligned}
 2. \quad \int \sin^2 x \cos^2 x dx &= \int \frac{1}{2}(1 - \cos 2x) \frac{1}{2}(1 + \cos 2x) dx \\
 &= \frac{1}{4} \int (1 + \cos 2x - \cos 2x - \cos^2 2x) dx \\
 &= \frac{1}{4} \int (1 - \cos^2 2x) dx \\
 &= \frac{1}{4} \int \left(1 - \frac{1}{2}(1 + \cos 4x) \right) dx, \\
 \text{since } \cos^2 x &= \frac{1}{2}(1 + \cos 2x) \Rightarrow \cos^2 2x = \frac{1}{2}(1 + \cos 4x) \\
 &= \frac{1}{4} \int \left(1 - \frac{1}{2} - \frac{1}{2} \cos 4x \right) dx \\
 &= \frac{1}{4} \int \left(\frac{1}{2} - \frac{1}{2} \cos 4x \right) dx = \frac{1}{8}x - \frac{1}{32} \sin 4x + c
 \end{aligned}$$

Synthesis

To integrate an integral of the form $\int \sin^m x \cos^n x dx$, we have two cases:

- If m or n is odd, save one cosine factor (or one sine factor) and use the relation $\cos^2 x = 1 - \sin^2 x$ (or $\sin^2 x = 1 - \cos^2 x$). Let $u = \sin x \Rightarrow du = \cos x dx$ (or let $u = \cos x \Rightarrow du = -\sin x dx$).
- If m and n are even, we use the identities:

$$\sin^2 x = \frac{1}{2}(1 - \cos 2x) \quad \text{and} \quad \cos^2 x = \frac{1}{2}(1 + \cos 2x).$$

Exercise 4.10 Learner's Book page 216

- $-\frac{1}{4} \cos^4 x + c$
- $\frac{1}{10} \sin^5 2x + c$
- $\frac{1}{3} \cos^3 x - \cos x + c$
- $-\frac{1}{12} \sin^3 4x + \frac{1}{4} \sin 4x + c$
- $\frac{1}{6} \cos^6 x - \frac{1}{4} \cos^4 x + c$
- $-\frac{1}{16} \cos^8 2x + \frac{1}{6} \cos^6 2x - \frac{1}{8} \cos^4 2x + c$

Lesson 4.11. Integrals of the form $\int \tan^m x \sec^n x dx$

Learning objectives

Given an integral of the form $\int \tan^m x \sec^n x dx$, learners should be able to find primitive function accurately.

Prerequisites

- ✔ Derivative of $\sec x$.
- ✔ Identity $\sec^2 x = 1 + \tan^2 x$.

Teaching Aids

Exercise book and pen

Activity 4.11 Learner's Book page 216



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad u = \tan x \Rightarrow du = (1 + \tan^2 x) dx = (1 + u^2) dx$$

$$\Rightarrow dx = \frac{du}{1 + u^2}$$

$$\begin{aligned} \int \tan^2 x dx &= \int \frac{u^2}{1 + u^2} du \\ &= \int \left(1 - \frac{1}{1 + u^2} \right) du \\ &= u - \arctan u + c \\ &= \tan x - \arctan(\tan x) + c \\ &= \tan x - x + c \end{aligned}$$

$$2. \quad u = \sec x \Rightarrow du = \sec x \tan x dx$$

$$dx = \frac{du}{\sec x \tan x} = \frac{du}{u \tan x}$$

$$\begin{aligned} \int \tan^5 x dx &= \int \tan^4 x \tan x dx = \int (\tan^2 x)^2 \tan x dx \\ &= \int (\sec^2 x - 1)^2 \tan x dx \\ &= \int (\sec^4 x - 2\sec^2 x + 1) \tan x dx \\ &= \int (u^4 - 2u^2 + 1) \tan x \frac{du}{u \tan x} \\ &= \int \frac{u^4 - 2u^2 + 1}{u} du \\ &= \int \left(u^3 - 2u + \frac{1}{u} \right) du \\ &= \frac{u^4}{4} - u^2 + \ln|u| + c = \frac{\sec^4 x}{4} - \sec^2 x + \ln|\sec x| + c \end{aligned}$$

Synthesis

Integration of the form $\int \tan^m x \sec^n x dx$, is in two types:

- If the power of secant is even, save a factor of $\sec^2 x$ and use $\sec^2 x = 1 + \tan^2 x$ to express the remaining factors in term of $\tan x$. Then substitute $u = \tan x$.
- If the power of tangent is odd, save a factor of $\sec x \tan x$ and use $\tan^2 x = \sec^2 x - 1$ to express the remaining factors in terms of $\sec x$. Then substitute $u = \sec x$.

Exercise 4.11 Learner's Book page 218

- $\frac{1}{2} \sec^2 x + c$
- $\frac{1}{4} \ln \left(\frac{1 - \sin x}{1 + \sin x} \right) + \frac{1}{2} \sec x \tan x + c$
- $\frac{1}{3} \sec^3 x + c$
- $-\frac{1}{3} \sec^3 x + \frac{1}{5} \sec^5 x + c$
- $\frac{1}{3} \tan^3 x + c$
- $\frac{1}{5} \tan^5 x + \frac{1}{3} \tan^3 x + c$

Lesson 4.12. Integrals containing $\sin x$, $\cos x$, $\tan x$ on denominator

Learning objectives

Given a function containing $\sin x$, $\cos x$, $\tan x$ on denominator, learners should be able to find primitive function moderately.

Prerequisites

- ✔ Identities

$$\sin x = \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \quad \cos x = \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \quad \tan x = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}}$$

from double angle formulae.

Teaching Aids

Exercise book and pen

Activity 4.12 Learner's Book page 218



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$\begin{aligned} \sin x + \cos x + 1 &= \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} + \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} + 1 \\ &= \frac{2 \tan \frac{x}{2} + 1 - \tan^2 \frac{x}{2} + 1 + \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}} = \frac{2 \tan \frac{x}{2} + 2}{1 + \tan^2 \frac{x}{2}} \end{aligned}$$

$$\frac{1}{\sin x + \cos x + 1} = \frac{1 + \tan^2 \frac{x}{2}}{2 \tan \frac{x}{2} + 2}$$

$$\text{Let } u = \tan \frac{x}{2} \Rightarrow \arctan u = \frac{x}{2} \Rightarrow \frac{du}{1+u^2} = \frac{dx}{2} \Rightarrow dx = \frac{2du}{1+u^2}$$

$$\begin{aligned} \int \frac{1}{\sin x + \cos x + 1} dx &= \int \frac{1+u^2}{2u+2} \times \frac{2du}{1+u^2} \\ &= \int \frac{du}{u+1} \\ &= \ln|u+1| + c \\ &= \ln \left| \tan \frac{x}{2} + 1 \right| + c \end{aligned}$$

Synthesis

To find an integral containing $\sin x$, $\cos x$, $\tan x$ on denominator, use the formulae

$$\sin x = \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \quad \cos x = \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}, \quad \tan x = \frac{2 \tan \frac{x}{2}}{1 - \tan^2 \frac{x}{2}}$$

$$\text{and let } u = \tan \frac{x}{2} \Rightarrow \arctan u = \frac{x}{2}$$

Exercise 4.12 Learner's Book page 220

1. $\frac{2}{\sqrt{3}} \arctan \left(\frac{1}{\sqrt{3}} \tan \frac{x}{2} \right) + c$
2. $\frac{2}{\sqrt{5}} \arctan \left(\frac{3 \tan \frac{x}{2} - 2}{\sqrt{5}} \right) + c$
3. $\frac{1}{\sqrt{3}} \arctan \left(\frac{2 \tan x + 1}{\sqrt{3}} \right) + c$
4. $\frac{1}{2} \arctan \left(2 \tan \frac{x}{2} \right) + c$
5. $\frac{2}{\sqrt{11}} \arctan \left(\frac{3 \tan \frac{x}{2} + 4}{\sqrt{11}} \right) + c$
6. $-\frac{1}{\tan \frac{x}{2} - 2} + c$

Lesson 4.13. Integrals containing $\sin^2 x, \cos^2 x$ on denominator

Learning objectives

Given an integral containing $\sin^2 x, \cos^2 x$ on denominator, learners should be able to find primitive function accurately.

Prerequisites

⑦ Identities $\cos x = \frac{1}{\sqrt{1+\tan^2 x}}$ and $\sin x = \frac{\tan x}{\sqrt{1+\tan^2 x}}$.

Teaching Aids

Exercise book and pen

Activity 4.13 Learner's Book page 220



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ⑦ Critical thinking
- ⑦ Communication
- ⑦ Self confidence
- ⑦ Cooperation, interpersonal management and life skills
- ⑦ Peace and values education
- ⑦ Inclusive education

Answers

We know that $\sec^2 x = 1 + \tan^2 x$ or $\sec x = \pm\sqrt{1 + \tan^2 x}$.
Then,

$$\cos x = \pm \frac{1}{\sqrt{1 + \tan^2 x}} \quad \text{and} \quad \sin x = \pm \frac{\tan x}{\sqrt{1 + \tan^2 x}}$$

$$\text{Now, } \frac{1}{\cos^2 x} = \frac{1}{\left(\pm \frac{1}{\sqrt{1 + \tan^2 x}}\right)^2} = \frac{1}{\frac{1}{1 + \tan^2 x}} = 1 + \tan^2 x$$

$$\int \frac{1}{\cos^2 x} dx = \int (1 + \tan^2 x) dx$$

$$\text{Let } u = \tan x \Rightarrow x = \arctan u \Rightarrow dx = \frac{du}{1 + u^2}$$

$$\int \frac{1}{\cos^2 x} dx = \int (1+u^2) \frac{du}{1+u^2} = \int du = u + c = \tan x + c$$

Synthesis

To integrate an integral containing $\sin^2 x$, $\cos^2 x$ on denominator, use identities $\cos x = \frac{1}{\sqrt{1+\tan^2 x}}$ and $\sin x = \frac{\tan x}{\sqrt{1+\tan^2 x}}$ and let $u = \tan x \Rightarrow x = \arctan u$

Exercise 4.13 Learner's Book page 222

- 1) $\frac{1}{3} \tan^3 x + \tan x + c$ 2) $\frac{1}{5} \tan^5 x + \frac{2}{3} \tan^3 x + \tan x + c$
 3) $-\cot x - \frac{2}{3} \cot^3 x - \frac{1}{5} \cot^5 x + c$ 4) $\frac{1}{7} \tan^7 x + \frac{3}{5} \tan^5 x + \tan^3 x + \tan x + c$

Lesson 4.14. Integrals containing $\sqrt[n]{ax+b}$

Learning objectives

Given an integral containing $\sqrt[n]{ax+b}$, learners should be able to find primitive function accurately.

Prerequisites

- ✔ Properties of radicals.

Teaching Aids

Exercise book and pen



Activity 4.14 Learner's Book page 223

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$u^2 = 3x - 1 \Rightarrow u = \sqrt{3x - 1}$$

$$\Rightarrow 2udu = 3dx \Rightarrow dx = \frac{2udu}{3}$$

$$\int \sqrt{3x - 1} dx = \int u \frac{2udu}{3} = \frac{2}{3} \int u^2 du$$

$$= \frac{2u^3}{9} + c = \frac{2}{9} (\sqrt{3x - 1})^3 + c$$

$$= \frac{2}{9} (\sqrt{3x - 1})^2 \sqrt{3x - 1} + c = \frac{2}{9} (3x - 1) \sqrt{3x - 1} + c$$

Synthesis

For integral containing $\sqrt[n]{ax + b}$, $a \neq 0$, let $u^n = ax + b$

Exercise 4.14 Learner's Book page 224

$$1. \quad \frac{1}{3} (2x + 1) \sqrt{6x + 3} + c$$

$$2. \quad \frac{2}{25} (5x - 2)^2 \sqrt{5x - 2} + c$$

$$3. \quad \frac{3}{16} \sqrt[3]{(8x + 1)^2} + c$$

$$4. \quad \frac{2}{3\sqrt{2 - 3x}} + c$$

$$5. \quad \frac{1}{3} \sqrt{(2x + 5)^3} + c$$

$$6. \quad \frac{1}{4} (3x - 8) \sqrt{3x - 8} + c$$

$$7. \quad 4\sqrt{2x + 3} + c$$

$$8. \quad \frac{4}{3\sqrt{1 - 3x}} + c$$

Lesson 4.15. Integrals containing $\sqrt{ax^2 + bx + c}$ **Learning objectives**

Given an integral containing $\sqrt{ax^2 + bx + c}$, learners should be able to find primitive function accurately.

Prerequisites

⑦ Use of the relation $ax^2 + bx + c = a \left[\left(x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2} \right]$.

Teaching Aids

Exercise book and pen



Activity 4.15 Learner's Book page 225

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

The relation to be used is

$$ax^2 + bx + c = a \left[\left(x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2} \right]$$

$$1. \int \frac{dx}{\sqrt{x^2 - 2x + 1}}$$

$$x^2 - 2x + 1 = (x - 1)^2 - \frac{(-2)^2 - 4 \times 1}{4} = (x - 1)^2$$

$$\text{Let } u = x - 1 \Rightarrow du = dx$$

$$\int \frac{dx}{\sqrt{x^2 - 2x + 1}} = \int \frac{du}{\sqrt{u^2}} = \int \frac{du}{u}$$

Using formula $\int \frac{u'}{u} du = \ln|u| + d$, we have

$$\int \frac{dx}{\sqrt{x^2 - 2x + 1}} = \int \frac{du}{u} = \ln|u| + d = \ln|x - 1| + d$$

$$2. \int \frac{dx}{\sqrt{x^2 - 5x + 6}}$$

$$x^2 - 5x + 6 = \left(x - \frac{5}{2} \right)^2 - \frac{(-5)^2 - 4 \times 6}{4} = \left(x - \frac{5}{2} \right)^2 - \frac{1}{4}$$

$$\text{Let } u = x - \frac{5}{2} \Rightarrow du = dx$$

$$\int \frac{dx}{\sqrt{x^2 - 5x + 6}} = \int \frac{du}{\sqrt{u^2 - \frac{1}{4}}}$$

Using formula $\int \frac{dx}{\sqrt{x^2 \pm k^2}} = \ln \left| x + \sqrt{x^2 \pm k^2} \right| + d$,
we have

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - 5x + 6}} &= \int \frac{du}{\sqrt{u^2 - \frac{1}{4}}} \\ &= \ln \left| u + \sqrt{u^2 - \frac{1}{4}} \right| + d = \ln \left| x - \frac{5}{2} + \sqrt{\left(x - \frac{5}{2}\right)^2 - \frac{1}{4}} \right| + d \\ &= \ln \left| \frac{2x - 5 + 2\sqrt{x^2 - 5x + 6}}{2} \right| + d \\ &= \ln \left| 2x - 5 + 2\sqrt{x^2 - 5x + 6} \right| + e \end{aligned}$$

$$\begin{aligned} 3. \int \frac{dx}{\sqrt{x^2 - 6x + 18}} \\ x^2 - 6x + 18 &= (x - 3)^2 - \frac{(-6)^2 - 4 \times 16}{4} \\ &= (x - 3)^2 + 9 \end{aligned}$$

Let $u = x - 3 \Rightarrow du = dx$

$$\int \frac{dx}{\sqrt{x^2 - 6x + 18}} = \int \frac{du}{\sqrt{u^2 + 9}}$$

Using formula $\int \frac{dx}{\sqrt{x^2 \pm k^2}} = \ln \left| x + \sqrt{x^2 \pm k^2} \right| + d$,
we have

$$\begin{aligned} \int \frac{dx}{\sqrt{x^2 - 6x + 18}} &= \int \frac{du}{\sqrt{u^2 + 9}} \\ &= \ln \left| u + \sqrt{u^2 + 9} \right| + d \\ &= \ln \left| x - 3 + \sqrt{(x - 3)^2 + 9} \right| + d \\ &= \ln \left| x - 3 + \sqrt{x^2 - 6x + 18} \right| + d \end{aligned}$$

Synthesis

For the integrals of the form $\int \frac{dx}{\sqrt{ax^2 + bx + c}}$, first, transform $ax^2 + bx + c$ in the form

$$a \left[\left(x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2} \right] \text{ and then;}$$

① If $b^2 - 4ac = 0$, then

$$\int \frac{dx}{\sqrt{ax^2 + bx + c}} = \frac{1}{\sqrt{a}} \int \frac{dx}{x + \frac{b}{2a}} \text{ and we let } u = x + \frac{b}{2a}$$

② If $b^2 - 4ac > 0$, then

$$\int \frac{dx}{\sqrt{ax^2 + bx + c}} = \frac{1}{\sqrt{a}} \int \frac{dx}{\sqrt{\left(x + \frac{b}{2a} \right)^2 + \frac{b^2 - 4ac}{4a^2}}}$$

We let $u = x + \frac{b}{2a}$, $k^2 = \frac{b^2 - 4ac}{4a^2}$ finally, use the integral

$$\int \frac{dx}{\sqrt{x^2 \pm k^2}} = \ln \left| x + \sqrt{x^2 \pm k^2} \right| + d$$

③ If $b^2 - 4ac < 0$, then

$$\int \frac{dx}{\sqrt{ax^2 + bx + c}} = \frac{1}{\sqrt{a}} \int \frac{dx}{\sqrt{\left(x + \frac{b}{2a} \right)^2 - \frac{b^2 - 4ac}{4a^2}}}$$

We let $u = x + \frac{b}{2a}$, $k^2 = -\frac{b^2 - 4ac}{4a^2}$ and use the integral

$$\int \frac{dx}{\sqrt{x^2 \pm k^2}} = \ln \left| x + \sqrt{x^2 \pm k^2} \right| + d$$

Exercise 4.15 Learner's Book page 232

1. $\ln \left| x + 1 + \sqrt{x^2 + 2x + 5} \right| + c$ 2. $\arcsin \frac{x+1}{\sqrt{5}} + c$

3. $\ln \left| x + 2 + \sqrt{x^2 + 4x + 2} \right| + c$ 4. $\arcsin \frac{x-3}{2} + c$

$$5. \frac{2x-1}{4}\sqrt{x-x^2} + \frac{1}{8}\arcsin(2x-1) + c$$

Lesson 4.16. Integration by parts

Learning objectives

Through examples, learners should be able to integrate by parts accurately.

Prerequisites

Product rule differentiation.

Teaching Aids

Exercise book and pen

Activity 4.16 Learner's Book page 232



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$1. \frac{d}{dx} f(x) = e^x \frac{d}{dx}(x-1) + (x-1) \frac{d}{dx} e^x \\ = e^x + (x-1)e^x = xe^x$$

$$2. \text{From 1), } \int xe^x dx = (x-1)e^x + c$$

$$3. \text{Let } u = x, v = e^x,$$

$$\text{Thus, } \int uv dx = \int xe^x dx \text{ while } \int u dx \int v dx = \int x dx \int e^x dx$$

From 2) we have

$$\int uv dx = \int xe^x dx = (x-1)e^x + c$$

Let us find $\int u dx \int v dx$:

$$\int u dx \int v dx = \int x dx \int e^x dx = \frac{x^2}{2} e^x + c$$

Therefore, $\int uv dx \neq \int u dx \int v dx$

Synthesis

Integration by parts use the formula $\int u dv = uv - \int v du$

The following table can be used:

u	v'
Logarithmic function	Polynomial function
Polynomial function	Exponential function
Polynomial function	Trigonometric function
Exponential function	Trigonometric function
Trigonometric function	Exponential function
Inverse trigonometric function	Polynomial function

Exercise 4.16 Learner's Book page 235

- $\frac{1}{2}x \sin 2x + \frac{1}{4} \cos 2x + c$
- $\frac{1}{9}(3x-1)e^{3x} + c$
- $\frac{1}{16} \sin 4x - \frac{1}{4}x \cos 4x + c$
- $\frac{1}{9}x^3(3 \ln x - 1) + c$
- $(x+1)e^{2x} + c$
- $-\frac{1}{4}(2x+1)e^{-2x} + c$

Lesson 4.17. Integration by reduction formulae

Learning objectives

Given integral I_m and by using integration by parts, learners should be able to find a reduction formula for I_m rightfully.

Prerequisites

- ☑ Integration by parts.

Teaching Aids

Exercise book and pen

**Activity 4.17 Learner's Book page 236**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$I_m = \int x^m \cos bx \, dx, J_m = \int x^m \sin bx \, dx$$

For $J_m = \int x^m \sin bx \, dx$, let $u = x^m \Rightarrow du = mx^{m-1} dx$

$$dv = \sin bx \, dx \Rightarrow v = -\frac{1}{b} \cos bx$$

$$J_m = -\frac{x^m \cos bx}{b} - \int \left(-\frac{1}{b} \cos bx\right) mx^{m-1} dx$$

$$\Rightarrow J_m = -\frac{x^m \cos bx}{b} + \frac{m}{b} \underbrace{\int \cos bx x^{m-1} dx}_{I_{m-1}}$$

$$\Rightarrow J_m = -\frac{x^m \cos bx}{b} + \frac{m}{b} I_{m-1}$$

$$\Rightarrow bJ_m = -x^m \cos bx + mI_{m-1} \Rightarrow bJ_m - mI_{m-1} = -x^m \cos bx$$

Synthesis

Knowing integral I_m , we can establish a general relation, integration by parts, which will help us to reduce the power and find $I_{m-1}, I_{m-2}, I_{m-3}, \dots, I_0$.

Exercise 4.17 Learner's Book page 238

$$1. \frac{1}{a} x^n e^{ax} - \frac{n}{a} I_{n-1}$$

$$2. I_n = \frac{\tan^{n-1} x}{n-1} - I_{n-2} \text{ and}$$

$$\int \tan^5 x \, dx = I_5 = \frac{\tan^4 x}{4} - \frac{\tan^2 x}{2} + \ln |\sec x| + c$$

$$3. I_n = -\frac{1}{n} \sin^{n-1} x \cos x + \frac{n-1}{n} I_{n-2}$$

$$4. I_n = \frac{1}{n} \cos^{n-1} x \sin x + \frac{n-1}{n} I_{n-2}$$

$$5. I_n = x(\ln x)^n - nI_{n-1}$$

Lesson 4.18. Integration by Maclaurin series**Learning objectives**

Using Maclaurin series, learners should be able to find primitive functions of some functions accurately.

Prerequisites

✔ Maclaurin series of a function.

Teaching Aids

Exercise book and pen

**Activity 4.18 Learner's Book page 238**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \dots$$

$$2. \int \ln(1+x) dx = \int \left(x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \dots \right) dx$$

$$= \frac{1}{2}x^2 - \frac{1}{6}x^3 + \frac{1}{12}x^4 - \frac{1}{20}x^5 + \dots + c$$

Synthesis

For some integrals, we proceed also by Maclaurin series of the function to be integrated.

Exercise 4.18 Learner's Book page 239

$$1. \int e^{-3x} dx = \int \left(1 - 3x + \frac{9}{2}x^2 - \frac{9}{2}x^3 + \dots \right) dx = x - \frac{3}{2}x^2 + \frac{3}{2}x^3 - \frac{9}{8}x^4 + \dots + c$$

$$2. \int \sin x dx = \int \left(x - \frac{1}{6}x^3 + \frac{1}{120}x^5 + \dots \right) dx = \frac{1}{2}x^2 - \frac{1}{24}x^4 + \frac{1}{720}x^6 + \dots + c$$

$$3. \int \cos x dx = \int \left(1 - \frac{1}{2}x^2 + \frac{1}{24}x^4 - \frac{1}{720}x^6 + \dots \right) dx = x - \frac{1}{6}x^3 + \frac{1}{120}x^5 - \frac{1}{5040}x^7 + \dots + c$$

$$4. \int \tan x dx = \int \left(x + \frac{1}{3}x^3 + \frac{2}{15}x^5 + \dots \right) dx = \frac{1}{2}x^2 + \frac{1}{12}x^4 + \frac{1}{45}x^6 + \dots + c$$

$$5. \int \sqrt{1+x} dx = \int \left(1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 + \dots \right) dx = x + \frac{1}{4}x^2 - \frac{1}{24}x^3 + \frac{1}{64}x^4 + \dots + c$$

Lesson 4.19. Definite integrals

Learning objectives

By the end of this lesson, learners should be able to define definite integrals.

Prerequisites

- ⑦ Indefinite integrals.

Teaching Aids

Exercise book, calculator and pen



Activity 4.19 Learner's Book page 239

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. F(x) = \int (x^2 - 2x + 3) dx = \frac{1}{3}x^3 - x^2 + 3x + c$$

$$\begin{aligned}
 2. F(1) - F(-1) &= \left[\frac{1}{3}(1)^3 - (1)^2 + 3(1) + c \right] \\
 &\quad - \left[\frac{1}{3}(-1)^3 - (-1)^2 + 3(-1) + c \right] \\
 &= \left(\frac{1}{3} - 1 + 3 + c \right) - \left(-\frac{1}{3} - 1 - 3 + c \right) \\
 &= \frac{1}{3} + 2 + c + \frac{1}{3} + 4 - c \\
 &= \frac{20}{3}
 \end{aligned}$$

Synthesis

The area of the strip between x_{i-1} and x_i is approximately equal to the area of a rectangle with width $l = \Delta x$ and length $L = f(x_i)$ i.e. as illustrated in figure 4.1.

The total area A is $\sum_{i=1}^n S_i = \sum_{i=1}^n f(x_i) \cdot \Delta x$ or

$A = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \cdot \Delta x$; this is known as **Sum of Riemann**.




Figure 4.1: Definite integral of the function

We define the definite integrals of the function $f(x)$ with respect to x from a to b to be

$\int_a^b f(x) dx = [F(x)]_a^b = F(b) - F(a)$. Where $F(x)$ is the anti-derivative of $f(x)$.

Exercise 4.19 Learner's Book page 242

- | | | | |
|-------------------|------------------|---------|------------------|
| 1. $\frac{9}{2}$ | 2. $\frac{5}{6}$ | 3. -2 | 4. $\frac{3}{4}$ |
| 5. $\frac{46}{3}$ | 6. 4 | 7. 25 | 8. 145 |

Lesson 4.20. Properties of definite integrals

Learning objectives

Through examples, learners should be able to use properties of definite integrals accurately.

Prerequisites

- ⑦ Definition of definite integrals.

Teaching Aids

Exercise book, calculator and pen



Activity 4.20 Learner's Book page 243

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \int_{-3}^0 f(x) dx = \int_{-3}^0 x^2 dx = \left[\frac{x^3}{3} \right]_{-3}^0 = 9$$

$$\int_0^{-3} f(x) dx = \int_0^{-3} x^2 dx = \left[\frac{x^3}{3} \right]_0^{-3} = -9$$

$$\text{Observation: } \int_{-3}^0 f(x) dx = -\int_0^{-3} f(x) dx$$

$$2. \int_{-2}^2 x^2 dx = \left[\frac{1}{3} x^3 \right]_{-2}^2 = \frac{8}{3} + \frac{8}{3} = \frac{16}{3}$$

$$\int_{-2}^0 x^2 dx + \int_0^2 x^2 dx = \left[\frac{1}{3} x^3 \right]_{-2}^0 + \left[\frac{1}{3} x^3 \right]_0^2 = 0 + \frac{8}{3} + \frac{8}{3} - 0 = \frac{16}{3}$$

Observation:

$$\int_{-2}^2 x^2 dx = \int_{-2}^0 x^2 dx + \int_0^2 x^2 dx$$

Synthesis

- ⑦ **Permutation of bounds:** If $f(x)$ is defined on (a, b) except may be at a finite number of points, then

$$\int_a^b f(x)dx = -\int_b^a f(x)dx$$

- ⑦ **Chasles relation:** For any arbitrary numbers a and b and any $c \in [a, b]$

$$\int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$

- ⑦ **Positivity:** Let f be a continuous function on interval $I = [a, b]$ the elements of I

If $f \geq 0$ on I and if $a \leq b$ then $\int_a^b f(x)dx \geq 0$

Also, if $f(x) \leq g(x)$ on $[a, b]$, then,

$$\int_a^b f(x)dx \leq \int_a^b g(x)dx$$

Exercise 4.20 Learner's Book page 245

1. 4 2. $\frac{7}{3}$ 3. $\frac{1}{32}\pi^2$ 4. $\frac{1}{2}\pi - 1$

Lesson 4.21. Improper integrals, Infinite limits of integration**Learning objectives**

Given an improper integral with infinite limits, learners should be able to determine whether it converges or diverges correctly.

Prerequisites

- ⑦ Limits concepts.

Teaching Aids

Exercise book and pen



Activity 4.21 Learner's Book page 245

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$\begin{aligned}
 1. \quad \lim_{n \rightarrow +\infty} \int_0^n \frac{dx}{x^2 + 4} &= \lim_{n \rightarrow +\infty} \left[\frac{1}{2} \arctan \left(\frac{1}{2}x \right) \right]_0^n \\
 &= \lim_{n \rightarrow +\infty} \left[\frac{1}{2} \arctan \left(\frac{1}{2}n \right) \right] \\
 &= \frac{1}{4} \pi
 \end{aligned}$$

$$\begin{aligned}
 2. \quad \lim_{n \rightarrow -\infty} \int_n^{-4} \frac{xdx}{\sqrt{1+3x^2}} &= \lim_{n \rightarrow -\infty} \left[\frac{1}{3} \sqrt{3x^2+1} \right]_n^{-4} \\
 &= \lim_{n \rightarrow -\infty} \frac{-\sqrt{3n^2+1}+7}{3} \\
 &= -\infty
 \end{aligned}$$

Synthesis

We define the improper integral as

$$\int_a^{+\infty} f(x)dx = \lim_{n \rightarrow +\infty} \int_a^n f(x)dx \quad \text{or} \quad \int_{-\infty}^b f(x)dx = \lim_{n \rightarrow -\infty} \int_n^b f(x)dx$$

If the limit exists, we say that the integral converges, otherwise it diverges.

Exercise 4.21 Learner's Book page 247

- | | |
|-----------------------------------|--------------------------------|
| 1) Convergent to π | 2) Convergent to $\frac{1}{2}$ |
| 3) Convergent to $\frac{1}{4}\pi$ | 4) Convergent to $\frac{1}{2}$ |
| 5) divergent | 6) Convergent to 0 |

Lesson 4.22. Discontinuous integrand**Learning objectives**

Given an improper integral with discontinuous integrand, learners should be able to determine whether it converges or diverges correctly.

Prerequisites

- ✔ Limits concepts.

Teaching Aids

Exercise book, calculator and pen

Activity 4.22 Learner's Book page 247

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Since this function is a rational function, the denominator cannot be zero. Then, $x - 1 \neq 0$ or $x \neq 1$. Thus, considering the given interval, the given function is discontinuous at $x = 1$.

2. Since this function is a rational function, the denominator cannot be zero. Then, $x^2 - 3x - 10 \neq 0$ or $x \neq -2$ and $x \neq 5$. Thus, considering the given interval, the given function is discontinuous at $x = -2$.
3. Since there is natural logarithm, then, $x > 0$ also $\ln x \neq 0$ or $x \neq 1$. Thus, considering the given interval, the given function is discontinuous at $x = 1$.

Synthesis

For a function $f(x)$ which is continuous on the interval $[a, b[$, we define the improper integral as

$$\int_a^b f(x) dx = \lim_{t \rightarrow b^-} \int_a^t f(x) dx.$$

Also, if $f(x)$ is continuous on the interval $]a, b]$, we

have the improper integral $\int_a^b f(x) dx = \lim_{t \rightarrow a^+} \int_t^b f(x) dx.$

If $f(x)$ is a continuous function for all real numbers x in the interval $]a, b[$, except for some point $c \in]a, b[$, then,

$$\begin{aligned} \int_a^b f(x) dx &= \int_a^c f(x) dx + \int_c^b f(x) dx \\ &= \lim_{t \rightarrow c^-} \int_a^t f(x) dx + \lim_{t \rightarrow c^+} \int_t^b f(x) dx \end{aligned}$$

Exercise 4.22 Learner's Book page 249

1. Diverges
2. a) converges to 3 b) diverges
- c) diverges d) converges to $5 + 5\sqrt{2}$

Lesson 4.23. Calculation of area of plane surfaces

Learning objectives

Given two functions and by using integration, learners should be able to find the area between two curves in a given interval precisely.

Prerequisites

- ✔ Curve sketching.
- ✔ Definite integrals.

Teaching Aids

Exercise book, calculator and pen

Activity 4.23 Learner's Book page 249




In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education


Answers

1. Curve

x	0	4
$f(x)$	0	4



2. Curve with shaded region



3. We see that the shaded region is a triangle whose base is 4 units and height is 4 units. Then, we know the area of a triangle with base B and height H is

$$A = \frac{B \times H}{2} .$$

Then, the area of the shaded region is $A = \frac{4 \times 4}{2} = 8 \text{ sq. units} .$

$$\begin{aligned}
 4. \quad \int_0^4 f(x) dx &= \int_0^4 x dx = \left[\frac{x^2}{2} \right]_0^4 \\
 &= \frac{4^2}{2} \\
 &= 8
 \end{aligned}$$

5. Results in 3) and 4) are the same.

Synthesis

Given function $f(x)$ which lies above the x -axis, the area enclosed by the curve of $f(x)$ and x -axis in interval $[a, b]$ is given by;

$$A = \int_a^b f(x) dx$$




Figure 4.2: Area enclosed by a curve of a function and x -axis

The area between two functions $f(x)$ and $g(x)$ where $f(x) \leq g(x)$ in $[a, b]$ is given by

$$\int_a^b [g(x) - f(x)] dx = \int_a^b g(x) dx - \int_a^b f(x) dx$$





Figure 4.3: Area between two curves


Exercise 4.23 Learner's Book page 255

1. a) $\frac{2}{3}$ sq. units b) $\frac{1}{2}$ sq. units c) $33\frac{1}{3}$ sq. units
 d) $\frac{4}{3}$ sq. units e) 9 sq. units
- 2) $\frac{16}{15}a^5$ sq. units 3) $143\frac{5}{6}$ sq. units
- 4) 3.75 sq. units 5) $4\sqrt{2}$ sq. units
- 6) a) Graph



Area is $\frac{32}{3}$ sq. units

b) Graph



Area is $\frac{355}{6}$ sq. units

Lesson 4.24. Calculation of volume of a solid of revolution

Learning objectives

Given a function and by using integration, learners should be able to find the volume of a solid obtained when a curve of a function is revolved around an axis precisely.

Prerequisites

- ✔ Curve sketching.
- ✔ Definite integrals.

Teaching Aids

Exercise book, calculator and pen

Activity 4.24 Learner's Book page 255




In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education


Answers

1. $y = 2$ for $0 \leq x \leq 3$

- a) The region enclosed by the curve $y = 2$ for $0 \leq x \leq 3$ and x -axis



- b) The region for which the area in (a) is rotated 360° about the x-axis



- c) Solid of revolution obtained in (b) is a cylinder of radius 2 and height 3.

Volume of cylinder is

$$\pi r^2 h = \pi (2)^2 (3) = 12\pi \text{ cubic units}$$


- d) Volume $V = \lim_{\delta x \rightarrow 0} \sum_{x=0}^{x=3} \pi y^2 \delta x = \int_0^3 \pi y^2 dx$

$$= \int_0^3 \pi (2)^2 dx = \int_0^3 4\pi dx = 4\pi [x]_0^3 = 12\pi \text{ cubic units}$$


- e) The results obtained in (c) and (d) are equal.

2. $y = 2x$ for $0 \leq x \leq 5$

- a) The region enclosed by the curve $y = 2x$ for $0 \leq x \leq 5$ and x -axis



- b) The region for which the area in (a) is rotated 360° about the x -axis



- c) Solid of revolution obtained in (b) is a cone of radius 10 and height 5.

Volume of cone is

$$\frac{1}{3}\pi r^2 h = \frac{1}{3}\pi (10)^2 (5) = \frac{500}{3}\pi \text{ cubic units}$$

- d) Volume $V = \lim_{\delta x \rightarrow 0} \sum_{x=0}^{x=5} \pi y^2 \delta x = \int_0^5 \pi y^2 dx$

$$= \int_0^5 \pi (2x)^2 dx = \int_0^5 4\pi x^2 dx$$

$$= 4\pi \left[\frac{x^3}{3} \right]_0^5 = \frac{500}{3}\pi \text{ cubic units}$$

- e) The results obtained in (c) and (d) are equal.

Synthesis

The volume of the solid of revolution bound by the curve $f(x)$ about the x -axis calculated from $x = a$ to $x = b$,

is given $V = \pi \int_a^b f^2(x) dx$.




Figure 4.4: Volume of revolution

Exercise 4.24 Learner's Book page 263

1. a) $\frac{32\pi}{5}$ cubic units b) $\frac{373\pi}{14}$ cubic units
 c) $\frac{1296\pi}{5}$ cubic units
2. a) $\frac{3\pi}{5}$ cubic units b) 8π cubic units
 c) 2π cubic units

Exercise 4.25 Learner's Book page 268

- 1) $\frac{1}{243}(85\sqrt{85} - 8)$ cubic units
- 2) $\frac{1}{27}(80\sqrt{10} - 13\sqrt{13})$ cubic units
- 3) $\frac{17}{6}$ cubic units 4) $\frac{1}{27}(13\sqrt{13} + 80\sqrt{10} - 16)$ cubic units

Lesson 4.25. Calculation of arc length of a curved surface

Learning objectives

Given a function and by using integration, learners should be able to find arc length of a curve in a given interval precisely.

Prerequisites

- ✔ Curve sketching.
- ✔ Definite integrals.

Teaching Aids

Exercise book, calculator and pen

Activity 4.25 Learner's Book page 269



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers




Figure 4.5: Arc length of a curved line

$$1. (\Delta l)^2 = (\Delta x)^2 + (\Delta y)^2$$

$$\Delta l = \sqrt{(\Delta x)^2 + (\Delta y)^2}$$

$$2. \Delta l = \sqrt{(\Delta x)^2 + (\Delta y)^2} \Rightarrow dl = \sqrt{(dx)^2 + (dy)^2}$$

$$dl = \sqrt{(dx)^2 + (dy)^2}$$

$$= \sqrt{(dx)^2 \left(1 + \frac{(dy)^2}{(dx)^2} \right)}$$

$$= \sqrt{1 + \left(\frac{dy}{dx} \right)^2} dx$$

We recognise the ratio inside the square root as the derivative, $\frac{dy}{dx} = f'(x)$, then we can rewrite this as

$$dl = \sqrt{1 + [f'(x)]^2} dx$$

But $f(x) = (x-1)^{\frac{3}{2}} \Rightarrow f'(x) = \frac{3}{2}(x-1)^{\frac{1}{2}}$, then,

$$dl = \sqrt{1 + \left[\frac{3}{2}(x-1)^{\frac{1}{2}} \right]^2} dx$$

$$= \sqrt{1 + \frac{9}{4}(x-1)} dx$$

$$= \sqrt{\frac{4+9x-9}{4}} dx$$

$$= \sqrt{\frac{9x-5}{4}} dx$$

$$3. \int dl = \int_2^5 \sqrt{\frac{9x-5}{4}} dx \Rightarrow l = \int_2^5 \sqrt{\frac{9x-5}{4}} dx$$

$$= \frac{1}{2} \int_2^5 \sqrt{9x-5} dx$$

$$\text{But } \frac{1}{2} \int_2^5 \sqrt{9x-5} dx = \left[\frac{1}{27} (9x-5)^{\frac{3}{2}} \right]_2^5$$

Then,

$$\begin{aligned}
 l &= \frac{1}{27} \left[(9x-5)^{\frac{3}{2}} \right]_2^5 \\
 &= \frac{1}{27} \left((45-5)^{\frac{3}{2}} - (18-5)^{\frac{3}{2}} \right) \\
 &= \frac{1}{27} \left((40)^{\frac{3}{2}} - (13)^{\frac{3}{2}} \right) \\
 &= \frac{1}{27} \left(\sqrt{(40)^3} - \sqrt{(13)^3} \right) \\
 &= \frac{1}{27} \left(\sqrt{40 \times (40)^2} - \sqrt{13 \times (13)^2} \right) \\
 &= \frac{1}{27} \left(\sqrt{4 \times 10 \times (40)^2} - \sqrt{13 \times (13)^2} \right) \\
 &= \frac{1}{27} \left(80\sqrt{10} - 13\sqrt{13} \right) \text{ units of length}
 \end{aligned}$$

Synthesis

Arc length of a curve of function $f(x)$ in interval $]a, b[$ is given by $L = \int_a^b \sqrt{1 + [f'(x)]^2} dx$.

Exercise 4.26 Learner's Book page 272

- 1) $\frac{85\sqrt{85} - 8}{243}$ units of length
- 2) $\frac{80\sqrt{10} - 13\sqrt{13}}{27}$ units of length
- 3) $\ln(\sqrt{2} + 1)$ units of length
- 4) $\frac{14}{3}$ units of length

Summary of the unit

1. Differentials

The exact change, Δy , in y is given by

$$\Delta y = f(x + \Delta x) - f(x).$$

But if the change Δx is small, then we can get a good approximation to Δy by using the fact that $\frac{\Delta y}{\Delta x}$ is approximately the derivative $\frac{dy}{dx}$. Thus,

$$\Delta y = \frac{\Delta y}{\Delta x} \Delta x \approx \frac{dy}{dx} \Delta x = f'(x) \Delta x$$

If we denote the change of x by dx instead of Δx , then the change Δy in y is approximated by the differential dy , that is, $\Delta y \approx dy = f'(x) dx$

Whenever one makes an approximation, it is wise to try and estimate how big the error might be.

Relative change in x is $\frac{\Delta x}{x}$

Percentage change in x is $100 \times \frac{\Delta x}{x}$

2. Indefinite integrals

Integration can be defined as the inverse process of differentiation.

If $y = f(x)$ then

$$\frac{dy}{dx} = f'(x) \Leftrightarrow \int \frac{dy}{dx} dx = f(x) + c$$

Or equivalently

$$\int \frac{dy}{dx} dx = y + c$$

This is called indefinite integration and c is the constant of integration.

3. Basic integration formula

Exponential functions

$$1. \int x^n dx = \frac{x^{n+1}}{n+1} + c, n \neq -1$$

$$2. \int e^x dx = e^x + c$$

$$3. \int a^x dx = \frac{a^x}{\ln a} + c$$

Rational functions

$$1. \int \frac{1}{x} dx = \ln|x| + c$$

$$2. \int \frac{dx}{a^2 + x^2} = \frac{1}{a} \arctan \frac{x}{a} + c$$

$$3. -\int \frac{dx}{a^2 + x^2} = \frac{1}{a} \operatorname{arccot} \frac{x}{a} + c$$

$$4. \int \frac{dx}{x^2 - a^2} = \frac{1}{2a} \ln \left| \frac{x-a}{x+a} \right| + c$$

$$5. \int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \ln \left| \frac{a+x}{a-x} \right| + c$$

Irrational functions

$$1. \int \frac{dx}{\sqrt{a^2 - x^2}} = \arcsin \frac{x}{a} + c$$

$$2. -\int \frac{dx}{\sqrt{a^2 - x^2}} = \arccos \frac{x}{a} + c$$

$$3. \int \frac{dx}{\sqrt{x^2 + a^2}} = \ln \left| \frac{x + \sqrt{x^2 + a^2}}{a} \right| + c$$

$$4. \int \frac{dx}{\sqrt{x^2 - a^2}} = \ln \left| \frac{x + \sqrt{x^2 - a^2}}{a} \right| + c$$

$$5. \int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \operatorname{arcsec} \frac{x}{a} + c$$

$$6. -\int \frac{dx}{x\sqrt{x^2 - a^2}} = \frac{1}{a} \operatorname{arccsc} \frac{x}{a} + c$$

Trigonometric functions

$$1. \int \sin x dx = -\cos x + c$$

$$2. \int \cos x dx = \sin x + c$$

$$3. \int \sec^2 x dx = \tan x + c$$

$$4. \int \csc^2 x dx = -\cot x + c$$

5. $\int \tan x \, dx = -\ln|\cos x| + c$ 6. $\int \cot x \, dx = \ln|\sin x| + c$
7. $\int \sec x \, dx = \ln|\sec x + \tan x| + c = \ln\left|\tan\left(\frac{x}{2} + \frac{\pi}{4}\right)\right|$
8. $\int \csc x \, dx = \ln|\csc x - \cot x| + c = \ln\left|\tan\frac{x}{2}\right|$
9. $\int \sec x \tan x \, dx = \sec x + c$
10. $\int \csc x \cot x \, dx = -\csc x + c$

4. Non-basic integration

I) Integration by substitution

In evaluating $\int f(x) \, dx$ when $f(x)$ is not a basic function:

if $f(x) = g'(x)g(x)$ or $f(x) = \frac{g'(x)}{g(x)}$ or

$f(x) = h(g(x))g'(x)$, you let $u = g(x)$.

II) Integration by parts

To integrate a product of functions, try the formula for

integration by parts $\int u \frac{dv}{dx} \, dx = uv - \int v \frac{du}{dx} \, dx$.

An effective strategy is to choose for $\frac{dv}{dx}$ the most complicated factor that can readily be integrated. Then we differentiate the other part, u , to find $\frac{du}{dx}$.

The following table can be used:

u	v'
Logarithmic function	Polynomial function
Polynomial function	Exponential function
Polynomial function	Trigonometric function
Exponential function	Trigonometric function
Trigonometric function	Exponential function
Inverse trigonometric function	Polynomial function

Applying the method of integration by parts, the power of integrand is reduced and the process is continued till we get a power whose integral is known or which can

be easily integrated. This process is called **Reduction formula**.

III) Integration by partial fractions

Remember that:

A **rational function** is a function of the form $f(x) = \frac{P(x)}{Q(x)}$, where $P(x)$ and $Q(x)$ are polynomials.

A **proper rational function** is a rational function in which the degree of $P(x)$ is strictly less than the degree of $Q(x)$.

The problem of integrating rational functions is really the problem of integrating proper rational functions since **improper rational functions** (i.e. those in which the degree of $P(x)$ is greater than or equal to the degree of $Q(x)$) and can always be rewritten as the sum of a polynomial and a proper rational function.

The integrals of proper rational functions are found by **partial fraction expansion** of the integrand into simple fractions.

There are 4 types of simple fractions:

a) Fractions of the type $\frac{A}{x-a}$.

The integrals of such fractions are easily found:

$$\int \frac{A}{x-a} dx = A \ln|x-a| + c$$

b) Fractions of the type $\frac{A}{(x-a)^n}$, where n is a natural number greater than 1.

The integrals of such fractions are easily found:

$$\int \frac{A}{(x-a)^n} dx = A \int (x-a)^{-n} dx = \frac{A}{1-n} (x-a)^{1-n} + c$$

- c) Fractions of the type $\frac{Ax+B}{x^2+px+q}$, where $p^2-4q < 0$

The integrals of such fractions are found by completing the square in the denominator and subsequent substitution which lead to rational integrals of the form

$$\int \frac{du}{u^2+k^2} \text{ or } \int \frac{du}{u^2-k^2} \text{ or } \int \frac{du}{k^2+u^2}.$$

- d) Fractions of the type $\frac{Ax+B}{(x^2+px+q)^n}$,

where $p^2-4q < 0$ and n is a natural number greater than 1.

Integration of this type of fraction will not be considered in this course.

Expansion of proper rational functions in partial fractions is achieved by first factoring the denominator and then writing the type of partial fraction (with unknown coefficients in the numerator) that corresponds to each term in the denominator:

- (i) if the denominator contains $(x-a)$, then the partial fraction expansion will contain $\frac{A}{x-a}$;
- (ii) if the denominator contains $(x-a)^n$, then the partial fraction expansion will contain

$$\frac{A}{(x-a)^n} + \frac{B}{(x-a)^{n-1}} + \frac{C}{(x-a)^{n-2}} + \dots + \frac{Z}{(x-a)}$$

- (iii) if the denominator contains (x^2+px+q) where $p^2-4q < 0$, then the partial fraction expansion will contain $\frac{Ax+B}{x^2+px+q}$.

The unknown coefficients (A, B, etc.) are then found by one of two ways: by inserting concrete values of , or by using the method of undetermined coefficients.

4. Integration of irrational functions

a) Integrals of the form $\int \frac{dx}{\sqrt{ax^2 + bx + c}}$

The integrals of such fractions are found by completing the square in the denominator and subsequent substitution which leads to irrational integrals of the form

$$\int \frac{du}{\sqrt{u^2 + k^2}} \text{ or } \int \frac{du}{\sqrt{u^2 - k^2}} \text{ or } \int \frac{du}{\sqrt{k^2 + u^2}}$$

b) Integrals of the form $\int \frac{px + q}{\sqrt{ax^2 + bx + c}} dx$

The numerator is written as the sum of two parts. One part is the derivative of radicand and the other part is a constant only, i.e.

$$\int \frac{px + q}{\sqrt{ax^2 + bx + c}} dx = k_1 \int \frac{\frac{d}{dx}(ax^2 + bx + c)}{\sqrt{ax^2 + bx + c}} dx + k_2 \int \frac{dx}{\sqrt{ax^2 + bx + c}}$$

c) Integrals of the form $\int \frac{px^2 + qx + r}{\sqrt{ax^2 + bx + c}} dx$

The numerator is written as the sum of three parts. One part is the same as radicand, the second part is derivative of radicand and the last part is a constant only, i.e.

$$\int \frac{px^2 + qx + r}{\sqrt{ax^2 + bx + c}} dx = k_1 \int \frac{ax^2 + bx + c}{\sqrt{ax^2 + bx + c}} dx + k_2 \int \frac{\frac{d}{dx}(ax^2 + bx + c)}{\sqrt{ax^2 + bx + c}} dx + k_3 \int \frac{dx}{\sqrt{ax^2 + bx + c}}$$

5. Integration of trigonometric functions

a) Integrals of the form $\int \frac{dx}{a \sin x + b \cos x + c}$

You can use t-formulae by letting $t = \tan \frac{x}{2}$.

b) Integrals of the form $\int \frac{dx}{a + b \cos^2 x}$ or $\int \frac{dx}{a + b \sin^2 x}$

Here also you can use t-formulae

In integrating the trigonometric functions containing product or power, transforming product or power into sum (or difference) leads to basic integration.

6. Definite integration

Remember that integrals containing an arbitrary constant c in their results are called **indefinite integrals** since their precise value cannot be determined without further information.

a) **Definite integrals** are those in which limits are applied.

If an expression is written as $[F(x)]_a^b$, 'b' is called the upper limit and 'a' the lower limit.

The operation of applying the limits is defined as:

$$[F(x)]_a^b = F(b) - F(a)$$

For example the increase in the value of the integral

$f(x)$ as x increases from 1 to 3 is written as $\int_1^3 f(x) dx$.

The **definite integral**, from $x = a$ to $x = b$, is defined as the area under the curve between those two values.

This is written as $\int_a^b f(x) dx$

b) The mean value of a function $y = f(x)$ over the range $]a, b[$ is the value the functions would have if it were constant over the range but with the same area under the graph. The mean value of $y = f(x)$ over the range $]a, b[$

$$\text{is } \overline{f(x)} = \frac{1}{b-a} \int_a^b f(x) dx .$$

c) The **root mean square** value (R.M.S. value) is the square root of the mean value of the square of y .

The r.m.s. value from $x = a$ to $x = b$ is given by;

$$R.M.S. = \sqrt{\frac{\int_a^b f^2(x) dx}{b-a}}$$

d) Improper integral

The definite integral $\int_a^b f(x)dx$ is called an improper integral if one of two situations occurs:

- ⑦ The limit a or b (or both bounds) are infinites.
- ⑦ The function $f(x)$ has one or more points of discontinuity in the interval $[a, b]$.

Let $f(x)$ be a continuous function on the interval $[a, +\infty[$ or $]-\infty, b]$.

We define the improper integral as

$$\int_a^{+\infty} f(x)dx = \lim_{n \rightarrow +\infty} \int_a^n f(x)dx$$

Or $\int_{-\infty}^b f(x)dx = \lim_{n \rightarrow -\infty} \int_n^b f(x)dx$ respectively.

If these limits exist and are finite, then we say that the improper integrals are convergent, otherwise, the integrals are divergent.

Let $f(x)$ be a continuous function for all real numbers.

By Chasles theorem $\int_{-\infty}^{+\infty} f(x)dx = \int_{-\infty}^c f(x)dx + \int_c^{+\infty} f(x)dx$

If for real number c , both integrals on the right side are convergent, then we say that the integral $\int_{-\infty}^{+\infty} f(x)dx$ is also **convergent**; otherwise it is **divergent**.

7. Applications

Integration has many applications, some of which are listed below:

- a) The area between two functions $f(x)$ and $g(x)$ where $f(x) \leq g(x)$ in $[a, b]$ is given by

$$\int_a^b [g(x) - f(x)]dx = \int_a^b g(x)dx - \int_a^b f(x)dx$$

- b) Volume

The volume of a solid of revolution can be found using one of the following methods:

- ⑦ disc method,

- ☐ washer method, and
- ☐ shell method.

In any of the methods, when finding volume, it is necessary to integrate along the axis of revolution; if the region is revolved about a horizontal line, integrate by x , and if the region is revolved about a vertical line, integrate with respect to y .

(i) **Disc method**

The volume of the solid of revolution bound by the curve $f(x)$ about the x -axis calculated from $x = a$ to $x = b$,

$$\text{is given by } \pi \int_a^b y^2 dx .$$

Volume of the solid generated by revolution of the area bound by the curve $y = f(x)$ about the y -axis is given

$$\text{by } \pi \int_a^b x^2 dy .$$

If the axis of revolution is the line parallel to x -axis (say $y = k$), the volume will be

$$\pi \int_a^b (y - k)^2 dx$$

(ii) **Washer method**

If the region bound by outer radius $y_U = g(x)$ (on top) and inner radius $y_L = f(x)$ and then lines $x = a$, $x = b$ is revolved about x -axis, then the volume of revolution is

$$\text{given by: } V = \pi \int_a^b ([g(x)]^2 - [f(x)]^2) dx$$

(iii) **Shell method**

The volume of the solid generated by revolving the region between the curve x -axis, $y = f(x) \geq 0, L \leq a \leq x \leq b$, about a vertical line $x = L$ is

$$V = 2\pi \int_a^b \left(\begin{matrix} \text{shell} \\ \text{radius} \end{matrix} \right) \left(\begin{matrix} \text{shell} \\ \text{height} \end{matrix} \right) dx$$

HINT for shell method:

Regardless of the position of the axis of revolution (horizontal or vertical), the steps for implementing the shell method are the following:

- ⑦ Draw the region and sketch a line segment across it, parallel to the axis of revolution. Label the segment's height or length (shell height) and distance from the axis of revolution (shell radius).
- ⑦ Find the limits of integration for the thickness variable.
- ⑦ Integrate the product $2\pi \left(\begin{matrix} \text{shell} \\ \text{radius} \end{matrix} \right) \left(\begin{matrix} \text{shell} \\ \text{height} \end{matrix} \right)$ with respect to the thickness variable (*x or y*) to find the volume.
- ⑦ Length of arc of the curve $y = f(x)$ between the points whose abscissas are a and b is

$$s = \int_a^b \sqrt{1 + \left(\frac{dy}{dx} \right)^2} dx$$

- ⑦ The **work** done by a variable force $F(x)$ in the direction of motion along the x -axis over the interval $[a, b]$ is

$$W = \int_a^b F(x) dx.$$

Hook's law says that the force required to hold a stretched or compressed spring x units beyond its equilibrium position pulls back with a force $F(x) = kx$ where k is constant called **spring constant** (or **force constant**).

End of Unit Assessment answers Learner's Book page 285

1. a) $\frac{x^2}{2} + 9x + 125 \ln|x-5| - 64 \ln|x-4| + c$
 - b) $x^3 + x^2 - 5x + 18 \ln|x+3| + c$
 - c) $-\frac{16}{x-2} - \frac{2}{3} \ln|x-2| + \frac{5}{3} \ln|x+1| + c$
 - d) $\ln|x-1| - \frac{1}{2} \ln(x^2+1) + \tan^{-1} x + c$
 - e) $4 \ln|x+2| - \frac{3}{2} \ln(x^2+2x+2) + \tan^{-1}(x+1) + c$
 - f) $\frac{1}{2} \ln(x^2+1) - \ln|x+1| - \frac{3}{x+1} + c$
 - g) $x \sin x + \cos x + c$
 - h) $\frac{5}{4} e^{4x} \left(x - \frac{1}{4} \right) + c$
 - i) $\frac{x^2}{4} (2 \ln x - 1) + c$
 - j) $\frac{\cos 3x}{27} (2 - 9x^2) + \frac{2}{9} x \sin 3x + c$
 - k) $\frac{e^{ax}}{a^2 + b^2} (b \sin bx + a \cos bx) + c$
 - l) $\frac{e^{ax}}{a^2 + b^2} (a \sin bx - a \cos bx) + c$
2. a) $\frac{4}{3}$ b) $\frac{10}{9}$ c) $\ln 2$
 - d) $\frac{\pi}{4}$ e) $\frac{\sqrt{2}}{2}$ f) 1
3. To be proved 4. $20 \frac{5}{6}$
 5. 4 6. $\frac{16}{3}$

7. $\frac{56\pi}{27}$

8. $\frac{32\pi}{3}$

9. $\frac{208\pi}{15}$

10. a) $\frac{\pi}{30}$

b) $\frac{\pi}{6}$

c) $\frac{\pi}{2}$

d) $\frac{5\pi}{6}$

e) $\frac{11\pi}{30}$

f) $\frac{19\pi}{30}$

11. a) $\frac{48\pi}{5}$

b) $\frac{24\pi}{5}$

12. a) $kt = \frac{1}{2a} \ln|a-x| - \ln|a+3x|$

b) $x = \frac{a(e^{2akt} - 1)}{3e^{2akt} - 1}$

c) $x \rightarrow \frac{a}{3}, t \rightarrow \infty$

13. $0.632N_o$

14. 7.26

15. $1.17J$

16. $\frac{9}{5}$

17. a) $30 N/m$


b) $60 J$

c) $1.5 m$

18. a) $926,640$

b) 0.468





Differential Equations

Learner's Book pages 289 - 328

Key unit competence

Use ordinary differential equations of first and second order to model and solve related problems in Physics, Economics, Chemistry, Biology.

Vocabulary or key words concepts

Differential equation (D.E): Equation that involves a function and its derivatives.

First order differential equation: Differential equation containing only first derivatives apart from dependent variable.

Second order differential equation: Differential equation containing second derivatives (and possibly first derivative also).

Particular solution: A solution found at particular values.

Guidance on the problem statement

Let the number of organisms at any time t be $x(t)$. The rate at which new organisms are produced written as $\left(\frac{dx}{dt}\right)$ is proportional to the number of organism present at time, t .

The differential equation is $\frac{dx}{dt} = kx$. k is the constant of proportionality. The problem is to think about what this situation means.

Here, if the number doubles in one day, then the second day there are twice as many available reproduce, so the organisms will double again on the second day and so on. This tells us what solution we are looking for.

List of lessons

No	Lesson title	Number of periods
1	Definition and classification	1
2	Differential equations with separable variables	1
3	Simple homogeneous equations	2
4	Linear equations	2
5	Particular solution	1
6	Second order differential equations: Definition	1
7	Second order differential equations with constant coefficient: two distinct real roots	1
8	Characteristic equation has a double root	1
9	Characteristic equation has complex roots	1
10	Non-homogeneous linear differential equations of the second order with constant coefficients	2
11	Non-homogeneous linear differential equations of the second order with the right hand side $r(x) = Pe^{\alpha x}$	2
12	Non-homogeneous linear differential equations of the second order with the right hand side $r(x) = Pe^{\alpha x} \cos \beta x + Qe^{\alpha x} \sin \beta x$	2
13	Application: Newton's law of cooling	2
14	Application: Electrical circuits	2
Total periods		21

Lesson development

Lesson 5.1. Definition and classification

Learning objectives

Through examples, learners should be able to define and classify given differential equations correctly.

Prerequisites

- ④ Differentiation

Teaching Aids

Exercise book and pen

Activity 5.1 Learner's Book page 290



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. On differentiation; $\frac{dy}{dx} = A$

The given equation becomes $y = x \frac{dy}{dx} + \left(\frac{dy}{dx}\right)^2$

Order of the highest derivative is 1.

2. On differentiation; $\frac{dy}{dx} = -A \sin x + B \cos x$

Again differentiating:

$$\frac{d^2y}{dx^2} = -(A \cos x + B \sin x) = -y$$

$$\text{Or } \frac{d^2y}{dx^2} + y = 0$$

Order of the highest derivative is 2.

3. On differentiation $2y \frac{dy}{dx} = 2Ax + B$

Again differentiating $2y \frac{d^2y}{dx^2} + 2 \left(\frac{dy}{dx} \right)^2 = 2A$

On differentiating again:

$$y \frac{d^3y}{dx^3} + \frac{dy}{dx} \frac{d^2y}{dx^2} + 2 \frac{dy}{dx} \frac{d^2y}{dx^2} = 0$$

$$\text{Or } y \frac{d^3y}{dx^3} + 3 \frac{dy}{dx} \frac{d^2y}{dx^2} = 0$$

Order of the highest derivative is 3.

Synthesis

An equation involving a differential coefficient i.e.

$\frac{dy}{dx}, \frac{d^2y}{dx^2}, \frac{dr}{dt}$ and so on is called a “**differential equation**”.

Order of the differential equation is the highest derivative of function that appears in a differential equation and is said to be the **order of differential equation**.

Given a function with arbitrary constants, you form differential equation by eliminating its arbitrary constants using differentiation.

Exercise 5.1 Learner's Book page 292

1. a) $y^2 \left(\frac{dy}{dx} \right)^2 + 2xy \frac{dy}{dx} - y^2 = 0$ b) $xy \frac{d^2y}{dx^2} + x \left(\frac{dy}{dx} \right)^2 - y \frac{dy}{dx} = 0$

c) $\frac{d^2y}{dx^2} - 4 \frac{dy}{dx} + 3y = 0$ d) $\frac{d^2y}{dx^2} - 2 \frac{dy}{dx} + 2y = 0$

e) $\frac{dy}{dx} = -\tan(x+3)$

2. a) This DE has **order 2** (the highest derivative appearing is the **second** derivative) and **degree 1** (the **power** of the highest derivative is 1).
- b) This DE has **order 1** (the highest derivative appearing is the **first** derivative) and **degree 4** (the **power** of the highest derivative is 4).

- c) This DE has **order 2** (the highest derivative appearing is the **second** derivative) and **degree 3** (the **power** of the highest derivative is 3).
- d) order 2; degree 1
- e) order 2; degree 1
- f) order 3; degree 1
- g) order 2; degree 1
- h) order 2; degree 3

Lesson 5.2. Differential equations with separable variables

Learning objectives

Through examples, learners should be able to identify and solve differential equations with separable variables accurately.

Prerequisites

- ☑ Integration

Teaching Aids

Exercise book and pen

Activity 5.2 Learner's Book page 293



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ☑ Critical thinking
- ☑ Communication
- ☑ Self confidence
- ☑ Cooperation, interpersonal management and life skills
- ☑ Peace and values education
- ☑ Inclusive education

Answers

$$1. \frac{dy}{dx} = \frac{x}{y} \Rightarrow ydy = xdx \Rightarrow \int ydy = \int xdx \Rightarrow \frac{y^2}{2} = \frac{x^2}{2} + k$$

$$\text{Or } y^2 = x^2 + 2k \Rightarrow y^2 = x^2 + c, \quad c \in \mathbb{R}$$

$$2. \frac{dy}{dx} = x^2 y^3 \Rightarrow \frac{dy}{y^3} = x^2 dx \Rightarrow \int \frac{dy}{y^3} = \int x^2 dx$$

$$\Rightarrow \int y^{-3} dy = \int x^2 dx \Rightarrow \frac{y^{-2}}{-2} = \frac{x^3}{3} + k \Rightarrow -\frac{1}{2y^2} = \frac{x^3}{3} + k, \quad k \in \mathbb{R}$$

Synthesis

To solve the integral $\frac{dy}{dx} = g(x)h(y)$, we write it in the separated form $\frac{dy}{h(y)} = g(x)dx$ and integrate.

Exercise 5.2 Learner's Book page 294

1. $y^2 = cx, c \in \mathbb{R}$
2. $x^3 - y^3 = c, c \in \mathbb{R}$
3. $\arctan y = x + c, c \in \mathbb{R}$
4. $\tan^{-1} y = x - \ln|1+x| + c, c \in \mathbb{R}$

Lesson 5.3. Simple homogeneous equations

Learning objectives

Through examples, learners should be able to identify and solve simple homogeneous equations accurately.

Prerequisites

- ✔ Integration

Teaching Aids

Exercise book and pen



Activity 5.3 Learner's Book page 294

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad f(tx, ty) = (tx)^2 + (tx)(ty)$$

$$f(tx, ty) = t^2x^2 + t^2xy$$

$$f(tx, ty) = t^2(x^2 + xy)$$

$$f(tx, ty) = t^2 f(x, y)$$

The value of n is 2.

$$2. \quad z = \frac{y}{x} \Rightarrow y = zx$$

$$\frac{dy}{dx} = z + x \frac{dz}{dx}$$

$$\frac{dy}{dx} = f(x, y) \text{ becomes } z + x \frac{dz}{dx} = f(x, y)$$

$$\text{But } f(x, y) = f(1, z)$$

$$3. \quad f(tx, ty) = f(x, y)$$

$$\text{But } t = \frac{1}{x}, \text{ then } f\left(1, \frac{y}{x}\right) = f(x, y) \Rightarrow f(1, z) = f(x, y)$$

$$\text{Then, } z + x \frac{dz}{dx} = f(1, z)$$

Separating variables, we have

$$x \frac{dz}{dx} = f(1, z) - z \Rightarrow x \frac{dz}{f(1, z) - z} = dx \Rightarrow \frac{dz}{f(1, z) - z} = \frac{dx}{x}$$

Synthesis

A function $f(x, y)$ is called homogeneous of degree n if $f(tx, ty) = t^n f(x, y)$ for all suitably restricted x, y and t .

The differential equation $M(x, y)dx + N(x, y)dy = 0$ is said to be homogeneous if M and N are homogeneous functions of the same degree.

This equation can be written in the form $\frac{dy}{dx} = f(x, y)$.

Where $f(x, y) = \frac{-M(x, y)}{N(x, y)}$ is clearly homogeneous of degree 0. We solve this equation by letting $z = \frac{y}{x}$.

Exercise 5.3 Learner's Book page 297

1. $x^2 - y^2 = c, c \in \mathbb{R}$
2. $x^2 + y^2 = cx, c \in \mathbb{R}$
3. $y^2 = x^2(cx^2 - 4), c \in \mathbb{R}$
4. $(y - x)^2 + 2(y - x) = 2x + c, \in \mathbb{R}$

Lesson 5.4. Linear differential equations

Learning objectives

Through examples, learners should be able to identify and solve linear differential equations accurately.

Prerequisites

- ✔ Differentiation
- ✔ Integration

Teaching Aids

Exercise book and pen



Activity 5.4 Learner's Book page 298

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $y = uv$

$$\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$$

2. Now, $\frac{dy}{dx} + py = q$ becomes $u \frac{dv}{dx} + v \frac{du}{dx} + p(uv) = q$

3. $u \frac{dv}{dx} + v \frac{du}{dx} + p(uv) = q \Rightarrow u \frac{dv}{dx} + p(uv) + v \frac{du}{dx} = q$

$$\Rightarrow u \left(\frac{dv}{dx} + pv \right) + v \frac{du}{dx} = q$$

If $\frac{dv}{dx} + pv = 0$, $dv + pvdx = 0 \Rightarrow dv = -pvdx$

Separating variables, we have $\frac{dv}{v} = -pdx$

Integrating both sides, we have $\int \frac{dv}{v} = \int -pdx$

$$\ln|v| = -\int pdx + c \Rightarrow \ln|v| = \ln e^{-\int pdx} + \ln k$$

$$\Rightarrow \ln|v| = \ln ke^{-\int pdx} \Rightarrow |v| = ke^{-\int pdx}$$

Take $v = e^{-\int pdx}$

4. Now, the equation $u \left(\frac{dv}{dx} + pv \right) + v \frac{du}{dx} = q$ becomes

$$e^{-\int pdx} \frac{du}{dx} = q \text{ since } \frac{dv}{dx} + pv \text{ is assumed to zero.}$$

$$e^{-\int pdx} \frac{du}{dx} = q \Rightarrow \frac{du}{dx} = \frac{q}{e^{-\int pdx}}$$

$$\Rightarrow \frac{du}{dx} = q e^{\int pdx} \Rightarrow du = q e^{\int pdx} dx$$

Integrating both sides gives

$$\Rightarrow \int du = \int q e^{\int pdx} dx \Rightarrow u = \int q e^{\int pdx} dx$$

Synthesis

The general solution to the equation $\frac{dy}{dx} + py = q$ where p and q are functions in x or constants, is $y = uv$ where $u = \int q e^{\int p dx} dx$ and $v = e^{-\int p dx}$.

Short cut method:

The solution of $\frac{dy}{dx} + py = q$ is simply given by formula $y = e^{-\int p dx} \int q e^{\int p dx} dx$.

Exercise 5.4 Learner's Book page 299

1. $y = \frac{x}{2} + \frac{c}{x}$
2. $y = x^2 - 2 + ce^{-\frac{x^2}{2}}$
3. $y = (x+1)e^x + c(x+1)$
4. $y = \cos x + c \cos^2 x$

Lesson 5.5. Particular solution**Learning objectives**

Given a differential equation and initial condition, learners should be able to find a particular solution for that differential equation accurately.

Prerequisites

- ✔ Integration

Teaching Aids

Exercise book and pen

**Activity 5.5 Learner's Book page 300**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \frac{dy}{dx} = x + 4$$

$$\Rightarrow dy = (x + 4)dx \Rightarrow \int dy = \int (x + 4)dx$$

$$\Rightarrow y = \frac{x^2}{2} + 4x + c$$

$$2. \text{ If } x = 2 \text{ then } y = 4$$

$$\Rightarrow 4 = \frac{2^2}{2} + 4 \times 2 + c \Rightarrow 4 = 10 + c \Rightarrow c = -6$$

$$\text{New solution is } y = \frac{x^2}{2} + 4x - 6$$

Synthesis

If we want to determine a function, $y(x)$, such that the given equation is satisfied for $y(x_0) = y_0$ or $y|_{x=x_0} = y_0$, this equation is referred to as an **initial value problem** for the obvious reason that out of the totality of all solution of the differential equation, we are looking for the one solution which initially (at, x_0) has the value y_0 .

Exercise 5.5 Learner's Book page 301

$$1. y = \cos x - 2 \cos^2 x$$

$$2. y = \tan \left(\tan^{-1} x + \frac{\pi}{4} \right)$$

$$3. y = \frac{1}{\sqrt{1-x^2}}$$

$$4. y^2 = x^2 - 2 \ln x + 3$$

$$5. e^y = \frac{1}{2}e^{2x} + \frac{1}{2}$$

Lesson 5.6. Second order differential equations: Definition

Learning objectives

Through examples, learners should be able to define a second order differential equation accurately.

Prerequisites

- ✔ Differentiation

Teaching Aids

Exercise book and pen



Activity 5.6 Learner's Book page 301

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

Answer may vary; here are some.

1. Second order differential equation with degree greater than 1 is of the form

$$\left(\frac{d^2y}{dx^2}\right)^n + p(x)\left(\frac{dy}{dx}\right)^k + q(x)y = r(x),$$

where $p(x)$, $q(x)$ and $r(x)$ are functions of x alone (or perhaps constants) and $n, k \in \mathbb{Z}$ with $n \geq 1$.

2. Second order differential equation with degree 1 is of the form

$$\left(\frac{d^2y}{dx^2}\right) + p(x)\left(\frac{dy}{dx}\right)^k + q(x)y = r(x),$$

where $p(x)$, $r(x)$ and $r(x)$ are functions of x alone (or perhaps constants) and $k \in \mathbb{Z}$.

Synthesis

The general second order linear differential equation is

of the form $\frac{d^2y}{dx^2} + p(x)\frac{dy}{dx} + q(x)y = r(x)$ or more

simply, $y'' + p(x)y' + q(x)y = r(x)$;

where $p(x)$, $q(x)$ and $r(x)$ are functions of x alone (or perhaps constants).

Lesson 5.7. Second order differential equations with constant coefficient: two distinct real roots

Learning objectives

Given a second order differential equations with constant coefficient where characteristic equation has two distinct real roots, learners should be able to find its general solution perfectly.

Prerequisites

- ✔ Solving quadratic equation.

Teaching Aids

Exercise book and pen

Activity 5.7 Learner's Book page 302



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad y' + ky = 0 \Rightarrow \frac{dy}{dx} = -ky \Rightarrow \frac{dy}{y} = -kdx$$

$$\Rightarrow \int \frac{dy}{y} = \int -kdx \Rightarrow \ln|y| = -kx$$

$$\Rightarrow \ln|y| = \ln e^{-kx} \Rightarrow y = e^{-kx}$$

$$2. \quad y'' + py' + qy = 0$$

$$\text{But } y = e^{-kx} \Rightarrow y' = -ke^{-kx} \text{ and } y'' = (-k)^2 e^{-kx}$$

$$y'' + py' + qy = 0 \text{ becomes}$$

$$(-k)^2 e^{-kx} + p(-ke^{-kx}) + qe^{-kx} = 0$$

$$\Rightarrow [(-k)^2 - kp + q]e^{-kx} = 0$$

This relation is true if $(-k)^2 - kp + q = 0$ since e^{-kx} cannot be zero.

Then $(-k)^2 - kp + q = 0$. Putting $m = -k$, we have

$$m^2 + mp + q = 0$$

Thus, the solution of $y' + ky = 0$ is also a solution of

$y'' + py' + qy = 0$ if m satisfy the auxiliary equation

$$m^2 + mp + q = 0 \text{ for } m = -k.$$

Therefore, the solution of the form e^{mx} is the solution of $y'' + py' + qy = 0$.

Synthesis

In solving homogeneous linear equation of second order

$y'' + py' + qy = 0$, we first determine its characteristic

equation which is $m^2 + mp + q = 0$.

If m_1 and m_2 are solutions of the characteristic

equation, then the general solution of $y'' + py' + qy = 0$ is

$$y = c_1 e^{m_1 x} + c_2 e^{m_2 x} \text{ where } m_1, m_2 = \frac{-p \pm \sqrt{p^2 - 4q}}{2}.$$

Exercise 5.6 Learner's Book page 304

1. $y = c_1 e^{3x} + c_2 e^{5x}$

2. $y = c_1 e^{-2x} + c_2 e^x$

3. $y = c_1 e^{5x} + c_2 e^{-6x}$

4. $y = c_1 e^{-3x} + c_2 e^{-7x}$

Lesson 5.8. Characteristic equation with a double root**Learning objectives**

Given a second order differential equations with constant coefficient where characteristic equation has one double root, learners should be able to find its general solution correctly.

Prerequisites

- ④ Solving quadratic equation.

Teaching Aids

Exercise book and pen

Activity 5.8 Learner's Book page 304

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. Characteristic equation:

$$m^2 + 2m + 1 = 0$$

$$\Delta = 4 - 4 = 0$$

$$\text{Thus, } m_1 = m_2 = -\frac{2}{2} = -1$$

One of solutions of $\frac{d^2 y}{dx^2} + 2\frac{dy}{dx} + y = 0$ is $y_1 = e^{-x}$

$$2. \quad y_2 = xy_1 \Rightarrow y_2 = xe^{-x}$$

Since $y_2 = xe^{-x}$, then

$$\frac{dy_2}{dx} = \frac{d(xe^{-x})}{dx} = e^{-x} - xe^{-x}$$

$$\frac{d^2y_2}{dx^2} = \frac{d(e^{-x} - xe^{-x})}{dx} = -e^{-x} - e^{-x} + xe^{-x} = -2e^{-x} + xe^{-x}$$

Substituting y by $y_2 = xe^{-x}$ in $\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0$ yield

$$\begin{aligned} & (-2e^{-x} + xe^{-x}) + 2(e^{-x} - xe^{-x}) + xe^{-x} \\ &= \cancel{-2e^{-x}} + \cancel{xe^{-x}} + \cancel{2e^{-x}} - \cancel{2xe^{-x}} + \cancel{xe^{-x}} = 0 \end{aligned}$$

We note that $y_2 = xe^{-x}$ is also a solution of

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0$$

The ratio $\frac{y_1}{y_2} = \frac{e^{-x}}{xe^{-x}} = \frac{1}{x}$ is not constant, thus,

$y_1 = e^{-x}$ and $y_2 = xe^{-x}$ are linearly independent and $y = c_1e^{-x} + c_2xe^{-x}$ is the general solution of

$$\frac{d^2y}{dx^2} + 2\frac{dy}{dx} + y = 0$$

(c_1 and c_2 being arbitrary constants).

Synthesis

In solving homogeneous linear equation of second order $y'' + py' + qy = 0$, if the characteristic equation $m^2 + mp + q = 0$ has a double root equal to m , the general solution of equation $y'' + py' + qy = 0$ will be $y = c_1e^{mx} + c_2xe^{mx}$.

Exercise 5.7 Learner's Book page 305

$$1. \quad y = c_1e^{3x} + c_2xe^{3x}$$

$$2. \quad y = c_1e^{-3x} + c_2xe^{-3x}$$

$$3. \quad y = (c_1 + c_2x)e^{4x}$$

$$4. \quad y = (c_1 + xc_2)e^{\frac{1}{6}x}$$

Lesson 5.9. Characteristic equation with complex roots

Learning objectives

Given a second order differential equations with constant coefficient where the characteristic equation has complex roots, learners should be able to find its general solution correctly.

Prerequisites

- ✔ Solving quadratic equation in complex numbers.

Teaching Aids

Exercise book and pen

Activity 5.9 Learner's Book page 305



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Characteristic equation is $m^2 - 4m + 25 = 0$

$$\Delta = 16 - 100 = -64 \quad \sqrt{\Delta} = \pm 8i$$

$$m_1 = \frac{4+8i}{2} = 2+4i \quad m_2 = \frac{4-8i}{2} = 2-4i$$

a) The basis of $\frac{d^2y}{dx^2} - 4\frac{dy}{dx} + 25y = 0$ are

$$y_1 = c_1 e^{(2+4i)x} \quad \text{and} \quad y_2 = c_2 e^{(2-4i)x}$$

b) Its general solution is $y = c_1 e^{(2+4i)x} + c_2 e^{(2-4i)x}$

2. $y = c_1 e^{(2+4i)x} + c_2 e^{(2-4i)x}$

$$\Leftrightarrow y = c_1 e^{2x+4ix} + c_2 e^{2x-4ix} \Leftrightarrow y = e^{2x} (c_1 e^{4ix} + c_2 e^{-4ix})$$

$$\Leftrightarrow y = e^{2x} [c_1 (\cos 4x + i \sin 4x) + c_2 (\cos 4x - i \sin 4x)]$$

$$\Leftrightarrow y = e^{2x} [(c_1 + c_2) \cos 4x + (c_1 - c_2) i \sin 4x]$$

$$\Leftrightarrow y = e^{2x} (c_1 + c_2) \cos 4x + e^{2x} (c_1 - c_2) i \sin 4x$$

$$\Leftrightarrow y = e^{2x} [(c_1 + c_2) \cos 4x + (c_1 - c_2) i \sin 4x]$$

3. Real basis are $y_1 = A e^{2x} \cos 4x$ and $y_2 = B e^{2x} \sin 4x$

4. General solution is $y = e^{2x} (A \cos 4x + B \sin 4x)$

Synthesis

If the characteristic equation has complex roots, $\alpha \pm i\beta$ then, the general solution is $y = e^{\alpha x} (A \cos \beta x + B \sin \beta x)$, where α and β are respectively, real and imaginary part of root of characteristic equation.

Exercise 5.8 Learner's Book page 307

$$1. \quad y = e^{-2x} (c_1 \cos 3t + c_2 \sin 3t) \quad 2. \quad y = e^{-2x} (c_1 \cos x + c_2 \sin x)$$

$$3. \quad y = e^x (c_1 \cos x + c_2 \sin x) \quad 4. \quad y = -e^{\pi-2x} \sin 3x$$

$$5. \quad y = \frac{2}{9} e^{2x} - \frac{2}{9} e^{-\frac{5x}{2}}$$

Lesson 5.10. Non-homogeneous linear differential equations of the second order with constant coefficients

Learning objectives

Through examples, learners should be able to identify a non-homogeneous linear differential equation of the second order with constant coefficients and solve it where possible correctly.

Prerequisites

- ① Solving homogeneous differential equation of second order.

Teaching Aids

Exercise book, calculator and pen

Activity 5.10 Learner's Book page 307

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. $\frac{dy}{dx} - \frac{y}{x+1} = e^x(x+1)$ is a linear differential equation of 1st order.

$$\text{Let } y = u \cdot v, \text{ then } \frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}.$$

Substituting in the given equation, we get

$$u \frac{dv}{dx} + v \frac{du}{dx} - \frac{u \cdot v}{x+1} = e^x(x+1)$$

$$\text{Or } u \frac{dv}{dx} - \frac{u \cdot v}{x+1} + v \frac{du}{dx} = e^x(x+1)$$

$$\Leftrightarrow u \left(\frac{dv}{dx} - \frac{v}{x+1} \right) + v \frac{du}{dx} = e^x(x+1)$$

Taking $\frac{dv}{dx} - \frac{v}{x+1} = 0$, you get $\ln v = \ln(x+1)$ or

$$v = x+1.$$

$$\text{As } \frac{dv}{dx} - \frac{v}{x+1} = 0, u \left(\frac{dv}{dx} - \frac{v}{x+1} \right) + v \frac{du}{dx} = e^x(x+1)$$

$$(x+1) \frac{du}{dx} = e^x(x+1) \Leftrightarrow \frac{du}{dx} = e^x \text{ or } u = e^x + c.$$

The solution of the given equation is then,

$$y = (x+1)(e^x + c) \text{ or } y = (x+1)e^x + c(x+1)$$

$$2. \frac{d^2y}{dx^2} - 4\frac{dy}{dx} = 5y \Leftrightarrow \frac{d^2y}{dx^2} - 4\frac{dy}{dx} - 5y = 0$$

is homogeneous linear equation of second order.

Characteristic equation

$$m^2 - 4m - 5 = 0 \quad \Delta = 16 + 20 = 36$$

$$m_1 = \frac{4-6}{2} = -1, m_2 = \frac{4+6}{2} = 5$$

General solution is $y = c_1e^{-x} + c_2e^{5x}$

$$3. \frac{d^2y}{dx^2} - 4\frac{dy}{dx} - 5y = x \text{ is non-homogeneous}$$

linear equation of second order.

At this level, it is impossible for most learners to solve this type of equation.

[General solution is given by $y = \bar{y} + y^*$.

From 1) Complementary solution is $y = c_1e^{-x} + c_2e^{5x}$.

Let $y^* = Ax + B$ be particular solution of the given equation.

Then $y^{*'} = A$ and $y^{*''} = 0$.

Putting $y^* = Ax + B$ and its derivatives in

$$\frac{d^2y}{dx^2} - 4\frac{dy}{dx} - 5y = x, \text{ gives}$$

$$0 - 4A - 5(Ax + B) = x \Leftrightarrow -5Ax - 4A - 5B = x$$

Identifying the coefficients, we get

$$-5A = 1 \text{ and } -4A - 5B = 0$$

$$\text{Or } A = -\frac{1}{5} \text{ and } B = \frac{4}{25}.$$

$$\text{Thus, particular solution is } y^* = -\frac{1}{5}x + \frac{4}{25}$$

$$\text{The general solution is } y = c_1e^{-x} + c_2e^{5x} - \frac{1}{5}x + \frac{4}{25}$$

Synthesis

The general solution of the second order non-homogeneous linear equation $y'' + py' + qy = r(x)$ can be expressed in

the form $y = \bar{y} + y^*$ where y^* is any specific function that satisfies the non-homogeneous equation, and $\bar{y} = c_1 y_1 + c_2 y_2$ is a general solution of the corresponding homogeneous equation $y'' + py' + qy = 0$.

Lesson 5.11. Differential equations of the second order with the right hand side

$$r(x) = Pe^{\alpha x}$$

Learning objectives

Given a differential equation of second order where the right hand side is of the form $r(x) = Pe^{\alpha x}$, learners should be able to find its general solution correctly.

Prerequisites

- ④ Solving homogeneous differential equation of second order..

Teaching Aids

Exercise book, calculator and pen

Activity 5.11 Learner's Book page 308



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. Characteristic equation: $m^2 - 2m + 1 = 0$

$$\Delta = 4 - 4 = 0$$

$$m_1 = m_2 = \frac{2-0}{2} = 1 \quad \bar{y} = c_1 e^x + c_2 x e^x$$

2. The right hand side can be written as $e^x = 1e^{1x}$.

$$P=1 \text{ and } \alpha=1$$

$\alpha=1$, is double root of characteristic equation, so

$$k=2$$

$y^* = Ax^2e^x$, $Q(x) = A$ as $P=1$ on right hand side, $Q(x)$ has degree zero.

3. $y^{*'} = 2Axe^x + Ax^2e^x$

$$y^{*''} = 2Ae^x + 2Axe^x + 2Axe^x + Ax^2e^x$$

$$\Rightarrow 2Ae^x + 2Axe^x + 2Axe^x + Ax^2e^x - 4Axe^x - 2Ax^2e^x + Ax^2e^x = e^x$$

$$\Rightarrow 2A + 2Ax + 2Ax + Ax^2 - 4Ax - 2Ax^2 - Ax^2 = 1$$

$$\Rightarrow -2Ax^2 + 2Ax^2 + x(2A + 2A - 4A) - 2A = 1$$

$$2A = 1 \Rightarrow A = \frac{1}{2}$$

$$\text{Thus, } y^* = \frac{1}{2}x^2e^x$$

Synthesis

If the right hand side of the equation $y'' + py' + qy = r(x)$ is $r(x) = Pe^{\alpha x}$ where P is a polynomial, we take the particular solution to be

$$y^* = x^k Q_n(x)e^{\alpha x}, Q_n = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_n$$

Here, k is the number of roots of the associated homogeneous equation equal to α .

α ; coefficient of x in $e^{\alpha x}$ on the right hand side,

n ; degree of $Q(x)$, the same as degree of $P(x)$ on right hand side.

3 cases arise

⊙ If α is not a root of characteristic equation $k=0$

⊙ If α is a simple root of characteristic equation $k=1$

⊙ If α is a double root of characteristic equation $k=2$

Note that the simple root or double root in the last 2 cases must be real numbers.

Exercise 5.9 Learner's Book page 310

$$1. \quad y = c_1 e^{-3x} + c_2 x e^{-3x} + \frac{5e^{3x}}{36} \quad 2. \quad c_1 e^x + c_2 e^{2x} + \frac{e^{3x}}{2}$$

$$3. \quad y = c_1 e^{-x} + c_2 e^{-2x} + \frac{1}{4} e^{2x} \quad 4. \quad y = c_1 e^x + c_2 e^{2x} + \frac{1}{2} e^{3x}$$

$$5. \quad y = (c_1 + c_2 x) e^{3x} + \frac{x^2}{2} e^{3x}$$

Lesson 5.12. Differential equations of the second order with the right hand side

$$r(x) = P e^{\alpha x} \cos \beta x + Q e^{\alpha x} \sin \beta x$$

Learning objectives

Given differential equations of second order where the right hand side is of the form $r(x) = P e^{\alpha x} \cos \beta x + Q e^{\alpha x} \sin \beta x$, learners should be able to find its solution correctly.

Prerequisites

- ④ Solving homogeneous differential equations of the second order.

Teaching Aids

Exercise book, calculator and pen

Activity 5.12 Learner's Book page 311

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. Characteristic equation: $m^2 + 4 = 0$

$$m_1 = 2i, m_2 = -2i$$

$$\bar{y} = c_1 \cos 2x + c_2 \sin 2x$$

2. The right hand side of the given equation is written as $\cos 2x = 1e^{0x} \cos 2x + 0e^{0x} \sin 2x$

$$P = 1, Q = 0, \alpha = 0, \beta = 2,$$

$\alpha + \beta i = 0 + 2i = 2i$ is a root of characteristic equation, so $r = 1$

Highest degree of P and Q is zero since $P = 1, Q = 0$

Then, $u = A, v = B$ and $y^* = x(A \cos 2x + B \sin 2x)$

3. $y^{*'} = A \cos 2x + B \sin 2x + x(-2A \sin 2x + 2B \cos 2x)$

$$= A \cos 2x + B \sin 2x + 2x(-A \sin 2x + B \cos 2x)$$

$$y^{*''} = -2A \sin 2x + 2B \cos 2x - 2A \sin 2x + 2B \cos 2x$$

$$+ 2x(-2A \cos 2x - 2B \sin 2x)$$

$$= 4(-A \sin 2x + B \cos 2x) + 4x(-A \cos 2x - B \sin 2x)$$

$$\Rightarrow 4(-A \sin 2x + B \cos 2x) + 4x(-A \cos 2x - B \sin 2x)$$

$$+ 4x(A \cos 2x + B \sin 2x) = \cos 2x$$

$$\Rightarrow -4A \sin 2x + 4B \cos 2x - 4x(A \cos 2x + B \sin 2x)$$

$$+ 4x(A \cos 2x + B \sin 2x) = \cos 2x$$

$$\Rightarrow -4A \sin 2x + 4B \cos 2x = \cos 2x$$

$$\begin{cases} -4A = 0 \\ 4B = 1 \end{cases} \Rightarrow \begin{cases} A = 0 \\ B = \frac{1}{4} \end{cases}$$

$$y^* = x \left(0 \cos 2x + \frac{1}{4} \sin 2x \right) = \frac{1}{4} x \sin 2x$$

Synthesis

If the right hand side of the equation $y'' + py' + qy = r(x)$ is $r(x) = Pe^{\alpha x} \cos \beta x + Qe^{\alpha x} \sin \beta x$ where P and Q are polynomials, two cases arise:

- ❖ If $\alpha + i\beta$ is not a root of characteristic equation, the particular solution is

$$y^* = Ue^{\alpha x} \cos \beta x + Ve^{\alpha x} \sin \beta x$$

- ❖ If $\alpha + i\beta$ is a root of characteristic equation, the particular solution becomes,

$$y^* = x[Ue^{\alpha x} \cos \beta x + Ve^{\alpha x} \sin \beta x]$$

In all cases, U and V are polynomial for which their degree is equal to the highest degree of P and Q .

Exercise 5.10 Learner's Book page 316

1. $y = c_1 e^x + c_2 x e^x + \frac{1}{2}(x \cos x + \cos x - \sin x)$
2. $y = c_1 e^x + c_2 x e^x - e^x(x \sin x + 2 \cos x)$
3. $y = c_1 e^x + c_2 e^{-x} - \frac{1}{10} \left(\frac{3}{5} \cos 3x + x \sin 3x + 5 \cos x \right)$
4. $y = c_1 \cos x + c_2 \sin x - x \cos x + \sin x \ln |\sin x|$
5. $y = c_1 \cos x + c_2 \sin x - \cos x \ln |\sec x + \tan x|$

Lesson 5.13. Applications: Newton's law of cooling**Learning objectives**

By reading textbooks or accessing internet, learners should be able to use differential equations to solve problems involving Newton's law of cooling accurately.

Prerequisites

- ❖ Solving differential equations

Teaching Aids

Exercise book, calculator, library or internet if available and pen



Activity 5.13 Learner's Book page 317

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education
- ④ Research

Answers

To formulate a model, we need to know something about how a liquid cools.

Experimental evidence shows that the rate at which temperature changes is proportional to the difference in temperature between the liquid and the surrounding (ambient) air. If T is the temperature of the liquid at time t then in this case;

$$\frac{dT}{dt} = -k(T - 20) \text{ where } k \text{ is the constant of proportionality}$$

and the negative sign shows that the temperature is reducing.

When coffee is made, its temperature is $90^{\circ}C$. So

$$T = 90^{\circ}C \text{ when } t = 0.$$

In formulating this model, we assume that;

- ④ The temperature throughout the coffee is uniform.
- ④ The temperature of surrounding air is constant.
- ④ The rate of cooling of a body is proportional to the temperature of the body above that of the surrounding air.

Synthesis

Newton's law of cooling states that the rate at which an object cools is proportional to the difference between the temperature at the surface of the body, and the ambient air temperature.

Thus, if T is the surface temperature at time t and T_a is the ambient temperature, then $\frac{dT}{dt} = -\lambda(T - T_a)$ where $\lambda > 0$ is some experimentally determined constant of proportionality, and T_0 is the initial temperature.

Lesson 5.14. Applications: Electrical circuits**Learning objectives**

By reading textbooks or accessing internet, learners should be able to use differential equations in solving electrical circuit problems accurately.

Prerequisites

- ✔ Solving differential equations.
- ✔ Alternating current.

Teaching Aids

Exercise book, calculator, library or internet if available and pen

Activity 5.14 Learner's Book page 318

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Research
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

a) Rearranging $E - L\left(\frac{di}{dt}\right) = Ri$, gives $\frac{di}{dt} = \frac{E - Ri}{L}$

and separating the variables, we get

$$\frac{di}{E - Ri} = \frac{dt}{L}$$

Integrating both sides gives:

$$\int \frac{di}{E - Ri} = \int \frac{dt}{L} \Rightarrow -\frac{1}{R} \ln(E - Ri) = \frac{t}{L} + c$$

When $t = 0$, $i = 0$, thus $-\frac{1}{R} \ln E = c$

Thus, the particular solution is:

$$-\frac{1}{R} \ln(E - Ri) = \frac{t}{L} - \frac{1}{R} \ln E$$

Rearranging gives:

$$\Leftrightarrow -\frac{1}{R} \ln(E - Ri) + \frac{1}{R} \ln E = \frac{t}{L}$$

$$\Leftrightarrow -\frac{1}{R} \ln \frac{E - Ri}{E} = \frac{t}{L}$$

$$\Leftrightarrow \frac{1}{R} \ln \frac{E}{E - Ri} = \frac{t}{L} \Leftrightarrow \ln \frac{E}{E - Ri} = \frac{Rt}{L}$$

from which

$$\frac{E}{E - Ri} = e^{\frac{Rt}{L}} \Leftrightarrow \frac{E - Ri}{E} = e^{-\frac{Rt}{L}}$$

$$\Leftrightarrow E - Ri = Ee^{-\frac{Rt}{L}} \Rightarrow Ri = E - Ee^{-\frac{Rt}{L}}$$

Therefore, $i = \frac{E}{R} \left(1 - e^{-\frac{Rt}{L}}\right)$

b) $i = \frac{E}{R} (1 - e^{-3}) \approx 0.95 \frac{E}{R}$

c) $i = \frac{E}{R} (1 - e^{-2}) \approx 0.86 \frac{E}{R}$ i.e. 86%

Synthesis

In the R–L series circuit shown in figure 5.1, the supply p.d., E , is given by

$$E = V_R + V_L, \quad V_R = iR \quad \text{and} \quad V_L = L \frac{di}{dt}$$

Hence $E = iR + L \frac{di}{dt}$. From which $E - L \frac{di}{dt} = iR$




Figure 5.1: R-L series

The corresponding solution is $i = \frac{E}{R} \left(1 - e^{-\frac{Rt}{L}} \right)$ which represents the law of growth of current in an inductive circuit as shown in figure 5.2




Figure 5.2: Law of growth of current

The growth of the current in the RL circuit is the current's steady-state value. The number $t = \frac{L}{R}$ is the time constant of the circuit. The current gets to within 5% of its steady-state value in 3 times constant.

Exercise 5.11 Learner's Book page 320

1. P indicates number of rabbits, t time in months.

Differential equation:

$$\frac{dP}{dt} = 0.7P, P = 10 \text{ when } t = 0$$

2. To be proved

3. a) $\frac{dT}{dt} = k(1 - 32.2)$

$$T(0) = 34.8, T(1) = 34.1$$

T is temperature in $^{\circ}C$, t is time hours after 2:30 a.m. and k is constant.

Assume that the rate of temperature change is proportional to the difference between body temperature and room temperature. Assume room temperature is constant.

b) $T = 32.2 + 2.6e^{-0.31t}$

c) 0:33 a.m. ($t = -117$ minutes)

4. (a) $Q = Q_0 e^{-\frac{t}{CR}}$ (b) 9.30C, 5.81C

5. 273.3N, 2.31rads

Summary of the unit

1. Definition and classification

An equation involving one or more differential coefficients i.e.

$$\frac{dy}{dx}, \frac{d^2y}{dx^2}, \frac{dr}{dt} \text{ is called a **differential equation** .}$$

Order of the highest derivative of function that appears in a differential equation is said to be the **order of differential equation**.

The general ordinary differential equation of the n^{th} order is

$$F\left(x, y, \frac{dy}{dx}, \frac{d^2y}{dx^2}, \dots, \frac{d^n y}{dx^n}\right) = 0, \text{ OR}$$

$$F(x, y, y', y'', \dots, y^{(n)}) = 0$$

2. First order differential equations

The general differential equation of the 1st order is

$$F\left(x, y, \frac{dy}{dx}\right) = 0 \text{ or } \frac{dy}{dx} = f(x, y)$$

The simplest is that in which the variables are separable:

$$\frac{dy}{dx} = g(x)h(y).$$

A homogeneous equation of degree 0 can be expressed as a

function of $z = \frac{y}{x}$.

The general solution to the equation $\frac{dy}{dx} + py = q$ where p and q are functions in x or constants, is $y = uv$ where

$$u = \int q e^{\int p dx} dx \text{ and } v = e^{-\int p dx}.$$

3. Second order differential equations

The general second order linear differential equation is of the form

$$\frac{d^2y}{dx^2} + p(x)\frac{dy}{dx} + q(x)y = r(x)$$

Let $y'' + py' + qy = 0$ be a homogeneous linear equation of second order (right hand side is equal to zero) where p and q are constants.

The equation $m^2 + pm + q = 0$ is called the **characteristic auxiliary equation**.

⑦ If characteristic equation has two distinct real roots then, $y = c_1e^{m_1x} + c_2e^{m_2x}$ is the general solution of $y'' + py' + qy = 0$.

⑦ If characteristic equation has a real double root then, $y = c_1e^{mx} + c_2xe^{mx}$ is the general solution of $y'' + py' + qy = 0$.

⑦ If characteristic equation has complex roots then, $y = e^{\alpha x} (c_1 \cos \beta x + c_2 \sin \beta x)$ is the general solution of $y'' + py' + qy = 0$.

Let $y'' + py' + qy = r(x)$ be a non-homogeneous linear equation of second order (right hand side is different from zero) where p and q are real numbers.

⑦ If the right hand side of the equation $y'' + py' + qy = r(x)$ has the form $r(x) = Pe^{\alpha x}$ where P is a polynomial, then the particular solution will be

$$y^* = x^k Q_n(x)e^{\alpha x}, Q_n = a_0x^n + a_1x^{n-1} + a_2x^{n-2} + \dots + a_n,$$

where k - is the number of real roots of the associated homogeneous equation that equals to α ;

α is the coefficient of x in $e^{\alpha x}$ in the right hand side and n is the degree of $Q(x)$ that is the same as the degree of $P(x)$ for $r(x)$.

⑦ If the right hand side of the equation $y'' + py' + qy = r(x)$ is of the form $r(x) = Pe^{\alpha x} \cos \beta x + Qe^{\alpha x} \sin \beta x$ where P and Q are polynomials, two cases arise:

$\alpha + i\beta$ is not a root of characteristic equation.

Here, the particular solution will be

$$y^* = Ue^{\alpha x} \cos \beta x + Ve^{\alpha x} \sin \beta x$$

$\alpha + i\beta$ is a root of characteristic equation;

Then, the particular solution is

$$y^* = x [Ue^{\alpha x} \cos \beta x + Ve^{\alpha x} \sin \beta x].$$

In all cases, U and V are polynomials of degree that is equal to the highest degree of P and Q .

Alternative method: Variation of parameters

We know that the general solution of the characteristic equation associated with the equation $y'' + py' + qy = r(x)$ is found to be $\bar{y} = c_1y_1(x) + c_2y_2(x)$.

From $\bar{y} = c_1y_1(x) + c_2y_2(x)$, we can get particular solution y^* as follows:

- ⑦ We determine $W(y_1, y_2)$ known as **Wronskian** of two functions y_1 and y_2 defined by

$$W(y_1, y_2) = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} \neq 0, \text{ since } y_1 \text{ and } y_2 \text{ are linearly independent.}$$

- ⑦ We find out $v_1 = \int \frac{-y_2 r(x)}{W(y_1, y_2)} dx$, and $v_2 = \int \frac{y_1 r(x)}{W(y_1, y_2)} dx$ where $r(x)$ is the right hand side of the given equation.

Then, particular solution y^* is given by

$$y^* = v_1(x)y_1(x) + v_2(x)y_2(x).$$

Therefore, the general solution is $y = \bar{y} + y^*$

$$\text{Or } y = c_1y_1(x) + c_2y_2(x) + v_1(x)y_1(x) + v_2(x)y_2(x)$$

4. Applications


There are a number of well-known applications of first order equations which provide classic prototypes for mathematical

modeling. These mainly rely on the interpretation of $\frac{dy}{dt}$ as a rate of change of a function y with respect to time t . In everyday life, there are many examples of the importance of rates of change – speed of moving particles, growth and decay of populations and materials, heat flow, fluid flow, and so on. In each case, we can construct models of varying degrees of sophistication to describe given situations.

End of unit assessment answers Learner's Book page 325

1. a) $\frac{x^2}{2} + 1$ b) $\frac{1}{2}(x^2 + 1) + \ln x$ c) $-\cos - \frac{1}{\pi}x + 1$
 d) $ce^{-x} + \frac{1}{3}e^{2x}$ e) $e^x + ce^{-2x}$ f) $\frac{x^3 + c}{x - 1}$
 g) $cx^2 - \frac{x^2}{2}e^{-2x}$
2. a) $e^{3y} = \frac{3}{2}e^{2x} + c$ b) $y = cx^xe^{-x}$
 c) $(y^2 - 1)^2 = cx$ d) $y = x(cx - 3)$
 e) $x^2 + y^2 = kx$ f) $(2y - x)^4 = c(x + y)$
 g) $y = \frac{1}{5}e^{2x} + ce^{-3x}$ h) $y = \sqrt{1 - x^2} + c(1 - x^2)$
3. a) $\ln(x^2y) = 2x - y - 1$ b) $y = x(2 - \ln x)$
 c) $y = x\sqrt{8x + 1}$ d) $y = e^x(x - 1)$
 e) $y = \tan x + 2 \sec x$ f) $y = c_1e^{-2x} + c_2e^x$
4. a) $y = c_1e^{-2x} + c_2e^x$ b) $y = e^{-2x}(c_1 \cos x + c_2 \sin x)$
 c) $y = c_1e^{2x} + c_2xe^{2x}$ d) $c_1 \cos 2x + c_2 \sin 2x$
 e) $y = c_1e^{-3x} + c_2e^{3x}$
5. a) $y = c_1e^{-x} + c_2e^{-3x} + \frac{1}{8}(11 - 4x)$
 b) $y = c_1 \cos 2x + c_2 \sin 2x - \frac{e^{2x}}{4}(x + 2)$
 c) $y = e^{-x}(c_1 \cos x + c_2 \sin x) - \frac{14}{85} \cos 3x + \frac{12}{85} \sin 3x$
 d) $y = c_1 \cos 2x + c_2 \sin 2x + \frac{1}{10}e^{-x}(2 \cos x - \sin x)$
 e) $y = c_1 \cos 2x + c_2 \sin 2x + \frac{1}{5}e^x - \frac{x}{4} \cos 2x$
 f) $y = c_1e^x + c_2e^{3x} + \frac{1}{2}e^{3x}(x^2 - x) + \frac{3}{8}e^{3x}(\sin 2x - \cos 2x)$

6. a) $\frac{2}{9}\left(e^{2x} - e^{-\frac{5x}{2}}\right)$ b) $-e^{\pi-2x} \sin 3x$
7. a) $\frac{1}{4}(e^x - e^{-3x})$ b) $-\frac{1}{4} \cos 2x - \frac{\pi}{16} \sin 2x + \frac{1}{4}x + \frac{1}{4}$
- c) $\frac{2}{3} \sin x - \frac{1}{3} \sin 2x$ d) $\frac{1}{6}e^{3x} - \frac{3}{2}e^x + x + \frac{4}{3}$
- e) $y = \frac{4}{3}e^{5x} - \frac{10}{3}e^{2x} - \frac{1}{3}xe^{2x} + 2$
- f) $y = 2e^{\frac{3}{2}x} - 2e^{2x} + \frac{3}{29}e^x(3 \sin x - 7 \cos x)$
- g) $y = e^x(3 \cos x + \sin x) - e^x \cos 2x$
8. m is mass and t is time.
- $$\frac{dm}{dt} = -km, k \text{ is a constant.}$$
9. $\frac{dP}{dt} = kP(1,500 - P), k$ is a constant.
10. h is height in cm, t is time in days, $\frac{dh}{dt} = 0.25h, h = 2$ when $t = 0$.
11. a) $q(t) = EC + (q_0 - EC)e^{-\frac{t}{RC}}$ b) EC
- c) $-RC \ln\left(\frac{0.01EC}{q_0 - EC}\right)$
12. $47.22^\circ C$
13. $77.9^\circ C$





Intersection and Sum of Subspaces

Learner's Book pages 329 - 344

Key unit competence

Relate the sum and the intersection of subspaces of a vector space by the dimension formula.

Vocabulary or key words concepts

Dimension: Number of vectors of the basis of a vector space (or a subspace).

Grassmann's formula: Relation connecting dimensions of subspaces.

Guidance on the problem statement

From $H = \{(a - 2b, 3a + b, 2a + b) : a, b \in \mathbb{R}\}$ and

$K = \{(3b, b, 2b) : b \in \mathbb{R}\}$, how can one find the sum and intersection of H and K . Different subspaces can be added to make a new subspace. Also, we can find the intersection of $H \cap K$ and the result is also a subspace.

List of lessons

No	Lesson title	Number of periods
1	Definition of subspaces	4
2	Intersection of subspaces	2
3	Dimension of intersection of subspaces	2
4	Sum of subspaces	2
5	Dimension of sum of subspaces	2
6	Grassmann's formula of dimension for subspaces	2
Total periods		14

Lesson development

Lesson 6.1. Definition of subspaces

Learning objectives

Through examples, learners should be able to verify that a subset V of \mathbb{R}^n is a subspace of \mathbb{R}^n accurately.

Prerequisites

- ✔ Vector space

Teaching Aids

Exercise book and pen



Activity 6.1 Learner's Book page 330

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $(2x, 0, 5x) = (0, 0, 0)$

$$\begin{cases} 2x = 0 \\ 5x = 0 \end{cases} \Rightarrow x = 0$$

Thus, the value of x is 0.

2. $\alpha\vec{u} + \beta\vec{v} = \alpha(2a, 0, 5a) + \beta(2b, 0, 5b)$
 $= (2\alpha a, 0, 5\alpha a) + (2\beta b, 0, 5\beta b) = (2\alpha a + 2\beta b, 0, 5\alpha a + 5\beta b)$
 $= (2(\alpha a + \beta b), 0, 5(\alpha a + \beta b)) = (2x, 0, 5x) \text{ for } x = \alpha a + \beta b$

Hence, $\alpha\vec{u} + \beta\vec{v} \in V$

3. From results in 1) and 2) and since V is a subset of \mathbb{R}^3 , we conclude that V is a sub-vector space.

Synthesis

A subset V of \mathbb{R}^n is called a **sub-vector space**, or just a **subspace** of \mathbb{R}^n if it has the following properties:

- ① The null vector belongs to V .
- ② V is closed under vector addition, i.e if $\vec{u}, \vec{v} \in V$ then $\vec{u} + \vec{v} \in V$.
- ③ V is closed under scalar multiplication, i.e if $\alpha \in \mathbb{R}, \vec{u} \in V, \alpha\vec{u} \in V$.

Exercise 6.1 Learner's Book page 331

1. a) No, S is not closed under multiplication:

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix} \in S \text{ but } -\begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} -1 \\ -1 \end{bmatrix} \notin S$$

- b) Yes, all properties are verified.
2. a) This is a subspace. It contains $(1, 0, 0)$ and $(2, 1, 0)$.
- b) This is a subspace. It contains $(2, 1, 0)$ and $(3, 0, -3)$.
- c) This is not a subspace. It doesn't contain $(0, 0, 0)$.
3. From results in 1) and 2) and since V is a subset of \mathbb{R}^3 , we conclude that V is a sub-vector space.

Lesson 6.2. Intersection of subspaces**Learning objectives**

Given two subspaces, learners should be able to find their intersection and verify that this intersection is also a subspace correctly.

Prerequisites

- ① Subspace properties

Teaching Aids

Exercise book and pen



Activity 6.2 Learner's Book page 332

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$\begin{cases} 2x - y + 3z = 0 \\ x + y + z = 0 \end{cases}$$

$$3x + 4z = 0 \Rightarrow x = -\frac{4}{3}z$$

$$y = -x - z = -\left(-\frac{4}{3}z\right) - z = \frac{4}{3}z - z = \frac{1}{3}z$$

Then,

$$H \cap K = \left\{ \left(-\frac{4}{3}z, \frac{1}{3}z, z \right) : z \in \mathbb{R} \right\} \text{ or}$$

$$H \cap K = \{(-4x, x, 3x) : x \in \mathbb{R}\}$$

Synthesis

Let U and W be subspaces of a vector space V . The intersection of U and W , written $U \cap W$, consists of all vectors \vec{u} where $\vec{u} \in U$ and $\vec{u} \in W$.

Any intersection of subspaces of a vector space V is a subspace of V .

Properties:

- ✔ For any two subspaces U and W , $U \cap W = W \cap U$
- ✔ If U and W are subspaces of a vector space V , then $U \cap W$ is also a subspace of V .

Exercise 6.2 Learner's Book page 333

1. $U \cap W = \left\{ \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix} : a \in \mathbb{R} \right\}$
2. $H \cap K = \{ \text{functions } f \text{ on } \mathbb{R} : f(2) = f(1) = 0 \}$
3. $U \cap V = \{(0, 0)\}$
4. $U_1 \cap U_2 = \{(0, y, 0) : y \in \mathbb{R}\}$
5. $U_1 \cap U_2 = \{(0, 0, 0)\}$

Lesson 6.3. Dimensions of intersection of subspaces**Learning objectives**

Given two subspaces, learners should be able to find their intersection and the dimension of the intersection accurately.

Prerequisites

- ✔ Intersection of two subspaces.

Teaching Aids

Exercise book and pen

Activity 6.3 Learner's Book page 334

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

a) $U \cap W$

$$\begin{cases} x = x \\ y = y \\ 0 = z \end{cases}$$

$$b) U \cap W = \{(x, y, 0) : x, y \in \mathbb{R}\}$$

$$U \cap W = \{(x, y, 0) : x, y \in \mathbb{R}\}$$

$$\begin{aligned}(x, y, 0) &= (x, 0, 0) + (0, y, 0) \\ &= x(1, 0, 0) + y(0, 1, 0)\end{aligned}$$

The vectors $(1, 0, 0)$ and $(0, 1, 0)$ are linearly independent. Then basis of $U \cap W$ is

$$\{(1, 0, 0), (0, 1, 0)\} \text{ and hence } \dim(U \cap W) = 2.$$

Synthesis

A finite set S of vectors in a vector space V is called a basis for V provided that;

- ① The vectors in S are linearly independent.
 - ② The vector in S span V (or S is a generating set of V).
- The unique number of vectors in each basis for V is called the dimension of V and is denoted by $\dim(V)$.

The dimension of $U \cap W$ is the number of vectors of the basis for $U \cap W$.

Exercise 6.3 Learner's Book page 336

1. $\dim(U \cap W) = 1$
2. $\dim(U \cap W) = 1$
3. $\dim(H \cap K) = 2$

Lesson 6.4. Sum of subspaces

Learning objectives

Given two subspaces, learners should be able to find their sum and verify if the sum is a subspace accurately.

Prerequisites

- ① Properties of subspaces

Teaching Aids

Exercise book and pen



Activity 6.4 Learner's Book page 336

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

For $U = \{(a, 0, c) : a, c \in \mathbb{R}\}$ and $W = \{(0, b, b) : b \in \mathbb{R}\}$

$$\begin{aligned} U+W &= \{(a, 0, c) + (0, b, b) : a, b, c \in \mathbb{R}\} \\ &= \{(a, b, c+b) : a, b, c \in \mathbb{R}\} \\ &= \{(a, b, d) : a, b, d \in \mathbb{R}\} \end{aligned}$$

Clearly, $(0, 0, 0) \in U+W$. Let

$\vec{u} = (x, y, z), \vec{w} = (x', y', z') \in U+W$ and $\alpha, \beta \in U+W$

$$\begin{aligned} \alpha\vec{u} + \beta\vec{w} &= \alpha(x, y, z) + \beta(x', y', z') \\ &= (\alpha x, \alpha y, \alpha z) + (\beta x', \beta y', \beta z') \\ &= (\alpha x + \beta x', \alpha y + \beta y', \alpha z + \beta z') \\ &= (a, b, c) \in U+W \\ &\quad \text{for } a = \alpha x + \beta x', b = \alpha y + \beta y', c = \alpha z + \beta z' \end{aligned}$$

Thus, $U+W$ is a subspace of \mathbb{R}^3

Synthesis

Let U and W be subspaces of a vector space V . The sum of U and W , written $U+W$, consists of all sums $x+y$ where $x \in U$ and $y \in W$.

- ④ The sum $U+W$ of the subspaces U and V is also a subspace of V .
- ④ and W_2 are subspace of V , then W_1+W_2 is the smallest subspace that contains both W_1 and W_2 .

Exercise 6.4 Learner's Book page 337

1. Let $\vec{v} \in U + W$. Then $\vec{v} = \vec{u} + \vec{w}$, $\vec{u} \in U$ and $\vec{w} \in W$.
 Since $\{u_i\}$ generates U , \vec{u} is a linear combination of \vec{u}_i 's; and since $\{\vec{w}_j\}$ generates W , \vec{w} is a linear combination of \vec{w}_j 's.

Thus

$$\vec{v} = \vec{u} + \vec{w} = a_1\vec{u}_{i_1} + a_2\vec{u}_{i_2} + \dots + a_n\vec{u}_{i_n} + b_1\vec{w}_{j_1} + b_2\vec{w}_{j_2} + \dots + b_m\vec{w}_{j_m}$$

and so $\{\vec{u}_i, \vec{w}_j\}$ generates $U + W$.

2. $U + W = \left\{ \begin{pmatrix} e & b \\ d & 0 \end{pmatrix} : b, d, e \in \mathbb{R} \right\}$
3. $\{(2a - b, 3a + 4b, 5a + 3b) : a, b \in \mathbb{R}\}$

Lesson 6.5. Dimension of sum of subspaces

Learning objectives

Given two subspaces, learners should be able to find their sum and the dimension of the sum accurately.

Prerequisites

- ✔ Sum of subspaces.

Teaching Aids

Exercise book and pen



Activity 6.5 Learner's Book page 338

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \quad U + W = \{(a, 0, 0) + (0, b, 0) : a, b \in \mathbb{R}\} = \{(a, b, 0) : a, b \in \mathbb{R}\}$$

$$2. \quad U + W = \{(a, b, 0) : a, b \in \mathbb{R}\}$$

$$(a, b, 0) = (a, 0, 0) + (0, b, 0)$$

$$= a(1, 0, 0) + b(0, 1, 0)$$

The vectors $(1, 0, 0)$ and $(0, 1, 0)$ are linearly independent. Then, basis of $U + W$ is

$\{(1, 0, 0), (0, 1, 0)\}$ and hence $\dim(U + W) = 2$.

Synthesis

A finite set S of linearly independent vectors in the sum $U + V$ is called a basis for $U + V$ and the number of vectors in set S is the dimension of $U + V$.

Exercise 6.5 Learner's Book page 339

$$1. \quad \dim(H + K) = 2 \qquad 2. \quad \dim(U + V) = 2$$

$$3. \quad \dim(U_1 + U_2) = 2$$

Lesson 6.6. Grassmann's formula of dimension for subspaces**Learning objectives**

Given two subspaces, learners should be able to use Grassmann's formula to find the dimension of the sum or intersection correctly.

Prerequisites

- ✔ Sum of subspaces.
- ✔ Intersection of subspaces.

Teaching Aids

Exercise book and pen



Activity 6.6 Learner's Book page 340

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. a) $\dim(F) = 2$ and $\dim(G) = 2$
 b) $\dim(F) + \dim(G) = 2 + 2 = 4$
 c) $F \cap G = \{(0, 0, z) : z \in \mathbb{R}\}$ and $\dim(F \cap G) = 1$
 d) $\dim(F) + \dim(G) - \dim(F \cap G) = 4 - 1 = 3$
 e) $F + G = \{(x, y, z) : x, y, z \in \mathbb{R}\}$ and $\dim(F + G) = 3$
2. From results in d) and e),

$$\dim(F + G) = \dim(F) + \dim(G) - \dim(F \cap G)$$

Synthesis

- ✔ If $(\mathbb{R}, F, +)$ and $(\mathbb{R}, G, +)$ are two sub-vector spaces of $(\mathbb{R}, E, +)$, we have,

$$\dim(F + G) = \dim(F) + \dim(G) - \dim(F \cap G).$$

- ✔ If $\dim(F \cap G) = 0$, then

$$\dim(F + G) = \dim(F) + \dim(G).$$

In this case, F and G are said to be **complementary** and the sum $F + G$ is said to be a direct sum; and it is denoted by $F \oplus G$. Otherwise, F and G are said to be **supplementary**.

Exercise 6.6 Learner's Book page 342

1. $\dim(V) = 3, \dim(W) = 2, \dim(V \cap W) = 1$
 $\dim(V + W) = \dim(V) + \dim(W) - \dim(V \cap W) = 3 + 2 - 1 = 4$
2. \mathbb{R}^3 cannot be uniquely represented as a direct sum of W_1 and W_2 .
3. $F = W_1 + W_2$ is a direct sum. i.e, $F = W_1 \oplus W_2$.
4. No, since $\dim(F \cap G) = 3 \neq 0$.
5. Since U is not equal to W , the basis for U must have at least one vector linearly independent from U , so $\dim(U + W)$ is at least 4. But they are subspaces of \mathbb{R}^4 , so $\dim(U + W) = 4$. Using the fact that $\dim(U + W) = \dim(U) + \dim(W) - \dim(U \cap W)$.
Then,
 $4 = 3 + 3 - \dim(U \cap W) \Rightarrow \dim(U \cap W) = 6 - 4 = 2$.

Summary of the unit

1. Definition

If $(\mathbb{R}, F, +)$ is a subspace of $(\mathbb{R}, E, +)$, then

$$\text{⑦ } F \subset E$$

$$\text{⑦ } \vec{0} \in F$$

$$\text{⑦ } \vec{u}, \vec{v} \in F, \alpha, \beta \in \mathbb{R}; \quad \alpha\vec{u} + \beta\vec{v} \in F$$

2. Intersection and sum of two vector spaces

Let U and W be subspaces of a vector space V . The intersection of U and W , written $U \cap W$, consists of all vectors \vec{u} where $\vec{u} \in U$ and $\vec{u} \in W$.

Any intersection of subspaces of a vector space V is a subspace of V . W_1 and W_2 are subspaces of V , then $W_1 \cup W_2$ is a subspace $\Leftrightarrow W_1 \subseteq W_2$ or $W_2 \subseteq W_1$.

If F and G are two sub-vector spaces of E , then, the sum of F and G is also a sub-vector space of E . It is denoted as $F + G = \{x + y, x \in F, y \in G\}$.

Grassmann's formula of dimensions.

If $(\mathbb{R}, F, +)$ and $(\mathbb{R}, G, +)$ are two sub-vector spaces of $(\mathbb{R}, E, +)$, we have,

$$\dim(F + G) = \dim(F) + \dim(G) - \dim(F \cap G).$$

Remark


If $\dim(F \cap G) = 0$, then $\dim(F + G) = \dim(F) + \dim(G)$. In this case, F and G are said to be **complementary** and the sum $F + G$ is said to be a **direct sum**; and it is denoted by $F \oplus G$.

Otherwise, F and G are said to be **supplementary**.

End of Unit Assessment answers Learner's Book page 344

1. a) Yes, this is a plane through origin.
 b) No, this does not contain the origin.
 c) Yes, this is just the zero point.
 d) No, this is a conic which is not closed under addition.
2. $\dim(E) = 1$, $\dim(F) = 2$
3. $\dim(W) = 3$
4. Since U and W are distinct, $U + W$ properly contains U and W ; hence $\dim(U + W) > 4$. Since $\dim(V) = 6$, $\dim(U + W)$ cannot be greater than 6.
 Hence, there are two possibilities:
 a) $\dim(U + W) = 5 \Leftrightarrow 5 = 4 + 4 - \dim(U \cap W) \Rightarrow \dim(U \cap W) = 3$
 b) $\dim(U + W) = 6 \Leftrightarrow 6 = 4 + 4 - \dim(U \cap W) \Rightarrow \dim(U \cap W) = 2$
5. a) $\dim(U + W) = 3$ b) $\dim(U \cap W) = 1$
6. The set of the symmetric matrices W_1 and the set of the skew symmetric matrices W_2 are both subspaces of $M_{n \times n}$.
 $A \in W_1 \cap W_2, A = A^t = -A^t \Rightarrow A = 0, \therefore W_1 \cap W_2 = \{0\}$
 Let $\begin{cases} B = \frac{1}{2}(A + A^t) \\ C = \frac{1}{2}(A - A^t) \end{cases}$
 Then,
 $B \in W_1, C \in W_2, \therefore M_{n \times n} = W_1 \oplus W_2$
7. Yes, since $\dim(F \cap G) = 0$.





Transformation of Matrices

Learner's Book pages 345 - 385

Key unit competence

Transform matrices to an echelon form or to diagonal matrix and use the results to solve simultaneous linear equations or to calculate the n^{th} power of a matrix.

Vocabulary or key words concepts

Elementary row/column operations: Operations performed on row/column of a matrix (addition, scalar multiplication, and interchanging rows/columns) to obtain a new matrix.

Characteristic equation: Polynomial $|A - \lambda I| = 0$, $\lambda \in \mathbb{R}$ where A is a given matrix and I is identity matrix of the same order as A .

Eigenvalue: The real number λ that is a root in the characteristic polynomial $|A - \lambda I| = 0$, $\lambda \in \mathbb{R}$

Eigenvector: The vector \vec{u} such that $(A - \lambda I)\vec{u} = \vec{0}$.

Row echelon form: Matrix is in row echelon form when the first non-zero element in each row (called the leading entry) is 1 and this leading entry is in a column to the right of the leading entry in the previous row. Rows with all zero elements, if any, are below rows having a non-zero element.

Reduced row echelon form: A matrix is in reduced row echelon form when it is in row echelon form and the leading entry in each row is the only non-zero entry in its column.

Guidance on the problem statement

- ⑦ Given encoding matrix $A = \begin{bmatrix} -1 & 2 \\ 2 & -3 \end{bmatrix}$, can you decode the message 21, -22, 17, -18, 28, -42, 31, -39 and read the message?
- ⑦ Matrices and their inverse are used by programmers for coding or encrypting a message. Matrices are applied in the study of electrical circuits, quantum mechanics and optics.

List of lessons

No	Lesson title	Number of periods
1	Kernel and range	4
2	Elementary row/column operations	3
3	Eigenvalues and eigenvectors	4
4	Diagonalisation of a matrix	3
5	Echelon matrix	3
6	Inverse matrix	3
7	Rank of a matrix	3
8	Solving system of linear equations	3
9	Power of a matrix	3
Total periods		29

Lesson development

Lesson 7.1. Kernel and range

Learning objectives

By the end of this lesson, learners should be able to find kernel and range of a linear transformation.

Prerequisites

- ✔ Operation on vectors.
- ✔ Operations on matrices.

Teaching Aids

Exercise book and pen

Activity 7.1 Learner's Book page 347



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$(3x + y + 2, 3x - y + 1) = (0, 0)$$

$$\Rightarrow \begin{cases} 3x + y + 2 = 0 \\ 3x - y + 1 = 0 \end{cases}$$

$$\begin{cases} 3x + y + 2 = 0 \\ 3x - y + 1 = 0 \end{cases} \qquad 3\left(-\frac{1}{2}\right) + y + 2 = 0$$

$$6x + 3 = 0 \Rightarrow x = -\frac{1}{2} \qquad \Rightarrow y = -2 + \frac{3}{2} = -\frac{1}{2}$$

$$\text{Thus, } (x, y) = \left(-\frac{1}{2}, -\frac{1}{2}\right)$$

Synthesis

- ⑦ The **kernel** of a linear mapping $f: E \rightarrow F$ denoted $\text{Ker}(f)$ is a subset of E whose image by f is 0-vector of F . i.e, $\text{Ker}(f) = \{v \in E : f(v) = 0\}$.
- ⑦ The **nullity** of f denoted $n(f)$ is the dimension of $\text{Ker}(f)$. i.e, $n(f) = \dim \text{Ker}(f)$.
- ⑦ The **image** or **range** of a linear mapping $f: E \rightarrow F$ is the set of vectors in F to which points in E are mapped on. i.e, $\text{Im } f = \{u \in F : f(v) = u\}, v \in E$.
- ⑦ The **rank** of f denoted $\text{rank}(f)$ or $r(f)$ is the dimension of image of f . i.e, $\text{rank}(f) = \dim(\text{Im } f)$.
- ⑦ If $f: E \rightarrow F$, $\dim[\text{Ker}(f)] + \dim[\text{range}(f)] = \dim(E)$.

Exercise 7.1 Learner's Book page 350

1. a) $\text{Im } F = \{(a, b, c) : c = 0\} = xy\text{-plane}$.
b) $\text{Ker } F = \{(a, b, c) : a = 0, b = 0\} = z\text{-axis}$.
2. a) Basis is $\{(1, 0, 1), (0, 1, -1)\}$ and dimension is 2.
b) Basis is $\{(3, -1, 1)\}$ and dimension is 1.
3. a) Basis is $\{(1, 1, 1), (0, 1, 2)\}$ and dimension is 2.
b) Basis is $\{(2, 1, -1, 0), (1, 2, 0, 1)\}$ and dimension is 2.

Lesson 7.2. Elementary row/column operations

Learning objectives

Given a matrix, learners should be able to use row/column operations to transform correctly.

Prerequisites

- ⑦ Adding row/column of a matrix.
- ⑦ Multiplying a row/column of a matrix by a real number.

Teaching Aids

Exercise book, pen and calculator

**Activity 7.2 Learner's Book page 351**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$1. \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 3 \\ 3 & 2 & 1 \end{pmatrix}$$

$$2. \begin{pmatrix} 1 & 0 & 2 \\ 0 & 1 & 3 \\ 0 & 2 & -5 \end{pmatrix}$$

$$3. \begin{pmatrix} 1 & 2 & 0 \\ 0 & 3 & 1 \\ 0 & -5 & 2 \end{pmatrix}$$

$$4. \begin{pmatrix} 1 & 2 & -\frac{2}{3} \\ 0 & 3 & 0 \\ 0 & -5 & \frac{11}{3} \end{pmatrix}$$

Synthesis

Common row/column and their notations are;

Operation description	Notation
<u>Row operations</u>	
1. Interchange row i and j	$\rightarrow r_i \leftrightarrow r_j$
2. Multiply row i by $s \neq 0$	$\rightarrow \text{new } r_i \rightarrow sr_i$
3. Add s times row i to row j	$\rightarrow \text{new } r_j \rightarrow r_j + sr_i$
<u>Column operations</u>	
1. Interchange column i and j	$\rightarrow c_i \leftrightarrow c_j$
2. Multiply column i by $s \neq 0$	$\rightarrow \text{new } c_i \rightarrow sc_i$
3. Add s times column i to column j	$\rightarrow \text{new } c_j \rightarrow c_j + sc_i$

Exercise 7.2 Learner's Book page 353

$$1. \quad a) \begin{pmatrix} 1 & 2 & -3 & 0 \\ 0 & 0 & 4 & 2 \\ 3 & 6 & -4 & 3 \end{pmatrix} \quad b) \begin{pmatrix} 1 & 2 & -3 & 0 \\ 0 & 0 & 4 & 2 \\ 0 & 0 & 5 & 3 \end{pmatrix}$$

$$c) \begin{pmatrix} 1 & 2 & -3 & 0 \\ 0 & 0 & 4 & 2 \\ 0 & 0 & 0 & 2 \end{pmatrix}$$

$$2. \quad a) \begin{pmatrix} 6 & 9 & 0 & 7 & -2 \\ 0 & 0 & 3 & 2 & 5 \\ 0 & 0 & 0 & 0 & 2 \end{pmatrix} \quad b) \begin{pmatrix} 6 & 9 & 0 & 7 & 0 \\ 0 & 0 & 3 & 2 & 5 \\ 0 & 0 & 0 & 0 & 2 \end{pmatrix}$$

$$c) \begin{pmatrix} 6 & 9 & 0 & 7 & 0 \\ 0 & 0 & 6 & 4 & 0 \\ 0 & 0 & 0 & 0 & 2 \end{pmatrix}$$

Lesson 7.3. Eigenvalues and eigenvectors**Learning objectives**

Given a matrix, learners should be able to find eigenvalues and eigenvectors accurately.

Prerequisites

- ✔ Operation on matrices.
- ✔ Matrix determinant.
- ✔ Solving equation of second/third degree.

Teaching Aids

Exercise book, pen and calculator

**Activity 7.3 Learner's Book page 354**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence

- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$\begin{aligned}
 1. \quad \det(A - \lambda I) &= \det \left(\begin{pmatrix} 4 & 2 \\ 3 & -1 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right) \\
 &= \begin{vmatrix} 4 - \lambda & 2 \\ 3 & -1 - \lambda \end{vmatrix} \\
 &= (4 - \lambda)(-1 - \lambda) - 6 \\
 &= \lambda^2 - 3\lambda - 10
 \end{aligned}$$

$$2. \quad \lambda^2 - 3\lambda - 10 = 0 \Leftrightarrow (\lambda + 2)(\lambda - 5) = 0 \Rightarrow \lambda = -2 \text{ or } \lambda = 5$$

$$3. \quad (A - \lambda I)\vec{u} = \vec{0}$$

For $\lambda = -2$

$$\left(\begin{pmatrix} 4 & 2 \\ 3 & -1 \end{pmatrix} + 2 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right) \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} 6 & 2 \\ 3 & 1 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{cases} 6u_1 + 2u_2 = 0 \\ 3u_1 + u_2 = 0 \end{cases} \Rightarrow u_2 = -3u_1$$

Thus, $\vec{u} = k \begin{pmatrix} 1 \\ -3 \end{pmatrix}$, $k \in \mathbb{R}_0$

For $\lambda = 5$

$$\left(\begin{pmatrix} 4 & 2 \\ 3 & -1 \end{pmatrix} - 5 \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right) \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{pmatrix} -1 & 2 \\ 3 & -6 \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix} \quad \begin{cases} -u_1 + 2u_2 = 0 \\ 3u_1 - 6u_2 = 0 \end{cases} \Rightarrow u_1 = 2u_2$$

Thus, $\vec{u} = k \begin{pmatrix} 2 \\ 1 \end{pmatrix}$, $k \in \mathbb{R}_0$

Synthesis

The eigenvalues of square matrix A , are the roots of the polynomial $\det(A - \lambda I)$. The homogeneous system $(A - \lambda I)\vec{u} = \vec{0}$ gives the eigenvector \vec{u} associated with eigenvalue λ .

Cayley and Hamilton theorem

The Cayley–Hamilton theorem (named after the mathematicians Arthur Cayley and William Rowan Hamilton) states that every square matrix over a commutative ring (such as the real or complex field) satisfies its own characteristic equation.

Note that an eigenvector cannot be 0 , but an eigenvalue can be 0 . If 0 is an eigenvalue of A , there must be some non-trivial vector \vec{u} for which $(A - 0I)\vec{u} = \vec{0}$.

Exercise 7.3 Learner's Book page 360

- Eigenvalues: 7 and -4 , eigenvectors: $\vec{u} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ and $\vec{v} = \begin{pmatrix} 2 \\ -3 \end{pmatrix}$
 - Eigenvalues: 7 and -4 , eigenvectors: $\vec{u} = \begin{pmatrix} 3 \\ 1 \end{pmatrix}$ and $\vec{v} = \begin{pmatrix} 2 \\ -3 \end{pmatrix}$
 - No eigenvalues, no eigenvectors
- $\lambda_1 = 2, u = (1, -1, 0), v = (1, 0, -1); \lambda_2 = 6, w = (1, 2, 1)$
 - $\lambda_1 = 3, u = (1, 1, 0), v = (1, 0, 1); \lambda_2 = , w = (2, -1, 1)$ ()
 - $\lambda = 1, u = (1, 0, 0), v = (0, 0, 1)$
- Characteristic equation of matrix A is

$$|A - \lambda I| = 0. \quad (1)$$
 Characteristic equation of matrix A' is

$$|A' - \lambda I| = 0. \quad (2)$$

Clearly, both (1) and (2) are the same, as we know that $|A| = |A'|$.

Therefore, A and A' have the same eigenvalues.

4. Let us consider the triangular matrix,

$$A = \begin{pmatrix} a_{11} & 0 & \cdots & 0 \\ a_{21} & a_{22} & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{pmatrix}$$

Characteristic equation is $|A - \lambda I| = 0$.

$$\Leftrightarrow \begin{vmatrix} a_{11} - \lambda & 0 & \cdots & 0 \\ a_{21} & a_{22} - \lambda & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} - \lambda \end{vmatrix} = 0$$

$$\Leftrightarrow (a_{11} - \lambda)(a_{22} - \lambda) \cdots (a_{nn} - \lambda) = 0$$

Therefore, $\lambda = a_{11}, a_{22}, \dots, a_{nn}$ are the elements of diagonal entries.

5. $\frac{1}{9} \begin{pmatrix} 7 & 2 & -10 \\ -2 & 2 & -1 \\ -1 & 1 & 4 \end{pmatrix}$

6. $A^5 - 4A^4 - 7A^3 + 11A^2 - A - 10I = A + 5I$

$$\begin{aligned} &= (A^5 - 4A^4 - 5A^3) - 2A^3 + 11A^2 - A - 10I \\ &= A^3(A^2 - 4A - 5) - 2A^3 + 11A^2 - A - 10I \\ &= 0 - 2A^3 + 11A^2 - A - 10I \\ &= -(2A^3 - 8A^2 - 10A) + 3A^2 - 11A - 10I \\ &= 0 + 3A^2 - 11A - 10I \\ &= (3A^2 - 12A - 15I) + A + 5I \\ &= 3(A^2 - 4A - 5I) + A + 5I = A + 5I \end{aligned}$$

Therefore, $A^5 - 4A^4 - 7A^3 + 11A^2 - A - 10I = A + 5I$.

Lesson 7.4. Diagonalisation of a matrix

Learning objectives

Given a matrix, learners should be able to diagonalise that matrix accurately.

Prerequisites

- ✔ Operation on matrices

Teaching Aids

Exercise book, pen and calculator



Activity 7.4 Learner's Book page 361

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. A = \begin{pmatrix} -4 & -6 \\ 3 & 5 \end{pmatrix}$$

Eigenvalues:

$$\begin{aligned} \left| \begin{pmatrix} -4 & -6 \\ 3 & 5 \end{pmatrix} - \lambda \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right| = 0 &\Rightarrow \begin{vmatrix} -4-\lambda & -6 \\ 3 & 5-\lambda \end{vmatrix} = 0 \\ &\Rightarrow (-4-\lambda)(5-\lambda) + 18 = 0 \end{aligned}$$

$$\lambda^2 - \lambda - 2 = 0 \Rightarrow \lambda = -1 \text{ or } \lambda = 2$$

Eigenvalues are -1 and 2

Eigenvectors:

For $\lambda = -1$

$$\left[\begin{pmatrix} -4 & -6 \\ 3 & 5 \end{pmatrix} + \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \right] \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

$$\begin{aligned} \Rightarrow \begin{pmatrix} -3 & -6 \\ 3 & 6 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \begin{cases} -3x - 6y = 0 \\ 3x + 6y = 0 \end{cases} \\ \Rightarrow \begin{pmatrix} -3x - 6y \\ 3x + 6y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \Rightarrow 3x = -6y \Rightarrow x = -2y \end{aligned}$$

Eigenvector associated to $\lambda = -1$ has the form

$$\begin{pmatrix} -2y \\ y \end{pmatrix}, y \in \mathbb{R}_0. \text{ Take } \vec{u} = \begin{pmatrix} -2 \\ 1 \end{pmatrix}$$

For $\lambda = 2$

$$\begin{aligned} \left[\begin{pmatrix} -4 & -6 \\ 3 & 5 \end{pmatrix} - \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix} \right] \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} \\ \Rightarrow \begin{pmatrix} -6 & -6 \\ 3 & 3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \begin{cases} -6x - 6y = 0 \\ 3x + 3y = 0 \end{cases} \\ \Rightarrow \begin{pmatrix} -6x - 6y \\ 3x + 3y \end{pmatrix} &= \begin{pmatrix} 0 \\ 0 \end{pmatrix} & \Rightarrow 3x = -3y \Rightarrow x = -y \end{aligned}$$

Eigenvector associated to $\lambda = 2$ has the form

$$\begin{pmatrix} -y \\ y \end{pmatrix}, y \in \mathbb{R}_0. \text{ Take } \vec{u} = \begin{pmatrix} -1 \\ 1 \end{pmatrix}$$

2. From 1), $P = \begin{pmatrix} -2 & -1 \\ 1 & 1 \end{pmatrix}$

3. $P^{-1} = \frac{1}{-1} \begin{pmatrix} 1 & 1 \\ -1 & -2 \end{pmatrix} = \begin{pmatrix} -1 & -1 \\ 1 & 2 \end{pmatrix}$

4.
$$\begin{aligned} D &= P^{-1}AP = \begin{pmatrix} -1 & -1 \\ 1 & 2 \end{pmatrix} \begin{pmatrix} -4 & -6 \\ 3 & 5 \end{pmatrix} \begin{pmatrix} -2 & -1 \\ 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 4-3 & 6-5 \\ -4+6 & -6+10 \end{pmatrix} \begin{pmatrix} -2 & -1 \\ 1 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} -2 & -1 \\ 1 & 1 \end{pmatrix} \\ &= \begin{pmatrix} -2+1 & -1+1 \\ -4+4 & -2+4 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 2 \end{pmatrix} \end{aligned}$$

Matrix D is a diagonal matrix. Also, elements of the leading diagonal are the eigenvalues obtained in 1).

Synthesis

To diagonalise matrix A , we perform the following steps:

1. Find the eigenvalues.
2. If there is a non-real eigenvalue, the matrix cannot be diagonalised.
3. If all eigenvalues are real, find their associated eigenvectors (they must be linearly independent).
4. If the number of eigenvectors is not equal to the order of matrix A , then this matrix cannot be diagonalised.
5. If the number of eigenvectors is equal to the order of matrix A , form matrix P whose columns are elements of eigenvectors.
6. Find the inverse of P .
7. Find D , diagonal matrix of A by relation; $D = P^{-1}AP$.

Theorem

A $n \times n$ matrix is diagonalisable if and only if it has n linearly independent eigenvectors.

Exercise 7.4 Learner's Book page 364

1. a) $\begin{pmatrix} 8 & 0 \\ 0 & -2 \end{pmatrix}$ b) $\begin{pmatrix} 4 & 0 \\ 0 & 1 \end{pmatrix}$ c) $\begin{pmatrix} 7 & 0 \\ 0 & -3 \end{pmatrix}$
 d) $\begin{pmatrix} 2 & 0 \\ 0 & 1 \end{pmatrix}$ e) $\begin{pmatrix} -5 & 0 & 0 \\ 0 & -5 & 0 \\ 0 & 0 & 16 \end{pmatrix}$
2. a) 2 and -4 b) $S = \{(1,1,0), (0,1,1)\}$
 c) A is not diagonalisable.

Lesson 7.5. Echelon matrix

Learning objectives

Given a matrix and using elementary row/column operations, learners should be able to transform that matrix into its echelon form accurately.

Prerequisites

- ④ Elementary row/column operations.

Teaching Aids

Exercise book, pen and calculator

Activity 7.5 Learner's Book page 364



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$A = \begin{pmatrix} 8 & 3 \\ 1 & 2 \end{pmatrix}$$

Change first element of first row to 1.

$$\underline{r_1 = r_1 - 7r_2} \begin{pmatrix} 1 & -11 \\ 1 & 2 \end{pmatrix}$$

The first non-zero element in second row is 1 but it is not in a column to the right of the other in first row. So, this has to be changed to 0.

$$\underline{r_2 = r_2 - r_1} \begin{pmatrix} 1 & -11 \\ 0 & -13 \end{pmatrix}$$

Now, second element in second row has to be changed to 1.

$$\underline{r_2 = r_2 - \frac{14}{11}r_1} \begin{pmatrix} 1 & -11 \\ 0 & 1 \end{pmatrix}$$

Now, the first two conditions are satisfied.

For the third condition:

The first non-zero element in second row is not the only non-zero entry in its column. So -11 , in first row, has to be changed to 0.

$$\underline{r_1 = r_1 + 11r_2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Now, all conditions are satisfied.

Synthesis

A matrix is in **row echelon form** (ref) when it satisfies the following conditions:

- ④ The first non-zero element in each row, called the **leading entry (or pivot)**, is 1.
- ④ Each leading entry is in a column to the right of the leading entry in the previous row.
- ④ Rows with all zero elements, if any, are below rows having a non-zero element.

A matrix is in **reduced row echelon form** (rref) when it satisfies the following conditions:

- ④ The matrix is in row echelon form (i.e., it satisfies the three conditions listed above).
- ④ The leading entry in each row is the only non-zero entry in its column.

Exercise 7.5 Learner's Book page 367

$$1. \begin{pmatrix} 1 & -1 & 2 & 0 \\ 0 & 1 & 4 & -5 \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 & 6 & 0 \\ 0 & 1 & 4 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$2. \begin{pmatrix} 1 & -2 & 3 & 9 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 2 & 4 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 2 \end{pmatrix}$$

$$3. \begin{pmatrix} 2 & -2 & 4 & -2 \\ 0 & 3 & 6 & 9 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 & 4 & 0 \\ 0 & 1 & 2 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$4. \begin{pmatrix} 1 & -3 & 4 & 10 \\ 0 & 1 & -\frac{8}{7} & -\frac{23}{7} \\ 0 & 0 & 0 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 & \frac{4}{7} & 0 \\ 0 & 1 & -\frac{8}{7} & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

$$5. \begin{pmatrix} 1 & 2 & 3 & -2 & 3 \\ 0 & 1 & 1 & -1 & \frac{4}{3} \\ 0 & 0 & 0 & 1 & \frac{4}{9} \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 & 1 & 0 & \frac{1}{3} \\ 0 & 1 & 1 & 0 & \frac{16}{9} \\ 0 & 0 & 0 & 1 & \frac{4}{9} \end{pmatrix}$$

$$6. \begin{pmatrix} 1 & \tan \theta \\ 0 & 1 \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Lesson 7.6. Matrix inverse

Learning objectives

Given a square matrix and using elementary row/column operations, learners should be able to find the inverse of that matrix correctly.

Prerequisites

- ✔ Use of elementary row operations.
- ✔ Properties of inverse matrix.

Teaching Aids

Exercise book, pen and calculator



Activity 7.6 Learner's Book page 367

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. M = \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 1 & 2 & 4 & 0 & 0 & 1 \end{array} \right)$$

We need to transform matrix A such that all elements of leading diagonal become 1 and other elements become zero

$$\xrightarrow{r_3 \rightarrow r_3 - r_1} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 1 & 3 & -1 & 0 & 1 \end{array} \right) \xrightarrow{r_3 \rightarrow r_3 - r_2} \left(\begin{array}{ccc|ccc} 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 & -1 & 1 \end{array} \right)$$

$$\xrightarrow{r_1 \rightarrow r_1 - r_2} \left(\begin{array}{ccc|ccc} 1 & 0 & -1 & 1 & -1 & 0 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 & -1 & 1 \end{array} \right)$$

$$\xrightarrow{r_1 \rightarrow r_1 + r_3} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & -2 & 1 \\ 0 & 1 & 2 & 0 & 1 & 0 \\ 0 & 0 & 1 & -1 & -1 & 1 \end{array} \right)$$

$$\xrightarrow{r_2 \rightarrow r_2 - 2r_3} \left(\begin{array}{ccc|ccc} 1 & 0 & 0 & 0 & -2 & 1 \\ 0 & 1 & 0 & 2 & 3 & -2 \\ 0 & 0 & 1 & -1 & -1 & 1 \end{array} \right)$$

2. Matrix obtained in 1) is $\begin{pmatrix} 0 & -2 & 1 \\ 2 & 3 & -2 \\ -1 & -1 & 1 \end{pmatrix}$. Multiplying it by the given matrix, gives

$$\begin{pmatrix} 0 & -2 & 1 \\ 2 & 3 & -2 \\ -1 & -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 2 \\ 1 & 2 & 4 \end{pmatrix} = \begin{pmatrix} 0+0+1 & 0-2+2 & 0-4+4 \\ 2+0-2 & 2+3-4 & 2+6-8 \\ -1+0+1 & -1-1+2 & -1-2+4 \end{pmatrix} \\ = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Observation: Multiplying matrix obtained in 1) by matrix A gives identity matrix.

Therefore, the new matrix is the inverse of the matrix A .

Synthesis

To calculate the inverse of A , denoted as A^{-1} , follow these steps:

Construct a matrix of type $M = (A|I)$, that is to say, A is in the left half and the identity matrix I is on the right.

Using elementary row operations, transform the left half, A , to the identity matrix located to the right, and the matrix that results in the right side will be the inverse of matrix.

Exercise 7.6 Learner's Book page 369

1. $\begin{pmatrix} -1 & 2 & -4 \\ 1 & -1 & 3 \\ 0 & 0 & 1 \end{pmatrix}$

2. No inverse

3. $\begin{pmatrix} 1 & -3 & 2 \\ -3 & 3 & -1 \\ 2 & -1 & 0 \end{pmatrix}$

4. No inverse

Lesson 7.7. Rank of matrix

Learning objectives

Given a matrix and by using elementary row/column operations, learners should be able to find the rank of that matrix accurately.

Prerequisites

- ✔ Transformation of matrix using elementary row/column operations.

Teaching Aids

Exercise book, pen and calculator



Activity 7.7 Learner's Book page 369

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \begin{pmatrix} 4 & -6 & 0 \\ -6 & 0 & 1 \\ 0 & 9 & -1 \\ 0 & 1 & 4 \end{pmatrix} \xrightarrow{r_2 \rightarrow r_2 + \frac{3}{2}r_1} \begin{pmatrix} 4 & -6 & 0 \\ 0 & -9 & 1 \\ 0 & 9 & -1 \\ 0 & 1 & 4 \end{pmatrix}$$

$$\left. \begin{matrix} r_3 \rightarrow r_3 + r_2 \\ r_4 \rightarrow r_4 + \frac{1}{9}r_2 \end{matrix} \right\} \begin{pmatrix} 4 & -6 & 0 \\ 0 & -9 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & \frac{37}{9} \end{pmatrix} \xrightarrow{r_3 \leftrightarrow r_4} \begin{pmatrix} 4 & -6 & 0 \\ 0 & -9 & 1 \\ 0 & 0 & \frac{37}{9} \\ 0 & 0 & 0 \end{pmatrix}$$

2. There are three non-zero rows.

Synthesis

To find rank of matrix,

- ④ Transform matrix in its row echelon form using elementary row operations.
- ④ The number of non-zero rows is the rank of matrix.

Exercise 7.7 Learner's Book page 372

- | | |
|-----------|-----------|
| 1. Rank 4 | 2. Rank 3 |
| 3. Rank 3 | 4. Rank 2 |

Lesson 7.8. Solving system of linear equations**Learning objectives**

Given a system of linear equations and by using Gaussian elimination method, learners should be able to find the solution of that system correctly.

Prerequisites

- ④ Elementary row operations.

Teaching Aids

Exercise book, pen and calculator

Activity 7.8 Learner's Book page 373

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. The system
$$\begin{cases} x + y + z = 6 \\ 2x + y - z = 1 \\ 3x + 2y + z = 10 \end{cases} \Leftrightarrow \begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & -1 \\ 3 & 2 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 6 \\ 1 \\ 10 \end{pmatrix}$$

$$\text{Thus, } A = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 1 & -1 \\ 3 & 2 & 1 \end{pmatrix}$$

$$2. \begin{pmatrix} 1 & 1 & 1 & : & 6 \\ 2 & 1 & -1 & : & 1 \\ 3 & 2 & 1 & : & 10 \end{pmatrix}$$

$$3. \text{ From } \begin{pmatrix} 1 & 1 & 1 & : & 6 \\ 2 & 1 & -1 & : & 1 \\ 3 & 2 & 1 & : & 10 \end{pmatrix}, \text{ using elementary row}$$

operations; $r_2 \rightarrow r_2 - 2r_1, r_3 \rightarrow r_3 - 3r_1$, we get

$$\begin{pmatrix} 1 & 1 & 1 & : & 6 \\ 0 & -1 & -3 & : & -11 \\ 0 & -1 & -2 & : & -8 \end{pmatrix}$$

$$\text{Now, } r_3 \rightarrow r_3 - r_2, \text{ yields } \begin{pmatrix} 1 & 1 & 1 & : & 6 \\ 0 & -1 & -3 & : & -11 \\ 0 & 0 & 1 & : & 3 \end{pmatrix}$$

4. We have the system

$$\begin{cases} x + y + z = 6 \\ -y - 3z = -11 \\ z = 3 \end{cases}$$

Then,

$$-y - 9 = -11 \Rightarrow y = 2$$

$$x + 2 + 3 = 6 \Rightarrow x = 1$$

$$S = \{(1, 2, 3)\}$$

Synthesis

For the system

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = c_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = c_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = c_m \end{cases}$$

The Gauss elimination method is used to transform a system of equations into an equivalent system that is in row echelon form.

To facilitate the calculation, transform the system into a matrix and place the coefficients of the variables and the independent terms into the matrix as follows:

$$(A:C) = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} & :c_1 \\ a_{21} & a_{22} & \dots & a_{2n} & :c_2 \\ \vdots & \vdots & \dots & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & :c_m \end{pmatrix}$$

And then, transform the system in the form where the elements above and below the leading diagonal of matrix A become zeros. The system is now reduced to the simplest system.

Exercise 7.8 Learner's Book page 376

1. $S = \{(-4, -5, 2)\}$
2. No solution
3. Infinity number of solution.
4. Infinity number of solution.

Lesson 7.9. Power of matrix

Learning objectives

Given a square matrix and by using diagonalisation method, learners should be able to find the power of that matrix accurately.

Prerequisites

- ⑦ Finding eigenvalues and eigenvectors of a matrix.

Teaching Aids

Exercise book and pen



Activity 7.9 Learner's Book page 377

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

1. $A^2 = AA = PDP^{-1}PDP^{-1} = PD^2P^{-1}$
2. $A^3 = A^2A = PD^2P^{-1}PDP^{-1} = PD^3P^{-1}$
3. $A^4 = A^3A = PD^3P^{-1}PDP^{-1} = PD^4P^{-1}$
4. $A^5 = A^4A = PD^4P^{-1}PDP^{-1} = PD^4DP^{-1} = PD^5P^{-1}$
 $A^n = A^{n-1}A = PD^{n-1}P^{-1}PDP^{-1} = PD^nP^{-1} \Rightarrow A^n = PD^nP^{-1}$

Synthesis

The power of matrix A is given by $A^n = PD^nP^{-1}$ for an invertible matrix P whose columns are elements of eigenvectors of matrix A , and D is diagonal matrix of A .

Where,

$$D^n = \begin{pmatrix} \lambda_1^n & 0 & \cdots & 0 \\ 0 & \lambda_2^n & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_k^n \end{pmatrix}$$

λ_k are eigenvalues

Exercise 7.9 Learner's Book page 378

1. $P = \begin{pmatrix} -3 & 4 \\ 4 & 3 \end{pmatrix}$, $D = \begin{pmatrix} 20 & 0 \\ 0 & -5 \end{pmatrix}$, $A^3 = \begin{pmatrix} 2800 & -3900 \\ -3900 & 5075 \end{pmatrix}$
2. $P = \begin{pmatrix} 1 & 2 \\ 1 & 1 \end{pmatrix}$, $D = \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix}$, $A^5 = \begin{pmatrix} -245 & 488 \\ -244 & 487 \end{pmatrix}$

$$3. \quad P = \begin{pmatrix} -1 & 1 \\ 1 & 0 \end{pmatrix}, \quad D = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad A^{20} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

$$4. \quad P = \begin{pmatrix} 1 & 3 & 3 \\ 0 & 1 & 0 \\ 0 & 1 & 1 \end{pmatrix}, \quad D = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad A^5 = \begin{pmatrix} 32 & 0 & 633 \\ 0 & 243 & 0 \\ 0 & 0 & 243 \end{pmatrix}$$

Summary of the unit

1. Kernel and range

- ⑦ The kernel of a linear mapping $f: E \rightarrow F$ denoted $Ker(f)$ is a subset of E whose image f is 0-vector of F . i.e, $Ker(f) = \{v \in E: f(v) = 0\}$.

A linear transformation f is called singular if there exists a non-zero vector whose image is zero vector. Thus, it is non-singular if the only zero vector has zero vector as image, or equivalently, if its kernel consists only of the zero vector: $Ker(f) = \{0\}$.

A linear transformation $f: E \rightarrow F$ is one-to-one (1-1) if and only if $Ker(f) = \{0\}$.

- ⑦ The nullity of f denoted $n(f)$ is the dimension of $Ker(f)$. i.e, $n(f) = \dim Ker(f)$.
- ⑦ The image or range of a linear mapping $f: E \rightarrow F$ is the set of points in F to which points in E are mapped on. i.e, $Im f = \{u \in F: f(v) = u\}, v \in E$.

A linear transformation $f: E \rightarrow F$ is onto if the range is equal to F .

- ⑦ The rank of f denoted $rank(f)$ or $r(f)$ is the dimension of image of f . i.e, $rank(f) = \dim(Im f)$.

2. Elementary row/column operations

When these operations are performed on rows, they are called elementary row operations; and when they are performed on columns, they are called elementary column operations.

Operation description	Notation
<u>Row operations</u>	
1. Interchange row i and j	$\rightarrow r_i \leftrightarrow r_j$
2. Multiply row i by $s \neq 0$	$\rightarrow \text{new } r_i \rightarrow sr_i$
3. Add s times row i to row j	$\rightarrow \text{new } r_j \rightarrow r_j + sr_i$
<u>Column operations</u>	
1. Interchange column i and j	$\rightarrow c_i \leftrightarrow c_j$
2. Multiply column i by $s \neq 0$	$\rightarrow \text{new } c_i \rightarrow sc_i$
3. Add s times column i to column j	$\rightarrow \text{new } c_j \rightarrow c_j + sc_i$

Two matrices are said to be row equivalent (or column equivalent) if one can be changed to the other by a sequence of elementary row (or column) operations.

Two matrices A and B are said to be similar if $B = P^{-1}AP$ for some invertible matrix P .

3. Diagonalisation of matrices

a) Eigenvalues and eigenvectors

The eigenvalues of f are the roots (in K) of the polynomial; $\det(f - \lambda I)$. This polynomial is a polynomial associated with f and is called characteristic polynomial. For any square matrix A , the polynomial $\det(A - \lambda I)$ is its characteristic polynomial.

The homogeneous system $(f - \lambda I)\vec{u} = \vec{0}$ gives the eigenvector \vec{u} associated with eigenvalue λ .

b) Diagonalisation

To diagonalise matrix A , we perform the following steps:

1. Find the eigenvalues.
2. If there is a non-real eigenvalue, the matrix cannot be diagonalised.
3. If all eigenvalues are real, find their associated eigenvectors (they must be linearly independent).
4. If the number of eigenvectors is not equal to the order of matrix A , then this matrix cannot be diagonalised.
5. If the number of eigenvectors is equal to the order of matrix A , form matrix P whose columns are elements of eigenvectors.
6. Find the inverse of P .
7. Find D , diagonal matrix of A by relation $D = P^{-1}AP$.

4. Applications**a) Echelon matrix**

A matrix is in **row echelon form** (ref) when it satisfies the following conditions:

- ▶ The first non-zero element in each row, called the leading entry, is 1.
- ▶ Each leading entry is in a column to the right of the leading entry in the previous row.
- ▶ Rows with all zero elements, if any, are below rows having a non-zero element.

A matrix is in **reduced row echelon form** (rref) when it satisfies the following conditions:

- ▶ The matrix is in row echelon form (i.e., it satisfies the three conditions listed above).
- ▶ The leading entry in each row is the only non-zero entry in its column.

b) Matrix inverse

A is a square matrix of order n . To calculate the inverse of A , denoted as A^{-1} , follow these steps:

- ▶ Construct a matrix of type $M = (A|I)$, that is to say, A is in the left half of M and the identity matrix I is on the right.
- ▶ Using the Gaussian elimination method, transform the left half, A , to the identity matrix located to the right, and the matrix that results in the right side will be the inverse of matrix A .

c) Rank of matrix

The rank of matrix is the number of linearly independent rows or columns. Using this definition, the Gaussian elimination method is used to find the rank.

A line can be discarded if:

- ▶ All the coefficients are zeros.
- ▶ There are two equal lines.
- ▶ A line is proportional to another.
- ▶ A line is a linear combination of others.

In general, eliminate the maximum possible number of lines, and the rank is the number of non-zero rows.

d) Solving system of linear equations

Consider the following system;

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = c_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = c_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = c_m \end{cases}$$

The Gauss elimination method is to transform a system of equations into an equivalent system that is in triangular form.

To facilitate the calculation, transform the system into a matrix and place the coefficients of the variables and the independent terms into the matrix as follows:

$$(A:C) = \begin{pmatrix} a_{11} & a_{12} \cdots a_{1n} & :c_1 \\ a_{21} & a_{22} \cdots a_{2n} & :c_m \\ \vdots & \vdots \cdots \vdots & \vdots \\ a_{m1} & a_{m2} \cdots a_{mn} & :c_m \end{pmatrix}$$

Where

The matrix $(A:C)$ is called **augmented matrix**.

Remarks

- ⑦ If $\text{rank}(A) \neq \text{rank}(A:C)$, the system is said to be inconsistent and there is no solution.
- ⑦ If $\text{rank}(A) = \text{rank}(A:C) = r$, the system is said to be consistent and there is solution.
 - » If $r = n$, as there are n unknowns, then the system has a unique solution.
 - » If $r < n$, the system has infinite solutions. (It is undetermined system).

5. Power of matrix

The power of matrix A is given by $A^n = PD^nP^{-1}$ for an invertible matrix P whose columns are elements of eigenvectors of matrix A and D is diagonal matrix of A .

Where,

$$D^n = \begin{pmatrix} \lambda_1^n & 0 & \cdots & 0 \\ 0 & \lambda_2^n & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_k^n \end{pmatrix}$$

λ_k are eigenvalues

End of Unit Assessment answers Learner's Book page 383

1. a) $\lambda^2 - 5\lambda + 1$ b) $\lambda^2 - 3\lambda - 18$
 c) $\lambda^2 + 9$ d) $\lambda^3 + \lambda^2 - 8\lambda + 62$
 e) $\lambda^3 - 6\lambda^2 - 35\lambda - 38$
2. a) Eigenvalues: 2 and -5 , eigenvectors: $\vec{u} = \begin{pmatrix} 4 \\ 1 \end{pmatrix}$ and $\vec{v} = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$
 b) $P = \begin{pmatrix} 4 & 1 \\ 1 & 2 \end{pmatrix}$, $D = \begin{pmatrix} 2 & 0 \\ 0 & -5 \end{pmatrix}$
3. a) Eigenvalues: 1 and 4, eigenvectors: $\vec{u} = \begin{pmatrix} 2 \\ -1 \end{pmatrix}$ and $\vec{v} = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$
 b) $P = \begin{pmatrix} 2 & 1 \\ -1 & 1 \end{pmatrix}$, $D = \begin{pmatrix} 1 & 0 \\ 0 & 4 \end{pmatrix}$
 c) $A^6 = \begin{pmatrix} 1366 & 2230 \\ 1365 & 2731 \end{pmatrix}$, $f(A) = \begin{pmatrix} 1 & 2 \\ -1 & 0 \end{pmatrix}$
 d) $\frac{1}{3} \begin{pmatrix} 2 + \sqrt[3]{4} & -2 + 2\sqrt[3]{4} \\ -1 + \sqrt[3]{4} & 1 + 2\sqrt[3]{4} \end{pmatrix}$
4. a) Eigenvalues: 3 and 5
 b) $S = \{(1, -1, 0), (1, 0, 1), (1, 2, 1)\}$
 c) $\begin{pmatrix} 3 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 5 \end{pmatrix}$
5. $A = \begin{pmatrix} 2 & 1 & -2 \\ 2 & 3 & -4 \\ 1 & 1 & -1 \end{pmatrix}$, $D = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}$
6. To be proved

$$7. \text{ a) } \begin{pmatrix} 1 & 0 & -1 & -2 \\ 0 & 1 & 2 & 3 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\text{b) } \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\text{c) } \begin{pmatrix} 1 & 0 & 1 & 0 & 1 \\ 0 & 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 \end{pmatrix}$$

$$\text{d) } \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & 2 \end{pmatrix}$$

8. a) Rank 2

b) Rank 2

c) Rank 3

d) Rank 2

9. a) $S = \{(-2, 5, 10)\}$

b) No solution

$$10. \text{ a) } \frac{1}{5} \begin{pmatrix} -2 & -3 & 5 \\ 1 & 4 & -5 \\ -6 & -29 & 40 \end{pmatrix}$$

$$\text{b) } \frac{1}{11} \begin{pmatrix} 2 & 5 & -14 \\ 1 & -3 & 4 \\ -1 & 3 & 7 \end{pmatrix}$$

$$\text{c) } \frac{1}{2} \begin{pmatrix} 2 & 1 & 0 \\ 2 & 0 & 0 \\ -4 & -1 & 2 \end{pmatrix}$$

11. Characteristic equation: $\lambda^3 - 5\lambda^2 + 7\lambda - 3 = 0$

From Cayley-Hamilton theorem, we have

$$A^3 - 5A^2 + 7A - 3I = 0.$$


$$\text{Now, } A^8 - 5A^7 + 7A^6 - 3A^5 + A^4 - 5A^3 + 8A^2 - 2A + I$$

$$= A^5(A^3 - 5A^2 + 7A - 3I) + A(A^3 - 5A^2 + 7A - 3A) + A^2 + A + I$$

$$= 0 + 0 + A^2 + A + I$$

$$= \begin{pmatrix} 5 & 4 & 4 \\ 0 & 1 & 0 \\ 4 & 4 & 5 \end{pmatrix} + \begin{pmatrix} 2 & 1 & 1 \\ 0 & 1 & 0 \\ 1 & 1 & 2 \end{pmatrix} + \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 8 & 5 & 5 \\ 0 & 3 & 0 \\ 5 & 5 & 8 \end{pmatrix}$$





Conics

Learner's Book pages 387 - 444

Key unit competence

- ④ Determine the characteristics and the graph of a conic given by its Cartesian, parametric or polar equation.
 - ④ Find the Cartesian, parametric and polar equations of a conic from its characteristics.
-

Vocabulary or key words concepts

Conic section: Curve obtained by intersecting a double cone with a plane.

Parabola: Conic section obtained when the plane is parallel to generator but not along the generator.

Ellipse: Conic section obtained when the plane cuts the cone obliquely.

Hyperbola: Conic section obtained when the plane is parallel to the axis but not along the axis.

Guidance on the problem statement

The problem statement:

“How can one find the equation representing a curve which generates a parabolic antenna? What can you say about motion of planets around the sun? How can one find the equation of their orbits around the sun?”

These questions lead us to the applications of conics.

List of lessons

No	Lesson title	Number of periods
1	Generalities on conic sections	3
2	Definition and equation of a parabola	3
3	Tangent line and normal line on a parabola	3
4	Definition and equation of an ellipse	3
5	Tangent line and normal line on an ellipse	3
6	Definition and equation of a hyperbola	3
7	Tangent line and normal line on a hyperbola	3
8	Definition of polar coordinates	3
9	Polar equation of a conic	3
10	Polar equation of a straight line	3
11	Polar equation of a circle	3
12	Applications of conics	2
Total periods		35

Lesson development

Lesson 8.1. Generalities on conic sections

Learning objectives

Given a double cone and a plane, learners should be able to define a conic and draw the shape of conic sections accurately.

Prerequisites

- ✔ Double cone
- ✔ Plane

Teaching Aids

Exercise book, pen, calculator and instruments of geometry




Activity 8.1 Learner's Book page 388

In this lesson, the following generic competence and cross-cutting issues are to be addressed:


- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers


1. The plane is parallel to a generator of the cone but not along the generator




2. The plane cuts the cone obliquely



3. The plane is parallel to the axis but not along the axis



4. The plane is parallel to the base but does not pass through the vertex



Synthesis

Conic is the name given to the shapes that we obtain by taking different plane slices through a double cone. The sections of a right circular cone by different planes give curves of different shapes: parabola, ellipse, hyperbola, circle, single point, single line, pair of lines.

A conic section is the set of all points which move in a plane such that its distance from a fixed point and a fixed line not containing the fixed point are in a constant ratio.




Figure 8.8: Conic section

From figure 8.8, we have

$$|PF| = e|PM| \text{ where } M \text{ is a foot of perpendicularity of}$$

line joining P to directrix, P point lying on conic and F focal point.

Focal axis is a line passing through the focus and perpendicular to the directrix.

Vertex is a point where the conic intersects its axis.

Exercise 8.1 Learner's Book page 391

1. Single point: This is formed when the plane passes through the vertex horizontally, i.e. parallel to the base.
2. Single line: This is formed when the plane passes through the vertex and along the generator.
3. Pair of lines: This is the section formed when the plane passes through the vertex. In this case, the section is a pair of straight lines passing through the vertex.

Lesson 8.2. Definition and equation of a parabola

Learning objectives

Through examples, learners should be able to define a parabola and determine its equation accurately.

Prerequisites

- ✔ Distance between two points.
- ✔ Distance from a point to a straight line.
- ✔ Curve sketching.

Teaching Aids

Exercise book, pen, calculator and instruments of geometry



Activity 8.2 Learner's Book page 392

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Choose some point having coordinates (x, y) . The distance between this point and $(5, 3)$ is given by

$$\sqrt{(x-5)^2 + (y-3)^2}.$$

The distance between point (x, y) and $(2, 1)$ is given

$$\text{by } \sqrt{(x-2)^2 + (y-1)^2}.$$

Equating these distances, as the point (x, y) is to be equidistant from the two given points, we have

$$\sqrt{(x-5)^2 + (y-3)^2} = \sqrt{(x-2)^2 + (y-1)^2}$$

Squaring both sides, we get

$$(x-5)^2 + (y-3)^2 = (x-2)^2 + (y-1)^2$$

Expanding, we have

$$x^2 - 10x + 25 + y^2 - 6y + 9 = x^2 - 4x + 4 + y^2 - 2y + 1$$

Cancelling and combining like terms, we get

$$4y + 5 = -6x + 34$$

Or

$$4y = -6x + 29$$

This is the equation of a straight line with slope $-\frac{3}{2}$

and y intercept $\frac{29}{4}$.

2. Choose some point on the curve having coordinates (x, y) .

The distance from the point (x, y) on the curve to the line $x = -3$ is $\sqrt{(x+3)^2 + (y-y)^2} = \sqrt{(x+3)^2}$.

The distance from the point (x, y) on the curve to the point $(3, 0)$ is $\sqrt{(x-3)^2 + (y-0)^2} = \sqrt{(x-3)^2 + y^2}$.

Equating the two distances yields


$$\sqrt{(x+3)^2} = \sqrt{(x-3)^2 + y^2}$$

Squaring and expanding both sides, we get

$$x^2 + 6x + 9 = x^2 - 6x + 9 + y^2$$


Cancelling and collecting like terms yields $y^2 = 12x$ which is an equation of a curve.

3. Curve



Synthesis

A **parabola** is set of points $P(x, y)$ in the plane equidistant from a fixed point F , called focus and a fixed line d , called **directrix**. In the figure below $\overline{PF} = \overline{PM}$, where $M \in d$.




The equation of a parabola, whose focus at point $(a,0)$ and directrix with equation $x = -a$, is given by $y^2 = 4ax$.

The standard forms of the equation of parabola with vertices at the point $V(h,k)$ are as follows:


1. $(y - k)^2 = 4a(x - h)$, parabola opens to the right.
2. $(y - k)^2 = -4a(x - h)$, parabola opens to the left.
3. $(x - h)^2 = 4a(y - k)$, parabola opens upward.
4. $(x - h)^2 = -4a(y - k)$, parabola opens downward.

Exercise 8.2 Learner's Book page 398


1. Focus is $(-2,0)$, directrix is $x = 2$
2. a) Sketch:




b) Sketch:



c) Sketch:



d) Sketch:



3. a) Focus $\left(\frac{25}{4}, 0\right)$; directrix $x = -\frac{25}{4}$; length of latus rectum 25; equation of latus rectum $x = \frac{25}{4}$; ends of latus rectum $\left(\frac{25}{4}, -\frac{25}{2}\right)$ and $\left(\frac{25}{4}, \frac{25}{2}\right)$.
- b) Focus $(0, 2)$; directrix $y = -2$; length of latus rectum 8; equation of latus rectum $y = 2$; ends of latus rectum $(4, 2)$ and $(-4, 2)$.
- c) Focus $\left(0, -\frac{5}{4}\right)$; directrix $y = \frac{5}{4}$; length of latus rectum 5; equation of latus rectum $y = -\frac{5}{4}$; ends of latus rectum $\left(\frac{5}{2}, -\frac{5}{4}\right)$ and $\left(-\frac{5}{2}, -\frac{5}{4}\right)$
4. $(y-2)^2 = 12(x-1)$
5. a) $x^2 = -8y$ b) $y^2 = 4(x+4)$
- c) $4x^2 + 4xy + y^2 + 4x + 32y + 16 = 0$
6. a) $\frac{7}{4}$ b) $\left(\frac{41}{16}, 1\right); (3, 1)$
7. $(1, 3)$

Lesson 8.3. Tangent line and normal line on a parabola

Learning objectives

Given equation of parabola, learners should be able to find equation of tangent line and normal line at a given point and draw them accurately.

Prerequisites

- ✔ Equation of tangent at a point on a curve.
- ✔ Equation of normal line at a point on a curve.
- ✔ Differentiation.

Teaching Aids

Exercise book, pen and calculator



Activity 8.3 Learner's Book page 399

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. $T \equiv y - y_o = m(x - x_o)$

Differentiating with respect to x yields

$$2y \frac{dy}{dx} = 4a \Rightarrow \frac{dy}{dx} = \frac{2a}{y} \text{ and } m = \frac{2a}{y_o}$$

Then,

$$T \equiv y - y_o = \frac{2a}{y_o}(x - x_o)$$

$$\Leftrightarrow y_o y - y_o y_o = 2a(x - x_o) \Leftrightarrow y_o y - 4ax_o = 2ax - 2ax_o$$

$$\Leftrightarrow y_o y - 4ax_o = 2ax - 2ax_o \text{ since } y_o y_o = y_o^2 = 4ax_o$$

$$\Leftrightarrow y_o y = 2ax - 2ax_o + 4ax \Leftrightarrow y_o y = 2ax + 2ax_o$$

$$\Leftrightarrow y_o y = 2a(x + x_o)$$

Therefore, $T \equiv y_o y = 2a(x + x_o)$

2. Equation of normal line:

$$N \equiv y - y_o = -\frac{1}{m}(x - x_o), \text{ with } m = \left. \frac{dy}{dx} \right|_{x=x_o}$$

Then,

$$N \equiv y - y_o = -\frac{y_o}{2a}(x - x_o) \Leftrightarrow 2ay - 2ay_o = -y_o x + y_o x_o$$

$$\Rightarrow 2ay_o y - 2ay_o y_o = -y_o y_o x + y_o y_o x_o$$

$$\Rightarrow 2ay_o y - 2ay_o^2 = -y_o^2 x + y_o^2 x_o$$

$$\Rightarrow 2ay_0y - 8a^2x_0 = -4ax_0x + 4ax_0^2$$

$$\Leftrightarrow y_0y - 4ax_0 = -2x_0x + 2x_0^2$$

$$\Leftrightarrow y_0y - y_0y_0 = -2x_0(x - x_0)$$


Therefore, $N \equiv y_0y - y_0y_0 = -2x_0(x - x_0)$.

3. The tangent line of $y^2 = 2x$ at $(0,0)$:

Since the tangent line at point (x_0, y_0) , on parabola $y^2 = 4ax$, is given by

$$T \equiv y_0y = 2a(x + x_0), \text{ here, } y^2 = 4\left(\frac{1}{2}\right)x. \text{ So } a = \frac{1}{2}.$$

Then the tangent line is $T \equiv 0y = 2\left(\frac{1}{2}\right)(x + 0)$ or tangent line has equation $x = 0$.



The line $x = 0$ touches the parabola $y^2 = 2x$ once at $(0,0)$, as it is its tangent at $(0,0)$.

Synthesis

The tangent line at point (x_0, y_0) , on parabola $y^2 = 4ax$, is given by $T \equiv y_0y = 2a(x + x_0)$, and the normal line at the same point is $N \equiv y_0y - y_0y_0 = -2x_0(x - x_0)$.

Exercise 8.3 Learner's Book page 401

1. -6
2. Focus is $F(-2, 4)$, vertex is $V(-2, -3)$, equation of axis is $x + 2 = 0$, equation of directrix is $y + 2 = 0$, equation of tangent at vertex is $y + 3 = 0$.
3. $2y = x + 8$, $y + 2x + 1 = 0$
4. $y + tx = 2at + at^3; 0, \pm 2$

Lesson 8.4. Definition and equation of an ellipse**Learning objectives**

Through examples, learners should be able to define an ellipse and determine its equation accurately.

Prerequisites

- ✔ Distance between two points.
- ✔ Distance from a point to a straight line.
- ✔ Curve sketching.

Teaching Aids

Exercise book, pen, calculator and instruments of geometry

Activity 8.4 Learner's Book page 401

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Choose some point on the curve having coordinates (x, y) .

The distance from the point (x, y) on the curve to the point $(3, 0)$ is $d_1 = \sqrt{(3-x)^2 + (0-y)^2} = \sqrt{(3-x)^2 + y^2}$.

The distance from the point (x, y) to the line $x = \frac{25}{3}$ is $d_2 = \sqrt{\left(\frac{25}{3} - x\right)^2 + (y-y)^2} = \frac{25}{3} - x$.

Since $\frac{d_1}{d_2} = \frac{3}{5} \Rightarrow d_1 = \frac{3}{5}d_2$, then,

$$\sqrt{(3-x)^2 + y^2} = \frac{3}{5}\left(\frac{25}{3} - x\right)$$

Squaring both sides and expanding, we get

$$9 - 6x + x^2 + y^2 = 25 - 6x + \frac{9}{25}x^2$$


Collecting like terms and transposing give

$$\frac{16}{25}x^2 + y^2 = 16$$

Dividing each term by 16, we see that $\frac{x^2}{25} + \frac{y^2}{16} = 1$.


This equation is of an **ellipse**.

2. Sketch of the curve:



Synthesis

We define an ellipse with eccentricity e (where $0 < e < 1$) to be the set of points P in the plane whose distance from a fixed point F is e times their distance from a fixed line.



For any point $P(x, y)$ on the ellipse, we have $\overline{PF} = e\overline{PM}$, $M \in D$, where D is the directrix with equation $x = \frac{a}{e}$.

$$\overline{PF} = d_1 = \sqrt{(ae - x)^2 + (0 - y)^2} = \sqrt{(ae - x)^2 + y^2}.$$

$$\overline{PM} = d_2 = \sqrt{(ae - x)^2 + (y - y)^2} = \frac{a}{e} - x.$$

Since $\overline{PF} = e\overline{PM}$, then,

$$\sqrt{(ae - x)^2 + y^2} = a - ex$$

Squaring both sides and expanding, we get

$$a^2e^2 - 2aex + x^2 + y^2 = a^2 - 2aex + e^2x^2$$

Collecting terms

$$(1 - e^2)x^2 + y^2 = a^2(1 - e^2)$$

Dividing each term by $a^2(1 - e^2)$, we get

$$\frac{x^2}{a^2} + \frac{y^2}{a^2(1 - e^2)} = 1$$

Writing $b^2 = a^2(1 - e^2)$, this gives

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

This is equation ellipse centred at $(0,0)$ in standard form.


For the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, $b^2 = a^2(1 - e^2)$ with $e < 1$, we have two foci at $(\pm ae, 0)$ and two directrices $x = \pm \frac{a}{e}$.

When the centre of ellipse is located at some point other than $(0,0)$, say the point (x_0, y_0) , the equation of


ellipse in standard form is $\frac{(x - x_0)^2}{a^2} + \frac{(y - y_0)^2}{b^2} = 1$.

Exercise 8.4 Learner's Book page 407


1. $(0, \sqrt{2})$ and $(0, -\sqrt{2})$
2. $3x^2 + 7y^2 = 115$
3. $\frac{x^2}{12} + \frac{y^2}{16} = 1$
4. a) Sketch



b) Curve



c) Curve



5. $16x^2 + 9y^2 - 64x - 54y + 1 = 0$

$$\Rightarrow \frac{(x-2)^2}{9} + \frac{(y-3)^2}{16} = 1$$

Foci are $(2, 3 + \sqrt{7})$ and $(2, 3 - \sqrt{7})$

6. a) $e = \frac{\sqrt{2}}{2}$ b) $e = \frac{\sqrt{3}}{3}$ c) $e = \frac{\sqrt{3}}{2}$

7. 7 or 13

8. 13:5

Lesson 8.5. Tangent line and normal line on ellipse

Learning objectives

Given equation of ellipse, learners should be able to find tangent line and normal line at a given point and draw them accurately.

Prerequisites

- ✔ Equation of tangent line.
- ✔ Equation of normal line.
- ✔ Differentiation.

Teaching Aids

Exercise book, pen, calculator and instruments of geometry



Activity 8.5 Learner's Book page 408

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Equation of tangent line $T \equiv y - y_o = m(x - x_o)$ where

$$m = \left. \frac{dy}{dx} \right|_{x=x_o}$$

Differentiating with respect to x gives

$$\frac{2}{a^2}x + \frac{2}{b^2}y \frac{dy}{dx} = 0 \quad \text{or} \quad \frac{dy}{dx} = -\frac{b^2}{a^2} \frac{x}{y}$$

$$\text{Then, } m = -\frac{b^2}{a^2} \frac{x_o}{y_o}$$

$$\text{And } T \equiv y - y_o = -\frac{b^2}{a^2} \frac{x_o}{y_o} (x - x_o)$$

$$\Leftrightarrow a^2 y_0 y - a^2 y_0 y_0 = -b^2 x_0 (x - x_0)$$

$$\Leftrightarrow a^2 y_0 y + b^2 x_0 x = b^2 x_0 x_0 + a^2 y_0 y_0,$$

$$\text{since } b^2 x_0 x_0 + a^2 y_0 y_0 = a^2 b^2$$

$$\text{Thus } a^2 y_0 y + b^2 x_0 x = a^2 b^2$$


Dividing each term by $a^2 b^2$, we get

$$\frac{y_0 y}{b^2} + \frac{x_0 x}{a^2} = 1 \text{ or } \frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} = 1$$

$$\text{Therefore, } T \equiv \frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} = 1$$

2. The tangent line to the curve $x^2 + \frac{y^2}{9} = 1$ at $(0, 3)$ is given by $T \equiv \frac{(0)x}{1} + \frac{3y}{9} = 1$ or $T \equiv y = 3$.

Curve:



Synthesis

The tangent line at point (x_0, y_0) , on ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, is given by: $T \equiv \frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} = 1$

Exercise 8.5 Learner's Book page 411

1. $\pm \frac{3}{2}$
2. $y = -1.155x + 4$
3. $k = -4$ at $(-2, -1)$; $k = 4$ at $(2, 1)$
4. At $(0, 0)$: $T \equiv y = -2x$, at $(0, 2)$: $T \equiv y = 2x + 2$,
5. a) $\frac{6 - \sqrt{6}}{4} < m < \frac{6 + \sqrt{6}}{4}$ b) $m = \frac{6 \pm \sqrt{6}}{4}$
 c) $m < \frac{6 - \sqrt{6}}{4}$ or $m > \frac{6 + \sqrt{6}}{4}$

Lesson 8.6. Definition and equation of a hyperbola**Learning objectives**

Through examples, learners should be able to define hyperbola and find its equation accurately.

Prerequisites

- ✔ Distance between two points.
- ✔ Distance from a point to a straight line.
- ✔ Differentiation.

Teaching Aids

Exercise book, pen, calculator and instruments of geometry

**Activity 8.6 Learner's Book page 411**

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

If the difference of the distances from any point $P(x, y)$ on conic to the two foci is $2a$, thus

$$\begin{aligned} PF_1 - PF_2 = 2a &\Leftrightarrow \sqrt{(c-x)^2 + (0-y)^2} - \sqrt{(-c-x)^2 + (0-y)^2} = 2a \\ &\Leftrightarrow \sqrt{(c-x)^2 + y^2} - \sqrt{(c+x)^2 + y^2} = 2a \end{aligned}$$

Transposing one term from the left side to the right side and squaring, we get

$$\Leftrightarrow (c-x)^2 + y^2 = 4a^2 + 4a\sqrt{(c+x)^2 + y^2} + (c+x)^2 + y^2$$

$$\Leftrightarrow c^2 - 2cx + x^2 = 4a^2 + 4a\sqrt{(c+x)^2 + y^2} + c^2 + 2cx + x^2$$

$$\Leftrightarrow -2cx = 4a^2 + 4a\sqrt{(c+x)^2 + y^2} + 2cx$$

$$\Leftrightarrow -4(cx + a^2) = 4a\sqrt{(c+x)^2 + y^2} \Leftrightarrow -(cx + a^2) = a\sqrt{(c+x)^2 + y^2}$$

Squaring again both sides and expanding, we have

$$\Leftrightarrow c^2x^2 + 2cxa^2 + a^4 = a^2(c^2 + 2cx + x^2 + y^2)$$

$$\Leftrightarrow c^2x^2 + 2cxa^2 + a^4 = a^2c^2 + 2cxa^2 + a^2x^2 + a^2y^2$$

$$\Leftrightarrow c^2x^2 - a^2x^2 - a^2y^2 = a^2c^2 - a^4 \Leftrightarrow x^2(c^2 - a^2) - a^2y^2 = a^2(c^2 - a^2)$$

Since $c = \sqrt{a^2 + b^2}$ thus $c^2 - a^2 = b^2$ and

$$x^2(c^2 - a^2) - a^2y^2 = a^2(c^2 - a^2) \Leftrightarrow x^2b^2 - a^2y^2 = a^2b^2$$


Dividing both sides by a^2b^2 , we get

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

This is equation of hyperbola.

Synthesis

We define a hyperbola to be the set of all points P in the plane, the difference of whose distances from two fixed points, called **foci**, is a constant equal to $2a$.




In standard form a hyperbola is given by the equation $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$; this hyperbola has two foci $(\pm c, 0)$, $c^2 = a^2 + b^2$ and two directrices $x = \pm \frac{a^2}{a}$.

Eccentricity of the hyperbola is $e = \frac{c}{a} = \frac{\sqrt{a^2 + b^2}}{a}$, $e > 1$.

If the hyperbola has centre at (h, k) , then the equation is $\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$.

Exercise 8.6 Learner's Book page 415

1. a) Curve




Vertices: $(0, \pm 2)$

Eccentricity is $e = \frac{\sqrt{13}}{2}$

Foci: $(0, \sqrt{13})$ and $(0, -\sqrt{13})$

Asymptotes: $y = \frac{2}{3}x$ and $y = -\frac{2}{3}x$.

b) Curve




Vertices: $(5, 4)$ and $(-1, 4)$

Eccentricity is $e = \frac{\sqrt{13}}{3}$

Foci: $(2 + \sqrt{13}, 4)$ and $(2 - \sqrt{13}, 4)$

Asymptotes: $y = \frac{2}{3}x + \frac{8}{3}$ and $y = -\frac{2}{3}x + \frac{16}{3}$

c) Curve



Vertices: $(-2, 3)$ and $(-2, -9)$

Eccentricity is $e = \frac{\sqrt{10}}{3}$

Foci: $(-2, -3 + 2\sqrt{10})$ and $(-2, -3 - 2\sqrt{10})$

Asymptotes: $y = 3x + 3$ and $y = -3x - 9$

2. Foci are $(0, -5)$ and $(0, 5)$, vertices are $(0, -4)$ and

$(0, 4)$, asymptotes: $y = \pm \frac{4}{3}x$

3. a) Length of transverse axis is $2a = 6$; conjugate axis is $2b = 8$.

Eccentricity $\frac{\sqrt{a^2 + b^2}}{a} = \frac{5}{3}$, coordinates of foci

$(\pm ae, 0) = (\pm 5, 0)$, coordinates of vertices

$V(\pm a, 0) = (\pm 3, 0)$.

b) Length of transverse axis is $2a = 2\sqrt{3}$; conjugate axis is $2b = 2\sqrt{2}$.

Eccentricity $\frac{\sqrt{a^2 + b^2}}{a} = \frac{\sqrt{15}}{3}$, coordinates of

foci $(\pm ae, 0) = (\pm\sqrt{5}, 0)$, coordinates of vertices

$V(\pm a, 0) = (\pm\sqrt{3}, 0)$.

c) Length of transverse axis is $2b = 8$; conjugate axis is $2a = 2$.

Eccentricity $\frac{\sqrt{a^2 + b^2}}{a} = \frac{\sqrt{17}}{4}$, coordinates of foci

$(0, \pm be) = (0, \pm\sqrt{17})$, coordinates of vertices

$V(0, \pm b) = (0, \pm 4)$.

4. $\frac{y^2}{64} - \frac{x^2}{36} = 1$

$$5. \quad x^2 - y^2 - 4x + 8y - 21 = 0$$

$$\Rightarrow \frac{(x-2)^2}{9} - \frac{(y-4)^2}{9} = 1$$

Vertices: $(-1, 4), (5, 4)$

Foci: $(2 - 3\sqrt{2}, 4), (2 + 3\sqrt{2}, 4)$

Asymptotes: $y - 4 = \pm(x - 2)$

$$6. \quad \text{a) } 7x^2 + 24xy - 56x - 6y + 68 = 0$$

$$\text{b) } 9x^2 - 16y^2 - 36x + 96y - 252 = 0$$

$$7. \quad 21x^2 - 4y^2 - 84 = 0$$

Lesson 8.7. Tangent line and normal line on hyperbola

Learning objectives

Given equation of hyperbola, learners should be able to find equation of tangent line and normal line at a given point and draw them accurately.

Prerequisites

- ✔ Equation of tangent line.
- ✔ Equation of normal line.
- ✔ Differentiation.

Teaching Aids

Exercise book, pen, calculator and instruments of geometry

Activity 8.7 Learner's Book page 416



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

Equation of tangent line $T \equiv y - y_0 = m(x - x_0)$ where

$$m = \left. \frac{dy}{dx} \right|_{x=x_0}$$

Differentiating with respect to x gives

$$\frac{2}{a^2}x - \frac{2}{b^2}y \frac{dy}{dx} = 0 \quad \text{or} \quad \frac{dy}{dx} = \frac{b^2}{a^2} \frac{x}{y}$$

$$\text{Then, } m = \frac{b^2}{a^2} \frac{x_0}{y_0}$$

$$\text{And } T \equiv y - y_0 = \frac{b^2}{a^2} \frac{x_0}{y_0} (x - x_0)$$

$$\Leftrightarrow a^2 y_0 y - a^2 y_0 y_0 = b^2 x_0 (x - x_0)$$

$$\Leftrightarrow a^2 y_0 y - b^2 x_0 x = -b^2 x_0 x_0 + a^2 y_0 y_0, \text{ as}$$

$$b^2 x_0 x_0 - a^2 y_0 y_0 = a^2 b^2$$

$$a^2 y_0 y - b^2 x_0 x = -a^2 b^2$$

Dividing each term by $-a^2 b^2$, we get

$$-\frac{y_0 y}{b^2} + \frac{x_0 x}{a^2} = 1 \quad \text{or} \quad \frac{x_0 x}{a^2} - \frac{y_0 y}{b^2} = 1$$

$$\text{Therefore, } T \equiv \frac{x_0 x}{a^2} - \frac{y_0 y}{b^2} = 1$$

Synthesis

The tangent line at point (x_0, y_0) , on hyperbola

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1, \text{ is given by } T \equiv \frac{x_0 x}{a^2} - \frac{y_0 y}{b^2} = 1.$$

Exercise 8.7 Learner's Book page 420

1. 4
2. $y = -\frac{1}{4}x + 5$
3. $15x + 4y + 9 = 0$
4. $\left(\frac{3}{2}\sqrt{13}, -9\right), \left(-\frac{3}{2}\sqrt{13}, -9\right)$

Lesson 8.8. Definition of polar coordinates

Learning objectives

Through examples, learners should be able to define polar coordinates, convert polar coordinates to Cartesian coordinates, and sketch a curve in polar form accurately.

Prerequisites

- ✔ Polar form of a complex number.
- ✔ Converting a complex number from polar form to algebraic form.
- ✔ Curve sketching.

Teaching Aids

Exercise book, pen, calculator and instruments of geometry

Activity 8.8 Learner's Book page 420



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

$$1. \text{ a) } |z| = \sqrt{1+1} = \sqrt{2} \quad \text{b) } |z| = \sqrt{1+1} = \sqrt{2}$$

$$2. \cos \theta = \frac{1}{\sqrt{2}} \Rightarrow \theta = \pm \frac{\pi}{4} + 2k\pi$$

$$\sin \theta = \frac{1}{\sqrt{2}} \Rightarrow \theta = \begin{cases} \frac{\pi}{4} + 2k\pi \\ \frac{3\pi}{4} + 2k\pi \end{cases}$$


$$\text{As } -\pi < \theta \leq \pi, \text{ we take } \theta = \frac{\pi}{4}.$$

Synthesis

To form a polar coordinate system in the plane, we fix a point O called the **pole** (or origin) and construct from O an initial ray called the **polar axis**. Then each point P in the plane can be assigned **polar coordinates** (r, θ) as follows:

- ① r is the directed distance from O to P .
- ② θ is the directed angle, counterclockwise from polar axis to the segment \overline{OP} .

To convert rectangular coordinates (a, b) to polar coordinates is the same as to find the modulus and argument of complex number $z = a + bi$.



From the above figure $r = \overline{OP}$, $x = r \cos \theta$, $y = r \sin \theta$; ox is a polar axis.

Exercise 8.8 Learner's Book page 425

1. a) and e); b) and g); c) and h); d) and f)
2. a) $(2, 2k\pi)$ and $(-2, (2k+1)\pi), k \in \mathbb{Z}$
 b) $(2, (2k+1)\pi)$ and $(-2, 2k\pi), k \in \mathbb{Z}$
 c) $\left(2, \frac{\pi}{2} + 2k\pi\right)$ and $\left(-2, \frac{\pi}{2} + (2k+1)\pi\right), k \in \mathbb{Z}$
 d) $\left(2, \frac{3\pi}{2} + 2k\pi\right)$ and $\left(-2, \frac{3\pi}{2} + (2k+1)\pi\right), k \in \mathbb{Z}$

3. a) $(3,0)$ b) $(-3,0)$ c) $(-1,\sqrt{3})$
 d) $(1,\sqrt{3})$ e) $(3,0)$ f) $(1,\sqrt{3})$
 g) $(-3,0)$ h) $(-1,\sqrt{3})$
4. a) $x+y=1$ b) $x=3$ c) $x=y$
 d) $x-3y=3$ e) $x^2+y^2=9$ f) $\frac{x^2}{16}+\frac{y^2}{25}=1$
 g) $x^2+(y-2)^2=4$ h) $(x-3)^2+(y+1)^2=4$
 i) $y^2=3x$

Lesson 8.9. Polar equation of a conic

Learning objectives

Through examples, learners should be able to find polar equation of a conic or change from polar equation to Cartesian equation accurately.

Prerequisites

- ✔ Cartesian equation of a conic.
- ✔ Conversion formulae from polar form to Cartesian form and vice versa.

Teaching Aids

Exercise book, pen and calculator

Activity 8.9 Learner's Book page 426



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Polar coordinates

$$x = r \cos \theta, \quad y = r \sin \theta, \quad r^2 = x^2 + y^2, \quad \tan \theta = \frac{y}{x}, \quad x \neq 0$$

$$y^2 = 1 + 2x \Leftrightarrow y^2 = 2x + 1 \Leftrightarrow y^2 - 2x = 1$$

$$\Leftrightarrow r^2 \sin^2 \theta - 2r \cos \theta = 1$$

$$2. \text{ a) } r = \frac{6}{2 + \cos \theta} \Leftrightarrow 2r + r \cos \theta = 6$$

$$\Leftrightarrow 2\sqrt{x^2 + y^2} + x = 6$$

$$\Leftrightarrow 2\sqrt{x^2 + y^2} = 6 - x$$

Squaring both sides gives

$$4(x^2 + y^2) = (6 - x)^2 \Leftrightarrow 4x^2 + 4y^2 = 36 - 12x + x^2$$

$$\Leftrightarrow 3(x^2 + 4x) + 4y^2 = 36 \Leftrightarrow 3(x^2 + 4x) + 4y^2 = 36$$

$$\Leftrightarrow 3(x + 2) + 4y = 48 \Leftrightarrow 3(x + 2)^2 + 4y^2 = 48$$

Dividing each term by 48, we get

$$\Leftrightarrow \frac{(x + 2)^2}{16} + \frac{y^2}{12} = 1$$

b) This is equation of a horizontal ellipse of centre $(-2, 0)$, major axis 8, minor axis $4\sqrt{3}$, eccentricity

$$e = \frac{\sqrt{a^2 - b^2}}{a} = \frac{\sqrt{16 - 12}}{4} = \frac{1}{2}, \text{ vertices } (2, 0), (-6, 0),$$

foci $(0, 0), (-4, 0)$.

Synthesis

Using polar coordinates, there is an alternative way to define a conic. In polar equation of a conic, the pole is the focus of the conic. We use the following relations:

$$x = r \cos \theta, \quad y = r \sin \theta, \quad r^2 = x^2 + y^2, \quad \tan \theta = \frac{y}{x}, \quad x \neq 0$$

The polar equation of conic with eccentricity e , focus at the origin, whose directrix $x = -p$ has equation $r = \frac{ep}{1 + e \cos \theta}$ where (r, θ) are polar coordinates of any point P lying on the conic.

The given conic is an **ellipse** if $e < 1$, a circle if $e = 1$, a **parabola** if $e = 1$, a hyperbola if $e > 1$.

Exercise 8.9 Learner's Book page 427

1. a) $r = \frac{1}{1 + \cos \theta}$ b) $r = \frac{2}{1 - 2 \sin \theta}$

2. a) $3x^2 + 4y^2 - 4x = 2$ b) $x^2 = 2 - 4y$

3. $r = \frac{k}{1 + e \cos \theta} \Leftrightarrow r + er \cos \theta = k$

$$\Leftrightarrow \sqrt{x^2 + y^2} + ex = k \Leftrightarrow \sqrt{x^2 + y^2} = k - ex$$

Squaring both sides, we get

$$x^2 + y^2 = k^2 - 2kex + e^2x^2 \Leftrightarrow x^2 - e^2x^2 + y^2 + 2kex - k^2 = 0$$

$$\Leftrightarrow (1 - e^2)x^2 + y^2 + 2kex - k^2 = 0 \text{ as required.}$$

Lesson 8.10. Polar equation of a straight line

Learning objectives

Given Cartesian equation of a straight line, learners should be able to find polar equation of that straight line or change from polar equation to Cartesian equation accurately.

Prerequisites

- ✔ Cartesian equation of a straight line.
- ✔ Conversion formulae from polar form to Cartesian form and vice versa.

Teaching Aids

Exercise book, pen and calculator



Activity 8.10 Learner's Book page 427

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

Polar coordinates are $x = r \cos \theta$, $y = r \sin \theta$.

Replacing x and y by their values from polar coordinates in $3x - 2y + 6 = 0$, we get

$$\begin{aligned} 3(r \cos \theta) - 2(r \sin \theta) + 6 &= 0 \Leftrightarrow 3r \cos \theta - 2r \sin \theta = -6 \\ \Leftrightarrow r(3 \cos \theta - 2 \sin \theta) &= -6 \Leftrightarrow -r(2 \sin \theta - 3 \cos \theta) = -6 \\ \Leftrightarrow 2 \sin \theta - 3 \cos \theta &= \frac{6}{r} \Leftrightarrow \frac{1}{r} = \frac{1}{3} \sin \theta - \frac{1}{2} \cos \theta \end{aligned}$$

Synthesis

The polar equation of a straight line is

$$\frac{1}{r} = A \cos \theta + B \sin \theta, \quad A, B \in \mathbb{R} \text{ and } A \text{ and } B \text{ are not}$$

all zero.

From general equation of a line in Cartesian plane, we get the polar equation of the given line.

$$\text{In fact, } Ax + By + C = 0 \Leftrightarrow Ax + By = -C \Leftrightarrow Ax + By = -C$$

$$\Leftrightarrow Ar \cos \theta + Br \sin \theta = -C$$

$$\Leftrightarrow r(A \cos \theta + B \sin \theta) = -C$$

$$\Leftrightarrow A \cos \theta + B \sin \theta = -\frac{C}{r}$$

Therefore, the polar equation of a straight line

$$Ax + By + C = 0 \text{ is } A \cos \theta + B \sin \theta = -\frac{C}{r}.$$

Exercise 8.10 **Learner's Book page 428**

1.
$$r = \frac{4}{\cos \theta + \sqrt{3} \sin \theta}$$

2.
$$r = \frac{2}{\cos \theta - \sin \theta}$$

3.
$$r = \frac{\sqrt{3}}{\sqrt{3} \cos \theta - 2 \sin \theta}$$

4.
$$r = \frac{\sqrt{5}}{\cos \theta - 2 \sin \theta}$$

Lesson 8.11. Polar form of a circle**Learning objectives**

Given a Cartesian equation of a circle, learners should be able to find polar equation of that circle or change from polar equation to Cartesian equation correctly.

Prerequisites

- ✔ Cartesian equation of a circle.
- ✔ Conversion formulae from polar form to Cartesian form and vice versa.

Teaching Aids

Exercise book, pen and calculator

Activity 8.11 Learner's Book page 429

In this lesson, the following generic competence and cross-cutting issues are to be addressed:


- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education



Answers

From the figure below,

$$\overline{OC} = \rho, \overline{OP} = r, \overline{CP} = R$$



Cosine law here is: $\overline{CP}^2 = \overline{CO}^2 + \overline{OP}^2 - 2\overline{OC}\overline{OP}\cos(\theta - \alpha)$

$$\Rightarrow R^2 = \rho^2 + r^2 - 2\rho r \cos(\theta - \alpha)$$

$$\Rightarrow r^2 = R^2 - \rho^2 + 2\rho r \cos(\theta - \alpha)$$

Synthesis

The polar equation of a circle with centre (ρ, α) and radius R is

$$r^2 = R^2 - \rho^2 + 2r\rho \cos(\theta - \alpha)$$

Exercise 8.11 Learner's Book page 430

1. $r = 6 \cos \theta$

2. $r = 4 \sin \theta$

3. $r = -\cos \theta$

4. $r = -2 \sin \theta$

Lesson 8.12. Applications of conics

Learning objectives

By reading textbooks or accessing internet, learners should be able to apply conics in real life problems perfectly.

Prerequisites

- ✔ Equations of conics.

Teaching Aids

Exercise book and pen

Activity 8.12 Learner's Book page 430



In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Research and problem solving
- ✔ Peace and values education
- ✔ Inclusive education

Answers



The Pythagorean concept of a spherical Earth offers a simple surface that is mathematically easy to deal with. Many astronomical and navigational computations use it as a surface representing the Earth.

The idea of a planar or flat surface for Earth, however, is still sufficient for surveys of small areas, as the local topography is far more significant than the curvature. Plane-table surveys are made for relatively small areas, and no account is taken of the curvature of the Earth. A survey of a city would likely be computed as though the Earth were a plane surface; the size of the city. For such small areas, exact positions can be determined relative to each other without considering the size and shape of the entire Earth.

The simplest model for the shape of the entire Earth is a sphere. The Earth's radius is the distance from Earth's centre to its surface, about 6,371 kilometres (3,959 mi). While "radius" normally is a characteristic of perfect spheres, the Earth deviates from a perfect sphere by only a third of a percent, sufficiently close to treat it as a sphere in many contexts and justifying the term "the radius of the Earth".

Synthesis

The orbits of planets are ellipses with the sun at one focus. For most planets, these ellipses have very small eccentricity, so they are nearly circular. However, the Mercury and Pluto, the innermost and outermost known planets, have visibly elliptical orbits.

Excercise 8.12 Learner's Book page 433

1. a) 6 cm b) 10 cm
2. a) (i) $7\sqrt{5}$ ii) $7\sqrt{5} - 13 = 133 \text{ mm}$
 b) (i) 1 ii) AC has equation $y = -x + 10$
3. $(x-5)^2 + (y-13)^2 = 9$
4. $\sqrt{5} \text{ m}$
5. The minimum altitude is 272 miles above the Earth
 The maximum altitude is 648 miles above the Earth.
6. $\frac{x^2}{900} - \frac{y^2}{14400.3636} = 1$
7. Then the equation for the elliptical ceiling is:

$$\frac{x^2}{625} + \frac{(y-5)^2}{400} = 1$$
 and the height of the ceiling above each whispering point is $y = 21$.

Summary of the unit

1. Generalities on conic sections

Parabolas, circles, ellipses and hyperbolas are called conics because they are curves in which planes intersect right circular cones.

2. Parabola

A parabola is the set of all points in plane that are equidistant from a fixed line (called directrix) and a fixed point (called focus) not on the line.

Important result relating to different parabolas

Equation	$y^2 = 4ax$	$x^2 = 4ay$
Focus	$(a, 0)$	$(0, a)$
Directrix	$x = -a$	$y = -a$
Principal axis(the line through the focus perpendicular to the directrix)	$y = 0$	$x = 0$
Vertex (point where the parabola crosses its principal axis)	$(0, 0)$	$(0, 0)$
Length of latus rectum (length of chord through a focus and perpendicular to the principal axis)	$4a$	$4a$
Equation of latus rectum	$x = a$	$y = a$
Ends of latus rectum	$(a, \pm 2a)$	$(\pm 2a, a)$

Replacing x with $(x-h)$ has the effect of shifting the graph of an equation by $|h|$ units to the right if h is positive, to the left if h is negative.

Similarly, replacing y with $(y-k)$ has the effect of shifting the graph by $|k|$ units up if k is positive and down if k is negative.

Equation	$(y-k)^2 = 4p(x-h)$	$(x-h)^2 = 4p(y-k)$
Focus	$(h+p, k)$	$(h, k+p)$
Directrix	$x = h-p$	$y = k-p$
Principal axis(the line through the focus perpendicular to the directrix)	$y = k$	$x = h$

Vertex (point where the parabola crosses its principal axis)	(h, k)	(h, k)
--------------------------------------------------------------	----------	----------

Parametric equations of parabola are

$$\begin{cases} x = at^2 \\ y = 2at \end{cases} \text{ where } t \text{ is a parameter.}$$

The tangent line at point (x_0, y_0) , on parabola $y^2 = 4ax$, is given by

$$T \equiv y_0 y = 2a(x + x_0)$$

3. Ellipse

Ellipse is a set of all points in the plane, the sum of whose distances from two fixed points (called foci) is a given positive constant.

Important facts to different ellipses

Equation of Standard form	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, a > b > 0$	$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, 0 < a < b$
Coordinates of centre	$(0, 0)$	$(0, 0)$
Coordinates of vertices	$(a, 0)$ and $(-a, 0)$	$(0, b)$ and $(0, -b)$
Length of major axis	$2a$	$2b$
Equation of major axis	$y = 0$	$x = 0$
Length of minor axis	$2b$	$2a$
Equation of minor axis	$x = 0$	$y = 0$
Eccentricity (ratio of semi-focal separation and the semi-major axis)	$b^2 = a^2(1 - e^2) \Rightarrow e = \frac{\sqrt{a^2 - b^2}}{a}$	$a^2 = b^2(1 - e^2) \Rightarrow e = \frac{\sqrt{b^2 - a^2}}{b}$

Coordinates of foci	$(ae, 0)$ and $(-ae, 0)$ $\Leftrightarrow (\pm\sqrt{a^2 - b^2}, 0)$	$(0, be)$ and $(0, -be)$ $\Leftrightarrow (0, \pm\sqrt{b^2 - a^2})$
Equation of directrices	$x = \pm \frac{a}{e}$	$y = \pm \frac{b}{e}$
Length of latus rectum	$\frac{2b^2}{a}$	$\frac{2a^2}{b}$
Equations of latus rectum	$x = \pm ae$	$y = \pm be$

Parametric equations of ellipse with centre (x_0, y_0) are

$$\begin{cases} x = x_0 + a \cos t \\ y = y_0 + b \sin t \end{cases} \text{ where } t \text{ is a parameter and } t \in (-\pi, \pi].$$

The tangent line at point (x_0, y_0) , on ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, is given by

$$T \equiv \frac{x_0 x}{a^2} + \frac{y_0 y}{b^2} = 1$$

4. Hyperbola

Hyperbola is a set of all points in the plane, the difference of whose distances from two fixed points (foci) is a given positive constant

Important facts to different hyperbolas

Equation of Standard form	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$	$\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$
Coordinates of centre	$(0, 0)$	$(0, 0)$
Coordinates of vertices	$(a, 0)$ and $(-a, 0)$	$(0, b)$ and $(0, -b)$
Length of transverse axis	$2a$	$2b$
Equation of transverse axis	$y = 0$	$x = 0$

Equation of conjugate axis	$x = 0$	$y = 0$
Coordinates of foci	$(ae, 0)$ and $(-ae, 0)$ $\Leftrightarrow (\pm\sqrt{a^2 + b^2}, 0)$	$(0, be)$ and $(0, -be)$ $\Leftrightarrow (0, \pm\sqrt{a^2 + b^2})$
Equation of directrices	$x = \pm \frac{a}{e}$	$y = \pm \frac{b}{e}$
Length of latus rectum	$\frac{2b^2}{a}$	$\frac{2a^2}{b}$
Equations of latus rectum	$x = \pm ae$	$y = \pm be$
Eccentricity	$b^2 = a^2(1 - e^2)$	$a^2 = b^2(1 - e^2)$

Parametric equations of hyperbola whose centre (x_0, y_0) are

$$\begin{cases} x = x_0 + a \sec t \\ y = y_0 + b \tan t \end{cases} \text{ where } t \text{ is a parameter and}$$

$$t \in \left] -\frac{\pi}{2}, \frac{\pi}{2} \right[\cup \left] \frac{\pi}{2}, \frac{3\pi}{2} \right[$$

The tangent line at point (x_0, y_0) , on hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$, is given by

$$T \equiv \frac{x_0 x}{a^2} - \frac{y_0 y}{b^2} = 1$$

5. Polar coordinates

To form a polar coordinate system in the plane, we fix a point O called the pole (or origin) and construct from O an initial ray called the polar axis. Then, each point P in the plane can be assigned polar coordinates (r, θ) as follows:

- ⑦ r is the directed distance from O to P .
- ⑦ θ is the directed angle, counterclockwise from polar axis to the segment \overline{OP} .

In polar coordinate system, the coordinates (r, θ) , $(r, \theta + 2k\pi)$, $k \in \mathbb{Z}$ and $(-r, \theta + (2k+1)\pi)$ represent the same point.

Coordinate conversion

The polar coordinates (r, θ) of a point are related to the rectangular coordinates (x, y) as follows:

$$x = r \cos \theta, \quad y = r \sin \theta, \quad r^2 = x^2 + y^2, \quad \tan \theta = \frac{y}{x}, \quad x \neq 0$$

Polar equation of a conic

A conic curve with eccentricity e , focus at the origin, whose directrix $x = -p$ has equation $r = \frac{ep}{1 + e \cos \theta}$ where (r, θ) are polar coordinates of any point P lying on the conic. It is an **ellipse** if $e < 1$, a parabola if $e = 1$, a **hyperbola** if $e > 1$.


6. Applications

Eccentricities of orbits of the planets


The orbits of planets are ellipses with the sun at one focus. For most planets, these ellipses have very small eccentricity, so they are nearly circular. However, the Mercury and Pluto, the innermost and outermost known planets, have visibly elliptical orbits.

End of Unit Assessment answers Learner's Book page 441


1. a) Parabola, *focus* $(0, -2)$, principal axis is $x = 0$.
 - b) Ellipse, *foci* $(\pm 1, 0)$, semi-major axis $\sqrt{2}$, semi-minor axis 1.
 - c) Parabola, *focus* $\left(-\frac{1}{12}, 0\right)$, principal axis is $y = 0$.
 - d) Hyperbola, *foci* $(0, \pm 4)$, transverse axis $x = 0$, conjugate axis $y = 0$, asymptotes $y = \pm x$.
 - e) Parabola, *focus* $\left(\frac{15}{4}, 1\right)$, principal axis is $y = 1$.
 - f) Hyperbola, *foci* $(\pm\sqrt{10}, 0)$, transverse axis is $y = 0$, conjugate axis $x = 0$, asymptotes $y = \pm 2x$.
2. a) Intersection: $\left\{\left(\frac{109}{10}, -\frac{91}{20}\right)\right\}$




b) Intersection: $\left\{ \left(\frac{\sqrt{10}}{2}, 5 \right), \left(-\frac{\sqrt{10}}{2}, 5 \right) \right\}$



c) Intersection: $\{(-2, 1), (-2, -1), (2, 1), (2, -1)\}$



d) Intersection: $\{(2, \sqrt{3}), (2, -\sqrt{3}), (-2, \sqrt{3}), (-2, -\sqrt{3})\}$



$$3. \frac{(x-4)^2}{32} + \frac{(y-3)^2}{36} = 1$$

$$4. 8xy - 4x - 4y + 1 = 0$$

5. a) The line $y = kx$ intersects the given conic once, when there is a unique solution.

As $AC - B^2 = 0$, the given conic is a parabola.

Solving the equations of line and conic taken together, we get equation $(2-k)^2 x^2 + 6x + 1 = 0$.

If $k = 2$, we get a linear equation: $6x + 1 = 0$.

$$\Rightarrow x = -\frac{1}{6} \text{ and } y = -\frac{1}{3}$$

That is, the line touches the conic at $\left(-\frac{1}{6}, -\frac{1}{3}\right)$.

If $k \neq 2$, we have quadratic equation which can be solved using discriminant;

$$\Delta = b^2 - 4ac \Rightarrow \Delta = 9 - (2-k)^2 = (5-k)(1+k).$$

We have a unique solution if $\Delta = 0$, i.e. $k = 5$ or $k = -1$.


Therefore, the line $y = kx$ intersects the given conic once if $k = 2$ or $k = 5$ or $k = -1$.

b) The line $y = kx$ cuts the given conics in two points if $\Delta > 0$, $-1 < k < 5$ and $k \neq 2$.


c) The line $y = kx$ does not intersect the given conics if $\Delta < 0$.

Thus, $k < -1$ or $k > 5$.


6. a)




b)




c)




d)



e)



f)



7. a) $(y+2)^2 = 4(x+3), V(-3, -2), F(-1, -3), D \equiv x = -3$
 b) $(x-1)^2 = 8(y+7), V(1, -7), F(1, -5), D \equiv y = -9$
 c) $\frac{(x+2)^2}{6} + \frac{(y+1)^2}{9} = 1, F(-2, \pm\sqrt{3}-1), V(-2, \pm 3-1), C(-2, -1)$
 d) $\frac{(x-2)^2}{3} + \frac{(y-3)^2}{2} = 1, F(3, 3) \text{ and } F(1, 3), V(\pm\sqrt{3}+2, 3), C(2, 3)$
 e) $\frac{(x-2)^2}{4} - \frac{(y-2)^2}{5} = 1, F(5, 2) \text{ and } F(-1, 2), V(4, 2) \text{ and } V(0, 2), C(2, 2),$
 $A \equiv (y-2) = \pm \frac{\sqrt{5}}{2}(x-2)$
 f) $(y+1)^2 - (x+1)^2 = 1, F(-1, \sqrt{2}-1) \text{ and } F(-1, -\sqrt{2}-1),$
 $V(-1, 0) \text{ and } V(-1, -2)$
 $C(-1, -1), A \equiv (y+1) = \pm(x+1)$
8. a) $(3\sqrt{3}, 3)$ b) $\left(-\frac{7}{2}, \frac{7\sqrt{3}}{2}\right)$ c) $(4\sqrt{2}, 4\sqrt{2})$
 d) $(5, 0)$ e) $\left(-\frac{7\sqrt{3}}{2}, \frac{7}{2}\right)$ f) $(0, 0)$

9. (i) a) $(5, \pi)$ b) $\left(4, \frac{11\pi}{6}\right)$ c) $\left(2, \frac{3\pi}{2}\right)$
 d) $\left(8\sqrt{2}, \frac{5\pi}{4}\right)$ e) $\left(6, \frac{2\pi}{3}\right)$ f) $\left(\sqrt{2}, \frac{\pi}{4}\right)$
- (ii) a) $(-5, 0)$ b) $\left(-4, \frac{5\pi}{6}\right)$ c) $\left(-2, \frac{\pi}{2}\right)$
 d) $\left(-8\sqrt{2}, \frac{\pi}{4}\right)$ e) $\left(-6, \frac{5\pi}{3}\right)$ f) $\left(-\sqrt{2}, \frac{5\pi}{4}\right)$
10. a) $x^2 + y^2 = 5$; circle b) $y = 4$; straight line
 c) $y^2 = 1 + 2x$; parabola d) $x^2 - 3y^2 - 8y = 4$; hyperbola
 e) $3y - 4x = 5$; straight line
 f) $3x^2 + 4y^2 - 12x = 36$; ellipse
 g) $x^2 + y^2 + 4x = 0$; circle
11. a) Proof
 b)

Planets	Perihelion (astronomical units)	Aphelion (astronomical units)
Mercury	0.3075	0.4667
Venus	0.7184	0.7282
Earth	0.9833	1.0167
Mars	1.3817	1.6663
Jupiter	4.9512	5.4548
Saturn	9.0210	10.0570
Uranus	18.2977	20.0623
Neptune	29.8135	30.3065

c)

Planets	Polar equation for the ellipse with eccentricity e and semi-major axis a : $r = \frac{a(1-e^2)}{1+e \cos \theta}$
Mercury	$r = \frac{0.3707}{1+0.2056 \cos \theta}$
Venus	$r = \frac{0.7233}{1+0.0068 \cos \theta}$
Earth	$r = \frac{0.9997}{1+0.0167 \cos \theta}$
Mars	$r = \frac{1.5107}{1+0.0934 \cos \theta}$
Jupiter	$r = \frac{5.1908}{1+0.0484 \cos \theta}$
Saturn	$r = \frac{9.5109}{1+0.0543 \cos \theta}$
Uranus	$r = \frac{19.1394}{1+0.0460 \cos \theta}$
Neptune	$r = \frac{30.0580}{1+0.082 \cos \theta}$


12. $7.25m$

13. $\frac{x^2}{25} + \frac{y^2}{9} = 1$

14. $\frac{x^2}{16} + \frac{y^2}{7} = 1$

15. $10\sqrt{15}$ inches

16. $\frac{x^2}{1.1025} - \frac{y^2}{7.8975}$



Random Variables

Learner's Book pages 445 - 493

Key unit competence

Calculate and interpret the parameters of a random variable (discrete or continuous) including binomial and the Poisson distributions.

Vocabulary or key words concepts

Random variable: A variable which can assume numerical values each of which can correspond to one and only one of the events.

Discrete random variable: Random variable which takes only finite values between its limits.

Binomial distribution: Probability distribution for which probabilities are of successive terms of the binomial expansion $(q + p)^n$.

Poisson distribution: Discrete distribution used as a model for the number of events in a specific time period.

Continuous random variable: Random variable for which the possible values are all real values in some interval.

Guidance on the problem statement

The problem statement: "A life insurance company has determined that on the average it receives 10 death claims per day. Find the probability that the company receives at least five death claims on a randomly selected day."

This problem is solved using a special distribution called Poisson distribution that will be studied in this unit.

List of lessons

No	Lesson title	Number of periods
1	Probability density function of a discrete random variable	3
2	Expected value, variance and standard deviation of a discrete random variable	3
3	Cumulative distribution function of a discrete random variable	3
4	Binomial distribution	4
5	Expected value, variance and standard deviation of a binomial distribution	4
6	Poisson distribution	4
7	Probability density function of a continuous random variable	4
8	Cumulative distribution function of a continuous random variable	4
9	Expected value, variance and standard deviation of a continuous random variable	4
Total periods		33

Lesson development

Lesson 9.1. Probability density function of a discrete random variable

Learning objectives

Through examples, learners should be able to identify a discrete random variable and to find its probability distribution correctly.

Prerequisites

Finding probability of an event.

Teaching Aids

Exercise book, pen and calculator



Activity 9.1 Learner's Book page 445

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

All balls are 6 with 4 red balls and 2 black balls

$$\text{Probability of 0 red balls: } P(BBB) = \frac{2}{6} \times \frac{2}{6} \times \frac{2}{6} = \frac{1}{27}$$

Probability of 1 red ball:

$$P(RBB) + P(BRB) + P(BBR) = \frac{4}{6} \times \frac{2}{6} \times \frac{2}{6} + \frac{2}{6} \times \frac{4}{6} \times \frac{2}{6} + \frac{2}{6} \times \frac{2}{6} \times \frac{4}{6} = \frac{2}{9}$$

Probability of 2 red balls:

$$P(RRB) + P(RBR) + P(BRR) = \frac{4}{6} \times \frac{4}{6} \times \frac{2}{6} + \frac{4}{6} \times \frac{2}{6} \times \frac{4}{6} + \frac{2}{6} \times \frac{4}{6} \times \frac{4}{6} = \frac{4}{9}$$

$$\text{Probability of 3 red balls: } P(RRR) = \frac{4}{6} \times \frac{4}{6} \times \frac{4}{6} = \frac{8}{27}$$

x	0	1	2	3
P	$\frac{1}{27}$	$\frac{2}{9}$	$\frac{4}{9}$	$\frac{8}{27}$

x takes on whole numbers only. The sum of obtained probabilities is 1. This is the probability of the sample space.

Synthesis

A variable X which can assume numerical values each of which can correspond to one and only one of the events is called a **random variable** (or **stochastic variable**). A random variable X is said to be a discrete random variable, if it takes only finite values between its limits.

It is convenient to introduce the probability function $P(X = x_i) = p_i$, also called the **probability distribution** satisfying

1. $P(X = x_i) \geq 0$
2. $\sum_i P(X = x_i) = 1$, where the sum is taken over all values of x_i .

The **probability density function** (p.d.f), $F(x)$, is a function that allocates probabilities to all distinct values that X can take on.

Exercise 9.1 Learner's Book page 449

1. X is a random variable if $\sum_{i=1}^3 P(X = x_i) = 1$.

$$\sum_{i=1}^3 P(X = x_i) = P(X = 2) + P(X = 3) + P(X = 4) = \frac{1}{6} + \frac{2}{6} + \frac{3}{6} = 1$$

Therefore, X is a random variable.

2. a) $p = \frac{1}{10}$ b) $P(X \geq 2) = \frac{4}{5}$
3. $a = \frac{1}{2}$

Lesson 9.2. Cumulative distribution of discrete random variable

Learning objectives

Given a discrete random variable, learners should be able to find its cumulative distribution precisely.

Prerequisites

- ⑦ Probability density function of a discrete random variable.

Teaching Aids

Exercise book, pen and calculator



Activity 9.2 Learner's Book page 449

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

Cumulative probabilities are found by adding the probability up to each column of the table. In the table, we find the cumulative probability for one head by adding the probabilities for zero and one. The cumulative probability for two heads is found by adding the probabilities for zero, one, and two. We continue with this procedure until we reach the maximum number of heads, in this case four, which should have a cumulative probability of 1.00 because 100% of trials must have four or fewer heads.

Then,

Heads	0	1	2	3	4
Probability	$\frac{1}{16}$	$\frac{4}{16}$	$\frac{6}{16}$	$\frac{4}{16}$	$\frac{1}{16}$
Cumulative Probability	$\frac{1}{16}$	$\frac{5}{16}$	$\frac{11}{16}$	$\frac{15}{16}$	$\frac{16}{16} = 1$

Synthesis

To find a **cumulative probability** we add the probabilities for all values qualifying as “less than or equal” to the specified value. Then,

The cumulative distribution function of a random variable X is the function $F(x) = P(X \leq x)$.

Exercise 9.2 Learner's Book page 451

1. Cumulative distribution function

$$F(x) = \begin{cases} 0, & x < 0 \\ \frac{1}{4}, & 0 \leq x < 1 \\ \frac{3}{4}, & 1 \leq x < 2 \\ 1, & x \geq 2 \end{cases}$$

2. Cumulative distribution

x	2	3	4	5	...	12
$F(x)$	$\frac{1}{36}$	$\frac{3}{36}$	$\frac{6}{36}$	$\frac{10}{36}$...	1

3. Probability distribution

x	1	2	3	4	5
$P(X=x)$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{5}$	$\frac{1}{5}$

Lesson 9.3. Expected value, variance and standard deviation of a discrete random variable**Learning objectives**

Given a discrete random variable, learners should be able to calculate the expected value, variance and standard deviation correctly.

Prerequisites

⑦ Probability distribution of a discrete random variable.

Teaching Aids

Exercise book, pen and calculator



Activity 9.3 Learner's Book page 452

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

x	$P(X = x)$	$xP(X = x)$	$x^2P(X = x)$
1	0.2	0.2	0.2
2	0.5	1	2.0
3	0.3	0.9	2.7
Sum	1.0	2.1	4.9

Synthesis

The **expected value** of random variable X , which is the **mean** of the probability distribution of X , is denoted and

defined by $\mu = E(X) = \sum_{i=1}^n x_i P(X = x_i)$.

The variance of random variable X is denoted and defined by

$$\sigma^2 = \text{Var}(X) = E(X^2) - [E(X)]^2 \text{ or}$$

$$\text{var}(X) = E(X^2) - [E(X)]^2$$

The standard deviation of random variable X , is

$$\sigma = \text{SD}(X) = \sqrt{\text{Var}(X)}.$$

Exercise 9.3 Learner's Book page 454

1. a) 1.9 b) 2.4 c) 0.4 d) 9.23
2. a) $\frac{2}{3}$ b) $\frac{26}{9}$ c) $\frac{152}{81}$ d) $\frac{2\sqrt{38}}{9}$
3. Expected value: 3.5, variance: 2.9, standard deviation: 1.7

Lesson 9.4. Binomial distribution

Learning objectives

Through examples, learners should be able to identify a binomial distribution and find its probability distribution accurately.

Prerequisites

- ✔ Powers.
- ✔ Combination of n objects taken from n objects.

Teaching Aids

Exercise book, pen and calculator



Activity 9.4 Learner's Book page 455

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Probability of the sequence $SSSSFFFFFF$ is

$$P(S)P(S)P(S)P(S)P(F)P(F)P(F)P(F)P(F)P(F) = ppppqqqqqq$$

$$= p^4q^6$$

2. From 1),

$$\underbrace{P(S)P(S)\dots}_{r \text{ times}} \times \underbrace{P(F)P(F)\dots}_{n-r \text{ times}} = \underbrace{pp\dots}_{r \text{ times}} \times \underbrace{qq\dots}_{n-r \text{ times}}$$

$$= p^r q^{n-r}$$

3. Different combinations that produce 4 heads are given by ${}^{10}C_4 p^4 q^6$.

4. Different combinations that produce r heads in n trials are given by ${}^n C_r p^r q^{n-r}$.

Synthesis

The probability of obtaining r successes in n independent trials is $b(r : n, p) = {}^n C_r p^r q^{n-r}$ for $0 \leq r \leq n$ where p is the probability of a success in each trial. This probability distribution is called the binomial distribution since the values of the probabilities are successive terms of the binomial expansion of $(q + p)^n$; that is why $b(r : n, p) = {}^n C_r p^r q^{n-r}$.

Each trial has two possible outcomes: success (p) and failure (q).

The outcome of the n trials are mutually independent and there will be r successes and $n - r$ failures.

Exercise 9.4 Learner's Book page 459

- | | | |
|---------------------------|--------------------------|------------|
| 1. $\frac{15}{64}$ | 2. 0.92 | 3. 0.65536 |
| 4. 0.19 | 5. 0.51 | |
| 6. a) 0.39 | b) 0.35 | c) 0.93 |
| 7. a) 0.26 | b) $\frac{41}{1679616}$ | c) 0.77 |
| 8. a) $\frac{8183}{8192}$ | b) $\frac{16807}{32768}$ | |

Lesson 9.5. Expected value, variance and standard deviation of a binomial distribution

Learning objectives

Given a binomial random variable, learners should be able to find expected value, variance and standard deviation correctly.

Prerequisites

- ▼ Binomial distribution

Teaching Aids

Exercise book, pen and calculator



Activity 9.5 Learner's Book page 460

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ▼ Critical thinking
- ▼ Communication
- ▼ Self confidence
- ▼ Cooperation, interpersonal management and life skills
- ▼ Peace and values education
- ▼ Inclusive education

Answers

$$1. E(X) = \sum_{i=1}^n x_i P(X = x_i) = 0 \times (1-p) + 1 \times p = p$$

$$\text{For } n \text{ trials, } E(X) = np$$

$$2. E(X^2) = 0^2 \times (1-p) + 1^2 p = p \text{ and for } n \text{ trials}$$

$$E(X^2) = np$$

$$\text{Var}(X) = E(X^2) - [E(X)]^2$$

$$= np - [np]^2$$

$$= np - n^2 p^2$$

$$= np(1-p)$$

$$= npq$$

Synthesis

Basing on the results from activity 9.5, the expected value (or mean) of a binomial distribution of a discrete random variable X is denoted and defined by $\mu = E(X) = np$ where n is the number of trials and p is the probability of success.

The variance of a binomial distribution of a discrete random variable X is denoted and defined by $\sigma^2 = Var(X) = npq$ where n is the number of trials, p is the probability of success and q is the probability of failure.

The standard deviation of a binomial distribution of a discrete random variable X is denoted and defined by $\sigma = \sqrt{Var(X)} = \sqrt{npq}$.

Exercise 9.5 Learner's Book page 463

1. $np > npq$, as $q < 1$
2. Mean is 30, standard deviation is 5
3. a) $E(X) = 2.5$ b) $var(X) = 1.875$
4. a) 0.117 b) 0.974
5. approximately 1 6. 11 7. ≈ 17

Lesson 9.6. Poisson distribution**Learning objectives**

Through examples, learners should be able to identify a poisson distribution and solve problems using poisson distribution correctly.

Prerequisites

- ⑦ Use of exponential and factorial notation.

Teaching Aids

Exercise book, pen and calculator



Activity 9.6 Learner's Book page 464

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. Dividing both sides of (1) by e^θ

$$\begin{aligned}
 1 &= \frac{\theta^0}{0!e^\theta} + \frac{\theta^1}{1!e^\theta} + \frac{\theta^2}{2!e^\theta} + \frac{\theta^3}{3!e^\theta} + \dots + \frac{\theta^n}{n!e^\theta} + \dots \\
 &= \frac{\theta^0 e^{-\theta}}{0!} + \frac{\theta^1 e^{-\theta}}{1!} + \frac{\theta^2 e^{-\theta}}{2!} + \frac{\theta^3 e^{-\theta}}{3!} + \dots + \frac{\theta^n e^{-\theta}}{n!} + \dots
 \end{aligned}$$

2. If we take $\lambda = \theta$, we have

$$1 = \frac{\lambda^0}{0!e^\lambda} + \frac{\lambda^1}{1!e^\lambda} + \frac{\lambda^2}{2!e^\lambda} + \frac{\lambda^3}{3!e^\lambda} + \dots + \frac{\lambda^n}{n!e^\lambda} + \dots$$

3. Using the general term, $\frac{\lambda^n}{n!e^\lambda}$, and putting $n = x$, we

$$\text{have } P(X = x) = \frac{\lambda^x}{x!e^\lambda} \text{ or } P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$$

Synthesis

The probability density function of Poisson distribution is denoted $X \sim P(\lambda)$ and defined by

$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}, \quad x = 0, 1, 2, \dots$$

Where λ is a parameter which indicates the average number (the expected value) of events in the given time interval and $e \approx 2.718\dots$

For Poisson distribution with parameter λ , $E(x) = \lambda$ and $Var(x) = \lambda$.

Exercise 9.6 Learner's Book page 468

1. Wrong statement, because $\sigma = \sqrt{\lambda}$
2. 0.827008
3. 0.052129
4. 0.160623
5. 0.128387
6. 0.00000546

$$7. \sum_{x=0}^{14} \frac{e^{-20} (20)^x}{x!}$$

8. Considering the given table:

x	0	1	2	3	4	5	6	Total
f	103	143	98	42	8	4	2	400
$x \cdot f$	0	143	126	126	32	20	12	529

$$\text{Mean} = \frac{529}{400} = 1.32$$

Number of cells	Probability $P(x) = \frac{0.2674 \times (1.32)^x}{x!}$	Theoretical frequency
0	$\frac{0.2674 \times (1.32)^0}{0!} = 0.2674$	$0.2674 \times 400 \approx 107$
1	$\frac{0.2674 \times (1.32)^1}{1!} = 0.353$	$0.353 \times 400 \approx 141$
2	$\frac{0.2674 \times (1.32)^2}{2!} = 0.233$	$0.233 \times 400 \approx 93$
3	$\frac{0.2674 \times (1.32)^3}{3!} = 0.1025$	$0.1025 \times 400 \approx 41$
4	$\frac{0.2674 \times (1.32)^4}{4!} = 0.0338$	$0.0338 \times 400 \approx 14$

5	$\frac{0.2674 \times (1.32)^5}{5!} = 0.00893$	$0.00893 \times 400 \approx 4$
6	$\frac{0.2674 \times (1.32)^6}{6!} = 0.00196$	$0.00196 \times 400 \approx 1$

The expected (theoretical) frequencies are 107,141,93,41,14,4,1.

Lesson 9.7. Probability density function of a continuous random variable

Learning objectives

Through examples, learners should be able to identify a continuous random variable, and to find its probability distribution accurately.

Prerequisites

- ✔ Integration.
- ✔ Curve sketching.

Teaching Aids

Exercise book, pen, calculator, instruments of geometry.



Activity 9.7 Learner's Book page 470

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. The area under the curve of $f(x)$ is given by;

$$\int_{-1}^0 k(x+1)^2 dx + \int_0^1 kx dx . \text{ Then}$$

$$\int_{-1}^0 k(x+1)^2 dx + \int_0^1 kx dx = 1 \Rightarrow k \int_{-1}^0 (x^2 + 2x + 1) dx + k \int_0^1 dx = 1$$

$$\Rightarrow k \left[\frac{x^3}{3} + x^2 + x \right]_{-1}^0 + k [x]_0^1 = 1$$

$$\Rightarrow k \left(\frac{1}{3} - 1 + 1 \right) + k = 1 \Rightarrow \frac{4}{3}k = 1 \text{ or } k = \frac{3}{4}$$


2. Graph of $f(x) = \begin{cases} \frac{3}{4}(x+1)^2 & -1 \leq x \leq 0 \\ \frac{3}{4} & 0 < x \leq 1 \end{cases}$

$$f(x) = \frac{3}{4}(x+1)^2 \quad -1 \leq x \leq 0$$

x	-1	-0.3	-0.7	0
$f(x) = y$	0	0.37	0.07	0.43

$$f(x) = \frac{3}{4} \quad 0 < x \leq 1$$

x	0	1
$f(x) = y$	0.43	0.43



Synthesis

A random variable X is said to be continuous if its possible values are all real values in some interval.

A function $f(x)$ defined on an interval $[a, b]$ is a probability density function for a continuous random variable X distributed on $[a, b]$ if, whenever x_1 and x_2

satisfy $a \leq x_1 \leq x_2 \leq b$, we have $p(x_1 \leq X \leq x_2) = \int_{x_1}^{x_2} f(x) dx$

If X is a continuous random variable, then the probability that the values of X will fall between the values a and b is given by the area of the region lying below the graph of $f(x)$ and above the x -axis between a and b and this area is equal to 1.

Normally, $\int_{-\infty}^{+\infty} f(x) dx = 1$.

Exercise 9.7 Learner's Book page 474

1. a) $c = \frac{3}{2}$ b) $P\left(X \geq \frac{1}{2}\right) = \frac{7}{16}$
2. 0.693
3. a) $k = \frac{1}{4}$
 - b) i) $p(x < 1) = \frac{1}{4}$ ii) $p(x = 1) = 0$
 - iii) $p(x > 2.5) = 0.3125$
 - iv) $p[(0 < x < 2) / x \geq 1] = \frac{p(1 \leq x \leq 2)}{p(1 \leq x \leq 3)}$

$$= \frac{\int_1^2 \frac{1}{4} dx}{\int_1^2 \frac{1}{4} dx + \int_2^3 \frac{1}{4}(2x-3) dx} = \frac{1}{3}$$

Lesson 9.8. Cumulative distribution of continuous random variable

Learning objectives

Given a continuous random variable, learners should be able to find its cumulative distribution accurately.

Prerequisites

- ✔ Finite integrals

Teaching Aids

Exercise book, pen and calculator



Activity 9.8 Learner's Book page 475

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ✔ Critical thinking
- ✔ Communication
- ✔ Self confidence
- ✔ Cooperation, interpersonal management and life skills
- ✔ Peace and values education
- ✔ Inclusive education

Answers

1. For $x < 1$, $f(x) = 0$ and then $F(x) = \int_1^x 0 dx = 0$

2. For $1 \leq x \leq 3$, $\frac{1}{4}x$ and then

$$F(x) = \int \frac{1}{4}x dx = \left[\frac{1}{8}x^2 \right] = \frac{1}{8}x^2 - \frac{1}{8} = \frac{1}{8}$$

3. For $x > 3$, $f(x) = 0$ and then

$$F(x) = F(3) + \int_1^x 0 dx = \frac{x^2 - 1}{8} \text{ and } F(3) = \frac{3^2 - 1}{8} = 1$$

4. Hence,

$$F(x) = \begin{cases} 0, & x < 1 \\ \frac{x^2 - 1}{8}, & 1 \leq x \leq 3 \\ 1, & x > 3 \end{cases}$$

Synthesis

The **cumulative distribution** function of a continuous

random variable X is defined as: $F(x) = \int_{-\infty}^x f(t) dt$.

Where $F(x) = 0$, for $x \rightarrow -\infty$ and $F(x) = 1$, for $x \rightarrow +\infty$.

Exercise 9.8 Learner's Book page 477

$$1. F(x) = \begin{cases} 0, & x \leq -1 \\ \frac{1}{2}(x+1)^2, & -1 < x \leq 0 \\ 1 - \frac{(1-x)^2}{2}, & 0 < x < 1 \\ 1, & x \geq 1 \end{cases} \quad 2. F(x) = \begin{cases} 0, & x < 0 \\ \frac{x^2}{16}, & 0 \leq x \leq 4 \\ 1, & x > 4 \end{cases}$$

$$3. F(x) = \begin{cases} 0, & x < 0 \\ \frac{x^2}{6}, & 0 \leq x < 2 \\ \frac{-x^2}{3} + 2x - 2, & 2 \leq x < 3 \\ 1, & x \geq 3 \end{cases}$$

Lesson 9.9. Variance and standard deviation of a continuous random variable**Learning objectives**

Given a continuous random variable, learners should be able to find expected value, variance and standard deviation correctly.

Prerequisites

- ⑦ Probability density function of a continuous random variable.

Teaching Aids

Exercise book, pen and calculator



Activity 9.9 Learner's Book page 477

In this lesson, the following generic competence and cross-cutting issues are to be addressed:

- ④ Critical thinking
- ④ Communication
- ④ Self confidence
- ④ Cooperation, interpersonal management and life skills
- ④ Peace and values education
- ④ Inclusive education

Answers

$$\begin{aligned}
 1. \quad A \quad \int xf(x)dx &= 6 \int x(1-x)dx \\
 &= 6 \int_0^1 (x^2 - x^3)dx = 6 \left[\frac{1}{3}x^3 - \frac{1}{4}x^4 \right]_0^1 \\
 &= 6 \left[\frac{1}{3} - \frac{1}{4} \right] = 0.5
 \end{aligned}$$

$$\begin{aligned}
 2. \quad B &= \int_0^1 x^2 f(x)dx = 6 \int_0^1 x^3(1-x)dx \\
 &= 6 \int_0^1 (x^3 - x^4)dx = 6 \left[\frac{1}{4}x^4 - \frac{1}{5}x^5 \right]_0^1 \\
 &= 6 \left[\frac{1}{4} - \frac{1}{5} \right] = 0.3
 \end{aligned}$$

$$3. \quad B - A^2 = 0.3 - 0.25 = 0.05$$

Synthesis

The **mean**, μ , (or **expected value**, $E(X)$), of X is denoted and defined by;

$$\mu = E(X) = \int_a^b xf(x)dx$$

Also, expectation of function g of X is

$$E(g(x)) = \int_a^b g(x)f(x)dx$$

The variance $Var(x)$ or σ^2 is denoted and defined by

$$\sigma^2 = Var(X) = E(X^2) - [E(x)]^2.$$

The standard deviation is

$$\sigma = SD = \sqrt{Var(X)}.$$

Properties of $E(X)$ and $Var(X)$

$$\forall a, b \in \mathbb{R}$$

1. $E(a) = a$
2. $E(aX) = aE(X)$
3. $E(aX + b) = aE(X) + b$
4. $E(X + Y) = E(X) + E(Y)$
5. $E(aX^2 + b) \neq aE(X^2) + b$
6. $var(a) = 0$
7. $var(aX + b) = a^2 var(X)$
8. $var(aX + b) = a var(X)$

Exercise 9.9 Learner's Book page 482

1. a) $k = 1.44$
b) mean is 0.443, variance is 0.0827
2. a) $c = \frac{3}{2}$ b) $E(X) = 0, var(X) = \frac{3}{5}$
3. a) $a = 0.01$
b) $E(x) = 10, Var(x) = 16.6667$
 $E(9x) = 9E(x) = 90$ $Var(9x) = 9^2 Var(x) = 1350$

Summary of the unit

1. Discrete and finite random variables

1. Probability density function

A random variable X is said to be a **discrete random variable**, if it takes only finite values between its limits; for example, the number of learners appearing in a festival consisting of 400 learners is a discrete random variable which can assume values other than 0, 1, 2, ..., 400.

The **probability density function** (p.d.f), $F(x)$, is a function that allocates probabilities to all distinct values that X can take on.

If the initial probability is known, you can find successive probabilities using the following recurrence

$$\text{relation } P(X = x+1) = \left(\frac{n-x}{x+1}\right) \left(\frac{p}{q}\right) P(X = x).$$

2. Expectation, variance and standard deviation

The expected value of random variable X , which is the mean of the probability distribution of X , is denoted and defined by

$$\mu = E(X) = \sum_{i=1}^n x_i P(X = x_i).$$

Also, the expectation of any function $g(X)$ of the random variable X is

$$\mu = E(g(X)) = \sum_{i=1}^n g(x) P(X = x_i).$$

The variance of random variable X is denoted and defined by

$$\sigma^2 = \text{Var}(X) = \sum_{i=1}^n [x_i - \mu]^2 P(X = x_i).$$

This can be simplified to

$$\sigma^2 = \text{Var}(X) = E(X^2) - \mu^2$$

The standard deviation of random variable X , denoted by $SD(X)$, is the square root of the variance. That is

$$\sigma = SD(X) = \sqrt{\text{Var}(X)}.$$

Properties for mean and variance

$\forall a, b \in \mathbb{R}$

1. $E(a) = a$
2. $E(aX) = aE(X)$
3. $E(aX + b) = aE(X) + b$
4. $E(X + Y) = E(X) + E(Y)$
5. $E(aX^2 + b) \neq aE(X^2) + b$
6. $\text{var}(a) = 0$
7. $\text{var}(aX) = a^2 \text{var}(X)$
8. $\text{var}(aX + b) = a^2 \text{var}(X)$

3. Binomial distribution (Law of Bernoulli)

For binomial probability distribution, we are interested in the probabilities of obtaining r successes in n trials, in other word r successes and $n-r$ failures in n attempts.

Binomial distribution is denoted

$$b(r : n, p) = {}^n C_r P^r q^{n-r}, r = 0, 1, 2, \dots, n$$

The constant n, p, q are called parameters of the binomial distribution.

The following assumptions are made:

- There is a fixed number (n) of trials.
- The probability of success is the same for each trial.
- Each trial is independent of all other trials.

Note that $p + q = 1$

For N set of n trial, the successes $0, 1, 2, \dots, r, \dots, n$ are given by $N(p + q)^n$, which is called binomial distribution.

The expected value (or mean) of a binomial distribution of a discrete random variable X is denoted and defined

by $\mu = E(X) = np$ where n is the number of trials and p is the probability of success.

The variance of a binomial distribution of a discrete random variable X is denoted and defined by $\sigma^2 = \text{Var}(X) = npq$ where n is the number of trials, p is the probability of success and q is the probability of failure.

The standard deviation of a binomial distribution of a discrete random variable X is denoted and defined by $\sigma = \sqrt{\text{Var}(X)} = \sqrt{npq}$.

4. Uncountable infinite discrete case: Poisson distribution

The **Poisson distribution** is a discrete distribution often used as a model for the number of events (such as the number of customers in waiting lines, number of defects in a given surface area, airplane arrivals, or the number of accidents at an intersection) in a specific time period.

Poisson distribution is a limiting form of the binomial distribution $(p + q)^n$ under the following **conditions**:

- (i) $n \rightarrow \infty$, i.e., the number of trials is indefinitely large.
- (ii) $p \rightarrow 0$, i.e., the constant probability of success for each trial is indefinitely small.
- (iii) np is a finite quantity, say λ .

Typical events which could have a Poisson distribution:

- (i) Number of customers arriving at a supermarket checkout per minute.
- (ii) Number of suicides or deaths by heart attack in a minute.
- (iii) Number of accidents that take place on a busy road in time t .
- (iv) Number of printing mistakes at each unit of the book.

- (v) Number of cars passing a certain street in time t .
- (vi) Number of α – particles emitted per second by a radioactive sources.
- (vii) Number of faulty blades in a packet of 1000.
- (viii) Number of person born blind per year in a certain village.
- (ix) Number of telephone calls received at a particular switch board in a minute.
- (x) Number of times a teacher is late for class in a given week.

The probability density function of Poisson distribution is defined by

$$P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}, \quad x = 0, 1, 2, \dots$$

where λ is a parameter which indicates the average number (the expected value) of events in the given time interval. We write $X \sim \text{Po}(\lambda)$

- ☛ If the initial probability is known, you can find successive probabilities using the following recurrence relation;

$$P(X = x+1) = \frac{\lambda}{x+1} P(X = x).$$

- ☛ For a Poisson distribution of a discrete random variable X , the mean μ (or expected value) and the variance σ^2 are the same and equal to λ . Thus, $\mu = \sigma^2 = \lambda$

5. Continuous random variables

a) Probability density function

A function defined on an interval $[a, b]$ is a probability density function for a continuous random variable X distributed on $[a, b]$ if, whenever x_1 and x_2 satisfy

$$a \leq x_1 \leq x_2 \leq b, \text{ we have } p(x_1 \leq X \leq x_2) = \int_{x_1}^{x_2} f(x) dx.$$

Properties of p.d.f $f(x)$

a) $f(x) > 0$ for all x

b) $\int_{all\ x} f(x) dx = 1$

The **cumulative distribution** function of a continuous random variable X is defined as: $F(x) = \int_{-\infty}^x f(t) dt$

where $F(x) = 0$, for $x \rightarrow -\infty$ and $x \rightarrow +\infty$ for $x \rightarrow +\infty$.

b) Expected value, variance and standard deviation

The mean μ (or expected value $E(X)$) of X is denoted and defined by

$$\mu = E(X) = \int_a^b xf(x) dx$$

Also, expectation of function g of X is

$$E(g(x)) = \int_a^b g(x)f(x) dx$$

The variance $Var(x)$ or σ^2 is denoted and defined by

$$\sigma^2 = Var(X) = E(X^2) - [E(x)]^2.$$

The standard deviation is

$$\sigma = SD = \sqrt{Var(X)}.$$

Properties of $E(X)$ and $Var(X)$

$$\forall a, b \in \mathbb{R}$$

1. $E(a) = a$
2. $E(aX) = aE(X)$
3. $E(aX + b) = aE(X) + b$
4. $E(X + Y) = E(X) + E(Y)$
5. $E(aX^2 + b) \neq aE(X^2) + b$
6. $var(a) = 0$
7. $var(aX) = a^2 var(X)$
8. $var(aX + b) = a^2 var(X)$

End of Unit Assessment answers Learner's Book page 488

1. $\frac{29}{32}$
2. a) 0.0145 b) 0.1887
c) 0.000000000000000000000001
3. 0.0863
4. Probability that it will work (0 defective components) is 0.896 . Probability that it will not work perfectly is 0.104
5. 0.00038
6. a) $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$
b) $p(HHH) = 0.166375$,
 $p(HHT) = p(HTH) = p(THH) = 0.136125$,
 $p(HTT) = p(THT) = p(TTH) = 0.111375$,
 $p(TTT) = 0.091125$
c) $f(0) = 0.911125$, $f(1) = 0.334125$, $f(2) = 0.408375$,
 $f(3) = 0.166375$
d) 0.908875 e) 1.650000
7. $\frac{176}{1024} \times 100 \approx 17$ 8. 0.99863
9. 0.0376 10. $\frac{na}{a+b}$
11. a) 0.5905 b) $E(X) = 7, Var(x) = 6.3$
12. a) 0.9997 b) 0.005
13. $\frac{3}{4}$ 14. a) 4.8, 0.98 b) 0.655
15. a) 0.0498×1000 b) 0.3526×1000
16. 2.3026 17. $\frac{e^{-100} (100)^x}{x!}$
18. 0.51
19. a) 0.147 b) 0.0408 c) 0.762

20. a) 0.122 b) 0.138 c) 0.224 d) 0.0273

21.

x	0	1	2	3	4	Total
f	122	60	15	2	1	200
$x \cdot f$	0	60	30	6	4	100

$$\text{Mean} = \frac{100}{200} = 0.5$$

The number of x deaths is given by $200 \times \frac{(e)^{-0.5} (0.5)^x}{x!}$ for 0, 1, 2, 3, 4.

Death/Frequencies	0	1	2	3	4
Probabilities $P(x) = \frac{e^{-0.5} \times (0.5)^x}{x!}$	0.27	0.35	0.23	0.10	0
Expected (Theoretical) frequency $200 \times \frac{(e)^{-0.5} (0.5)^x}{x!}$	121	61	15	3	0

The expected frequencies are 121,61,15,3 and 0.

22. 0.9998 23. 0.5620

24.

x	0	1	2	3	4	5	6	Total
f	143	90	42	12	9	3	1	300
$x \cdot f$	0	90	84	36	36	15	6	267

$$\text{Mean} = \frac{267}{300} = 0.89$$

The number of x mistakes per day is given by

$$300 \times \frac{(e)^{-0.89} (0.89)^x}{x!} \text{ for } 0, 1, 2, 3, 4, 5, 6.$$

Mistakes per day	1	1	2	3	4	5	6
Probabilities $P(x) = \frac{e^{-0.5} \times (0.5)^x}{x!}$	0.41	0.37	0.16	0.05	0.01	0	0
Expected (Theoretical) frequency $200 \times \frac{(e)^{-0.5} (0.5)^x}{x!}$	123	111	48	15	3	0	0

The expected frequencies are 123, 111, 48, 15, 3 and 0.

25. a) 0.25 b) $\frac{2}{3}$
26. a) $k = \frac{1}{2}$ b) 0.75
27. a) 0.125 b) 0.727
28. a) $k = \frac{3}{16}$ b) $E(x) = 1\frac{3}{4}$
- c) $P(1 \leq x \leq 3) = \frac{11}{16}$
29. a) (i) $k = \frac{2}{5}$ (ii) $E(x) = 1\frac{4}{15}$
- (iii) $\sigma = 0.75$ b) $p(x < \mu - \sigma) = 0.207$
30. (i) a) $k = \frac{2}{9}$ b) $\mu = E(x) = 2, \sigma^2 = \frac{1}{2}, \sigma = \frac{\sqrt{2}}{2}$
- c) $\frac{4\sqrt{2}}{9} \approx 0.63$
- (ii) a) $k = 3$ b) $\mu = E(x) = \frac{3}{4}, \sigma^2 = \frac{3}{80}, \sigma = \frac{3\sqrt{5}}{20}$
- c) $\frac{207\sqrt{5}}{400} \approx 0.668$
- (iii) a) $k = 6$ b) $\mu = E(x) = \frac{1}{2}, \sigma^2 = \frac{1}{20}, \sigma = \frac{\sqrt{5}}{20}$
- c) $\frac{7\sqrt{5}}{25} \approx 0.626$
31. a) $a = 12, b = 1$ b) 0.0523

Answers for Summative Evaluation One

Learner's Book pages 494 - 496

$$1. \left(\frac{1}{8}\right)^{x-2} = 4^{3-2x} \Leftrightarrow (2^{-3})^{x-2} = (2^2)^{3-2x}$$

$$\Leftrightarrow 2^{-3x+6} = 2^{6-4x} \Rightarrow -3x+6 = 6-4x$$

or $x = 0$

$$S = \{0\}$$

2. A quadratic function has a double root if and only if $\Delta = 0$.

For our case, $\Delta = b^2 - 4ac = 9 - 4m$.

$$\Delta = 0 \Leftrightarrow 9 - 4m = 0 \text{ or } m = \frac{9}{4}.$$

Therefore, $x^2 + 3x + m = 0$ admits a double root when

$$m = \frac{9}{4}.$$

For $m = \frac{9}{4}$, $x^2 + 3x + m = 0 \Rightarrow x^2 + 3x + \frac{9}{4} = 0$

The root is $x = -\frac{3}{2}$.

3. If the angle between $\vec{u} = (k, 3)$ and $\vec{v} = (4, 0)$ is 45° , thus

$$\cos 45^\circ = \frac{4k}{\sqrt{k^2 + 9}\sqrt{16}}$$

Or

$$\cos 45^\circ = \frac{4k}{4\sqrt{k^2 + 9}} \Leftrightarrow \cos 45^\circ = \frac{k}{\sqrt{k^2 + 9}}$$

Since $\cos 45^\circ = \frac{\sqrt{2}}{2}$, then,

$$\frac{\sqrt{2}}{2} = \frac{k}{\sqrt{k^2 + 9}} \Leftrightarrow 2k = \sqrt{2}\sqrt{k^2 + 9}$$

Squaring both sides yields

$$4k^2 = k^2 + 18 \Leftrightarrow k^2 = 9 \Rightarrow k = \pm 3$$

The value of k is 3 since $\cos 45^\circ > 0$.

$$\begin{aligned}
 4. \quad & 2 \cos^2 x - \cos x - 1 = 0 \Leftrightarrow 2 \cos^2 x - 2 \cos x + \cos x - 1 = 0 \\
 & \Leftrightarrow 2 \cos x (\cos x - 1) + \cos x - 1 = 0 \\
 & \Leftrightarrow (\cos x - 1)(2 \cos x + 1) = 0 \quad \cos x - 1 \\
 & \Rightarrow \cos x - 1 = 0 \quad \text{or} \quad 2 \cos x + 1 \\
 & \Leftrightarrow \cos x = 1 \quad \text{or} \quad \cos x = -\frac{1}{2} \\
 & \Rightarrow x = 2k\pi \quad \text{or} \quad x = \pm \frac{2\pi}{3} + 2k\pi, k \in \mathbb{Z} \\
 & \Rightarrow x = 2k\pi \quad \text{or} \quad x = \frac{2\pi}{3} + 2k\pi \quad \text{or} \quad x = -\frac{2\pi}{3} + 2k\pi \equiv \frac{4\pi}{3} + 2k\pi \\
 & \text{Hence, } S = \left\{ 2k\pi, \frac{2\pi}{3} + 2k\pi, \frac{4\pi}{3} + 2k\pi, k \in \mathbb{Z} \right\}.
 \end{aligned}$$

$$5. \quad \lim_{x \rightarrow 0} \frac{1 - \cos x}{\sin x} = \frac{1 - \cos 0}{\sin 0} = \frac{1 - 1}{0} = \frac{0}{0} \text{ I.F.}$$

Remove this indeterminate form by Hospital's rule

$$\lim_{x \rightarrow 0} \frac{1 - \cos x}{\sin x} = \lim_{x \rightarrow 0} \frac{(1 - \cos x)'}{(\sin x)'} = \lim_{x \rightarrow 0} \frac{\sin x}{\cos x} = 0$$

$$\text{Then, } \lim_{x \rightarrow 0} \frac{1 - \cos x}{\sin x} = 0$$

6. From sequence $\{u_n\}$ where $u_{n+1} = 3(u_n + 2)$ and $u_0 = 0$, we list;

$$u_0 = 0$$

$$u_1 = 3(u_0 + 2) = 6$$

$$u_2 = 3(u_1 + 2) = 3(6 + 2) = 24$$

$$u_3 = 3(u_2 + 2) = 3(24 + 2) = 78$$

$$u_4 = 3(u_3 + 2) = 3(78 + 2) = 240.$$

Therefore, the first five terms of the given sequence are 0, 6, 24, 78 and 240.

The sequence $\{u_n\}$ is arithmetic if $u_{n+1} - u_n = d, d \in \mathbb{R}$

and is geometric if $\frac{u_{n+1}}{u_n} = r, r \in \mathbb{R}$

Since $u_{n+1} = 3(u_n + 2)$, thus

$u_{n+1} - u_n = 3(u_n + 2) - u_n = 2u_n + 6$ and this is not a constant.

So, $\{u_n\}$ is not an arithmetic sequence.

$\frac{u_{n+1}}{u_n} = \frac{3(u_n + 2)}{u_n} = 3 + \frac{6}{u_n}$ and this is not a constant.

Thus, $\{u_n\}$ is not a geometric sequence.

Therefore, $\{u_n\}$ is neither arithmetic nor geometric sequence.

7. Let $f(x) = \sqrt{x^2 + 2\sqrt{x^2 - 1}} - \sqrt{x^2 - 2\sqrt{x^2 - 1}}$

a) Existence condition: $x^2 - 1 \geq 0$ and $x^2 - 2\sqrt{x^2 - 1} \geq 0$

$$x^2 - 1 \geq 0 \Leftrightarrow x \in (-\infty, -1] \cup [1, +\infty)$$

$$x^2 - 2\sqrt{x^2 - 1} \geq 0 \Leftrightarrow x^2 \geq 2\sqrt{x^2 - 1}$$

$$\Leftrightarrow x^4 \geq 4x^2 - 4 \Leftrightarrow x^4 - 4x^2 + 4 \geq 0 \Leftrightarrow (x^2 - 2)^2 \geq 0$$

$$\Leftrightarrow \forall x \in \mathbb{R}, (x^2 - 2)^2 \geq 0$$

Hence,

$$\text{Dom}f = (-\infty, -1] \cup [1, +\infty)$$

b) $f(x) = \sqrt{x^2 + 2\sqrt{x^2 - 1}} - \sqrt{x^2 - 2\sqrt{x^2 - 1}}$

$$f^2(x) = (x^2 + 2\sqrt{x^2 - 1}) - 2\sqrt{x^2 + 2\sqrt{x^2 - 1}}\sqrt{x^2 - 2\sqrt{x^2 - 1}} + (x^2 - 2\sqrt{x^2 - 1})$$

$$f^2(x) = 2x^2 - 2\sqrt{x^4 - 4x^2 + 4}$$

$$f^2(x) = 2x^2 - 2\sqrt{(x^2 - 2)^2}$$

$$f^2(x) = 2x^2 - 2|x^2 - 2|$$

$$f^2(x) = \begin{cases} 2x^2 - 2(x^2 - 2), & x \in]-\infty, -\sqrt{2}] \cup [\sqrt{2}, +\infty[\\ 2x^2 + 2(x^2 - 2), & x \in]-\sqrt{2}, -1] \cup [1, \sqrt{2}[\end{cases}$$

$$f^2(x) = \begin{cases} 4, & x \in]-\infty, -\sqrt{2}] \cup [\sqrt{2}, +\infty[\\ 4x^2 - 4, & x \in]-\sqrt{2}, -1] \cup [1, \sqrt{2}[\end{cases}$$

$$\Rightarrow f(x) = \begin{cases} 2, & x \in]-\infty, -\sqrt{2}] \cup [\sqrt{2}, +\infty[\\ 2\sqrt{x^2 - 1}, & x \in]-\sqrt{2}, -1] \cup [1, \sqrt{2}[\end{cases}$$

From $Domf = (-\infty, -1] \cup [1, +\infty)$, we get that

$$\Rightarrow f(x) = \begin{cases} 2, & x \in]-\infty, -\sqrt{2}] \cup [\sqrt{2}, +\infty[\\ 2\sqrt{x^2 - 1}, & x \in]-\sqrt{2}, -1] \cup [1, \sqrt{2}[\end{cases}$$

8. Let $P(T) = p$, then $P(H) = 3p$.

But $P(T) + P(H) = 1$.

Therefore, $4p = 1$ or $p = \frac{1}{4}$.

Thus, $P(T) = \frac{1}{4}$ and $P(H) = \frac{3}{4}$.

9. Tangent line:

$$T \equiv y - y_0 = y'_0(x - x_0)$$

Here, $x_0 = 2$ and $y_0 = 4$;

$$f'(x) = 3x^2 - 4x$$

$$y'_0 = f'(2) = 3(4) - 4(2) = 12 - 8 = 4$$

$$\text{Then, } T \equiv y - 4 = 4(x - 2) \Leftrightarrow y = 4x - 8 + 4 \Leftrightarrow y = 4x - 4$$

Normal line:

$$N \equiv y - y_0 = -\frac{1}{y'_0}(x - x_0);$$

$$\text{Thus, } N \equiv y - 4 = -\frac{1}{4}(x - 2) \Leftrightarrow y = 4x + \frac{1}{2} + 4$$

$$\Leftrightarrow y = 4x + \frac{9}{2}$$

10.a) Equation of sphere S whose centre (x_0, y_0, z_0) and

radius r has equation $(x - x_0)^2 + (y - y_0)^2 + (z - z_0)^2 = r^2$

For our case,

$$S \equiv (x-6)^2 + (y-5)^2 + (z+2)^2 = 70$$

$$\text{Or } S \equiv x^2 + y^2 + z^2 - 12x - 10y + 4z = 70 - 36 - 25 - 4$$

$$\text{Or } S \equiv x^2 + y^2 + z^2 - 12x - 10y + 4z = 5$$

$$\text{b) } x^2 + y^2 + z^2 + 4x - 8y + 6z + 7 = 0 \Leftrightarrow x^2 + 4x + y^2 - 8y + z^2 + 6z = -7$$

$$\Leftrightarrow (x+2)^2 + (y-4)^2 + (z+3)^2 = -6 + 4 + 16 + 9$$

$$\Leftrightarrow (x+2)^2 + (y-4)^2 + (z+3)^2 = 22$$

Centre is $C(-2, 4, -3)$, radius $r = \sqrt{22}$.

$$\text{c) Let us find } \overrightarrow{AB} = \begin{pmatrix} 1 \\ -4 \\ 5 \end{pmatrix}, \text{ the line } AB \equiv \frac{x-1}{1} = \frac{y-1}{-4} = \frac{z+1}{5}$$

$$\Rightarrow AB \equiv \begin{cases} -4x + 4 = y + 1 \\ 5x - 5 = z + 1 \end{cases} \Leftrightarrow AB \equiv \begin{cases} y = -4x + 3 \\ z = 5x - 6 \end{cases}$$

Substituting y, z with their values in S gives

$$x^2 + (-4x+3)^2 + (5x-6)^2 + 4x - 8(-4x+3) + 6(5x-6) + 7 = 0$$

$$\Leftrightarrow x^2 + 16x^2 - 24x + 9 + 25x^2 - 60x + 36 + 4x + 32x - 24 + 30x - 36 + 7 = 0$$

$$\Leftrightarrow 42x^2 - 18x - 8 = 0 \Leftrightarrow 21x - 9x - 4 = 0$$

$$\Delta = 81 + 336 = 417$$

$$x_{1,2} = \frac{9 \pm \sqrt{417}}{42}$$

$$x_1 = 5.7 \text{ and } x_2 = -14.7$$

For $x = 5.7$, we have $y = -19.8$ and $z = 22.5$.

Intersection point is then, $(5.7, -19.8, 22.5)$

For $x = -14.7$, we have $y = 61.8$ and $z = 79.5$

Intersection point is then, $(-14.7, 61.8, 79.5)$

$$11. \text{ Let } q(t) = q_0 e^{-tk}$$

Here $q_0 = 50$ and $q(5) = 20$.

$$q(5) = 20 \Rightarrow 20 = 50e^{-5k}$$

$$\Leftrightarrow \frac{2}{5} = e^{-5k} \Leftrightarrow -5k = \ln \frac{2}{5} \Leftrightarrow k = -\frac{1}{5} \ln \frac{2}{5} \Leftrightarrow k = 0.18326.$$

90% of the sugar being dissolved, it means that 10% of the sugar left i.e. 5kg.

Thus,

$$q(t) = 5 \Rightarrow 50e^{-0.18326t} = 5 \Leftrightarrow e^{-0.18326t} = \frac{1}{10}$$

$$\Leftrightarrow -0.18326t = \ln \frac{1}{10} \Leftrightarrow 0.18326t = -\ln 10 \Leftrightarrow t = 12.5647$$

12. In fact, $\sin y \cos(x-y) + \cos y \sin(x-y)$

$$= \sin y (\cos x \cos y + \sin x \sin y) + \cos y (\sin x \cos y - \sin y \cos x)$$

$$= \cancel{\sin y \cos x \cos y} + \sin x \sin^2 y + \sin x \cos^2 y - \cancel{\cos y \sin y \cos x}$$

$$= \sin x \sin^2 y + \sin x \cos^2 y = \sin x (\sin^2 y + \cos^2 y)$$

$$= \sin x \text{ as required.}$$

13. $\lim_{x \rightarrow 0} \frac{\sqrt[5]{(1+x)^3 - 1}}{x} = \frac{0}{0}$, *I.F.*

Remove this *I.F.* by Hospital's rule.

$$\lim_{x \rightarrow 0} \frac{\sqrt[5]{(1+x)^3 - 1}}{x} = \lim_{x \rightarrow 0} \frac{(\sqrt[5]{(1+x)^3 - 1})'}{x'} = \lim_{x \rightarrow 0} \frac{1}{5} [(1+x)^3 - 1]^{\frac{4}{5}} 3(1+x)^2$$

$$= \frac{1}{5} [(1+0)^3 - 1]^{\frac{4}{5}} 3(1+0)^2 = +\infty$$

14. If the mean is 50, thus, $\frac{56 + 37 + 54 + 52 + x + 48}{6} = 50$

$$\Leftrightarrow 247 + x = 300 \Leftrightarrow x = 300 - 247 \Rightarrow x = 53$$

15. Intersection points for $y^2 = 2px$ and $x^2 = 2py$:

$$y^2 = 2px \Rightarrow y = \sqrt{2px} \text{ and } x^2 = 2py \Rightarrow y = \frac{x^2}{2p}$$

$$\text{Then, } \sqrt{2px} = \frac{x^2}{2p} \Leftrightarrow 2px = \frac{x^4}{4p^2}$$


$$\Leftrightarrow 8p^3x = x^4 \Leftrightarrow 8p^3x - x^4 = 0 \Leftrightarrow x(8p^3 - x^3) = 0$$

$$\Rightarrow x = 0 \text{ or } 8p^3 - x^3 = 0$$

$$\Rightarrow x = 0 \text{ or } x = 2p$$

To be able to sketch the curve, let $p = 2$. Then we have

$$y^2 = 4x \text{ and } x^2 = 4y$$



$$A = \int_0^{2p} \left(\sqrt{2px} - \frac{x^2}{2p} \right) dx$$

$$A = \sqrt{2p} \int_0^{2p} \sqrt{x} dx - \frac{1}{2p} \int_0^{2p} x^2 dx$$

$$A = \sqrt{2p} \left[\frac{2}{3} x^{\frac{3}{2}} \right]_0^{2p} - \frac{1}{2p} \left[\frac{x^3}{3} \right]_0^{2p}$$

$$A = \frac{2}{3} \sqrt{2p} \cdot 2p \cdot \sqrt{2p} - \frac{8p^3}{6p}$$

$$A = \frac{8p^2}{3} - \frac{4p^3}{3p} = \frac{4p^3}{3p} = \frac{4p^2}{3}$$

Therefore, the area enclosed by the curves

$y^2 = 2px$ and $x^2 = 2py$ is $\frac{4p^2}{3}$ sq. unit

16. a) $r = \frac{\text{Cov}(x, y)}{\sigma_x \sigma_y}$

$$\text{Cov}(x, y) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})$$

$$\sigma_x = \sqrt{\text{Var}(x)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}$$

$$\sigma_y = \sqrt{\text{Var}(y)} = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2}$$

Mean:

$$\bar{x} = \frac{7+8+9+11+15}{5} = 10$$

$$\bar{y} = \frac{33+25+17+9+6}{5} = 18$$

x_i	y_i	$x_i - \bar{x}$	$y_i - \bar{y}$	$(x_i - \bar{x})^2$	$(y_i - \bar{y})^2$	$(x_i - \bar{x})(y_i - \bar{y})$
7	33	-3	-12	9	225	-45
8	25	-2	-9	4	49	-14
9	17	-1	-1	1	1	1
11	9	1	7	1	81	-9
15	6	5	15	25	144	-60
SUM				40	500	-127

$$Cov(x, y) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 (y_i - \bar{y})^2$$

$$\sigma_x = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2} = \sqrt{\frac{40}{5}} = \sqrt{8} = 2\sqrt{2}$$

$$\sigma_y = \sqrt{\frac{1}{n} \sum_{i=1}^n (y_i - \bar{y})^2} = \sqrt{\frac{500}{5}} = \sqrt{100} = 10$$

$$Cov(x, y) = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2 (y_i - \bar{y})^2 = \frac{-127}{5}$$


$$r = \frac{Cov(x, y)}{\sigma_x \sigma_y} = \frac{-127}{5(2\sqrt{2})10} = -0.89$$

b) $L_{y/x} \equiv y - \bar{y} = \frac{Cov(x, y)}{\sigma_x^2} (x - \bar{x})$

$$L_{y/x} \equiv y - 18 = \frac{-127}{40} (x - 10)$$

$$\Leftrightarrow y - 18 = -3.2(x - 10) \Leftrightarrow y = -3.2x + 50$$

c) Scatter diagram



$$17.a) \begin{cases} 3x + 2y - 5z = 2 \\ x + 2y = 3 \\ 2x - y + z = -3 \end{cases} \Leftrightarrow \begin{pmatrix} 3 & 2 & -5 \\ 1 & 2 & 0 \\ 2 & -1 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 2 \\ 3 \\ -3 \end{pmatrix}$$

$$\text{If } \Delta \neq 0, \text{ then } x = \frac{\Delta_y}{\Delta}, y = \frac{\Delta_x}{\Delta}, z = \frac{\Delta_z}{\Delta}$$

$$\Delta = \begin{vmatrix} 3 & 2 & -5 \\ 1 & 2 & 0 \\ 2 & -1 & 1 \end{vmatrix} = - \begin{vmatrix} 2 & -5 \\ -1 & 1 \end{vmatrix} + 2 \begin{vmatrix} 3 & -5 \\ 2 & 1 \end{vmatrix} = -2 + 5 + 6 + 20 = 29$$

$$\Delta_x = \begin{vmatrix} 2 & 2 & -5 \\ 3 & 2 & 0 \\ -3 & -1 & 1 \end{vmatrix} = -3 \begin{vmatrix} 2 & -5 \\ -1 & 1 \end{vmatrix} + 2 \begin{vmatrix} 2 & -5 \\ -3 & 1 \end{vmatrix} = -6 + 15 + 4 - 30 = -17$$

$$\Rightarrow x = -\frac{17}{29}$$

$$\Delta_y = \begin{vmatrix} 3 & 2 & -5 \\ 1 & 3 & 0 \\ 2 & -3 & 1 \end{vmatrix} = - \begin{vmatrix} 2 & -5 \\ -3 & 1 \end{vmatrix} + 3 \begin{vmatrix} 3 & -5 \\ 2 & 1 \end{vmatrix} = -2 + 15 + 9 + 30 = 52$$

$$\Rightarrow y = \frac{52}{29}$$

$$\Delta_z = \begin{vmatrix} 3 & 2 & 2 \\ 1 & 2 & 3 \\ 2 & -1 & -3 \end{vmatrix} = - \begin{vmatrix} 2 & 2 \\ -1 & -3 \end{vmatrix} + 2 \begin{vmatrix} 3 & 2 \\ 2 & -3 \end{vmatrix} - 3 \begin{vmatrix} 3 & 2 \\ 2 & -1 \end{vmatrix}$$

$$= 6 - 2 - 18 - 8 + 9 + 12 = -1 \Rightarrow z = -\frac{1}{29}$$

$$S = \left\{ \left(-\frac{17}{29}, \frac{52}{29}, -\frac{1}{29} \right) \right\}$$

- b) The area of a parallelogram whose adjacent sides are $\vec{a} = 6\vec{i} + 3\vec{j} - 2\vec{k}$ and $\vec{b} = 3\vec{i} - 2\vec{j} + 6\vec{k}$ is given by

$$A = \|\vec{a} \times \vec{b}\|$$

$$\text{Or } \vec{a} \times \vec{b} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 6 & 3 & -2 \\ 3 & -2 & 6 \end{vmatrix} = \vec{i} \begin{vmatrix} 3 & -2 \\ -2 & 6 \end{vmatrix} - \vec{j} \begin{vmatrix} 6 & -2 \\ 3 & 6 \end{vmatrix} + \vec{k} \begin{vmatrix} 6 & 3 \\ 3 & -2 \end{vmatrix}$$

$$= \vec{i}(18 - 4) - \vec{j}(36 + 6) + \vec{k}(-12 - 9) = 14\vec{i} - 42\vec{j} - 21\vec{k}$$

$$A = \sqrt{(14)^2 + (-42)^2 + (-21)^2} = \sqrt{196 + 1764 + 441}$$

$$= \sqrt{2401} = 49$$

Therefore, the area of the given parallelogram is *49 sq. unit*

18. We are given $F(x) = \begin{cases} cx(6-x)^2; & 0 \leq x \leq 6 \\ 0, & \text{elsewhere} \end{cases}$

- a) Since x is a random variable, thus, $\int_{All} f(x) dx = 1$

$$\Rightarrow 1 = \int_0^6 cx(6-x)^2 dx \Leftrightarrow 1 = \int_0^6 cx(36 - 12x + x^2) dx$$

$$\Leftrightarrow 1 = c \left[18x^2 - 4x^3 + \frac{x^4}{4} \right]_0^6 \Leftrightarrow 1 = c[18(36) - 4(216) + 324]$$

$$\Leftrightarrow 108c = 1$$

$$\Rightarrow c = \frac{1}{108}$$

- b) i) The mean is $E(x) = \int_{All} xf(x) dx$

$$\Rightarrow E(x) = \frac{1}{108} \int_0^6 x^2 (6-x)^2 dx$$

$$\begin{aligned} \Leftrightarrow E(x) &= \frac{1}{108} \int_0^6 (36x^2 - 12x^3 + x^4) dx \\ \Leftrightarrow E(x) &= \frac{1}{108} \left[12x^3 - 3x^4 + \frac{x^5}{5} \right]_0^6 \\ \Leftrightarrow E(x) &= \frac{1}{108} (2592 - 3888 + 1555.2) = 2.4 \\ \Leftrightarrow E(x) &= \frac{1}{108} (259.2) = 2.4 \\ E(x) &= 2.4 \end{aligned}$$

ii) The variance $Var(x) = E(x^2) - [E(x)]^2$
 or $E(x^2) = \int_{All} x^2 f(x) dx \Rightarrow E(x^2) = \frac{1}{108} \int_0^6 x^3 (6-x)^2 dx$

$$\begin{aligned} \Leftrightarrow E(x^2) &= \frac{1}{108} \int_0^6 (36x^3 - 12x^4 + x^5) dx \\ \Leftrightarrow E(x^2) &= \frac{1}{108} \left[9x^4 - 12 \frac{x^5}{5} + \frac{x^6}{6} \right]_0^6 \\ \Leftrightarrow E(x^2) &= \frac{1}{108} \left[9(6)^4 - 12 \frac{(6)^5}{5} + \frac{(6)^6}{6} \right]_0^6 = 7.2 \end{aligned}$$

Then, $Var(x) = 7.2 - (2.4)^2 = 1.44$

iii) Standard deviation of x is $\sigma = \sqrt{Var(x)} = \sqrt{1.44} = 1.2$

19. a) From $I_n = \int_0^{\frac{\pi}{2}} e^{-nx} \sin x dx$, let $u = \sin x$ and $dv = e^{-nx} dx$.

Hence, $du = \cos x dx$ and $v = -\frac{1}{n} e^{-nx}$.

Therefore, $I_n = \left[-\frac{1}{n} e^{-nx} \sin x \right]_0^{\frac{\pi}{2}} + \frac{1}{n} \int_0^{\frac{\pi}{2}} e^{-nx} \cos x dx$

$$\Leftrightarrow I_n = -\frac{1}{n} e^{-\frac{n\pi}{2}} + \frac{1}{n} J_n$$

$$\Leftrightarrow nI_n - J_n = -e^{-\frac{n\pi}{2}} \quad (1)$$

From $J_n = \int_0^{\frac{\pi}{2}} e^{-nx} \cos x dx$, let $t = \cos x$ and $dz = e^{-nx} dx$.

Then, $dt = -\sin x dx$ and $z = -\frac{1}{n} e^{-nx}$.

Therefore, $J_n = \left[-\frac{1}{n} e^{-nx} \cos x \right]_0^{\frac{\pi}{2}} - \frac{1}{n} \int_0^{\frac{\pi}{2}} e^{-nx} \sin x dx$

$$\Leftrightarrow J_n = \frac{1}{n} - \frac{1}{n} I_n$$

$$\Leftrightarrow nJ_n + I_n = 1 \quad (2)$$

Equation (1) and (2) give the simultaneous equations

$$\begin{cases} nI_n - J_n = -e^{-\frac{n\pi}{2}} \\ nJ_n + I_n = 1 \end{cases} \quad (3)$$

And (3) indicates two relations between I_n and J_n .

b) Multiply first equation of (3) by n to eliminate J_n

$$\begin{cases} n^2 I_n - nJ_n = -ne^{-\frac{n\pi}{2}} \\ nJ_n + I_n = 1 \end{cases} \Rightarrow n^2 I_n + I_n = 1 - ne^{-\frac{n\pi}{2}}$$

$$\text{Which gives } I_n = \frac{1 - ne^{-\frac{n\pi}{2}}}{n^2 + 1} \quad (4)$$

$$\text{From (1), } J_n = nI_n + e^{-\frac{n\pi}{2}}$$

$$\text{Then } J_n = \frac{n - n^2 e^{-\frac{n\pi}{2}} + n^2 e^{-\frac{n\pi}{2}} + e^{-\frac{n\pi}{2}}}{n^2 + 1}$$

$$\text{Or } J_n = \frac{n + e^{-\frac{n\pi}{2}}}{n^2 + 1} \quad (5)$$

20. To solve $y'' - y' - 2y = 6x$ with $y(0) = y'(0) = 1$

Homogeneous equation :

$$y'' - y' - 2y = 0$$

Characteristic equation

$$\lambda^2 - \lambda - 2 = 0 \Leftrightarrow (\lambda + 1)(\lambda - 2) = 0$$

$$\Leftrightarrow \lambda = -1 \text{ or } \lambda = 2.$$

General solution for solution for homogeneous equation is

$$y^* = c_1 e^{-x} + c_2 e^{2x}.$$

The complementary (particular) solution is given by

$$y = Ax + B.$$

Or $y' = A$ et $y'' = 0$.

The equation $y'' - y' - 2y = 6x$ becomes

$$-A - 2Ax - 2B = 6x$$

Identifying the coefficients, we get

$$\begin{cases} -A - 2B = 0 \\ -2A = 6 \end{cases} \Leftrightarrow \begin{cases} B = -\frac{A}{2} \\ A = -3 \end{cases} \Leftrightarrow \begin{cases} B = \frac{3}{2} \\ A = -3 \end{cases}$$

Thus, complementary solution is $y = -3x + \frac{3}{2}$

The general solution of the given equation is

$$y = c_1 e^{-x} + c_2 e^{2x} - 3x + \frac{3}{2}$$

From the initial conditions $y(0) = y'(0) = 1$, we get the values of c_1 and c_2 as follows:

$$y' = -c_1 e^{-x} + 2c_2 e^{2x} - 3$$

$$y(0) = y'(0) = 1 \Rightarrow \begin{cases} c_1 + c_2 + \frac{3}{2} = 1 \\ -c_1 + 2c_2 - 3 = 1 \end{cases} \Rightarrow \begin{cases} c_1 = -\frac{5}{3} \\ c_2 = \frac{7}{6} \end{cases}$$

Therefore, the required solution is

$$y = -\frac{5}{3} e^{-x} + \frac{7}{6} e^{2x} - 3x + \frac{3}{2}$$

Answers for Summative Evaluation Two

Learner's Book pages 497 - 499

$$1. \quad \begin{cases} x^2 + y^2 = \frac{37}{4} \\ xy = \frac{3}{2} \end{cases}$$

From 2nd equation, we get $x = \frac{3}{2y}$; putting this equality in 1st equation, we get

$$\left(\frac{3}{2y}\right)^2 + y^2 = \frac{37}{4} \Leftrightarrow \frac{9}{4y^2} + y^2 = \frac{37}{4}$$

$$\Leftrightarrow 9 + 4y^4 = 37y^2 \Rightarrow 4y^4 - 37y^2 + 9 = 0$$

$$\Delta = (-37)^2 - 16(9) = 1369 - 144 = 1225$$

$$y^2 = \frac{37+35}{8} = 9 \text{ or } y^2 = \frac{37-35}{8} = \frac{1}{4}$$

Solving for y , we get: $y_1 = -3, y_2 = 3, y_3 = -\frac{1}{2}, y_4 = \frac{1}{2}$

Substituting y with its values in $x = \frac{3}{2y}$, we get:

$$x_1 = -\frac{1}{2}, x_2 = \frac{1}{2}, x_3 = -3, x_4 = 3$$

And then, the solution set is

$$S = \left\{ \left(-3, -\frac{1}{2}\right), \left(3, \frac{1}{2}\right), \left(-\frac{1}{2}, -3\right), \left(\frac{1}{2}, 3\right) \right\}$$

$$2. \quad \ln\left(\frac{e^{\ln x}}{e^3}\right) + \ln\left(\frac{e}{x}\right) = \ln\left(\frac{e^{\ln x}}{e^3} \times \frac{e}{x}\right)$$

$$= \ln \frac{e^{\ln x}}{xe^2} = \ln x \ln e - \ln xe^2$$

$$= \ln x - (\ln x + \ln e^2) = \ln x - \ln x + 2 \ln e = 2$$

$$3. \quad \arctan x + \arctan \sqrt{3} = \frac{\pi}{4}$$

$$\Rightarrow \arctan x + \frac{\pi}{3} = \frac{\pi}{4} \Rightarrow \arctan x = -\frac{\pi}{12} \Rightarrow x = \tan\left(-\frac{\pi}{12}\right) \Rightarrow x = -\tan \frac{\pi}{12}$$

$$4. \quad f(x) = \frac{\ln(1+x^2)}{e^{x^2}} \Rightarrow f'(x) = \frac{[\ln(1+x^2)]' \times e^{x^2} - (e^{x^2})' \times \ln(1+x^2)}{e^{x^2}} \\ \Rightarrow f'(x) = \frac{\frac{2x}{1+x^2} \times e^{x^2} - 2xe^{x^2} \ln(1+x^2)}{[e^{x^2}]^2} = \frac{2x[1 - (1+x^2)\ln(1+x^2)]}{(1+x^2)e^{x^2}}$$

$$5. \quad \lim_{x \rightarrow 1} \frac{\sqrt{x}-1}{\sqrt[3]{x}-1} = \frac{0}{0} \quad (I.C) \\ \lim_{x \rightarrow 1} \frac{\sqrt{x}-1}{\sqrt[3]{x}-1} = \lim_{x \rightarrow 1} \frac{(\sqrt{x}-1)(\sqrt{x}+1)(\sqrt[3]{x^2} + \sqrt[3]{x} + 1)}{(\sqrt[3]{x}-1)(\sqrt{x}+1)(\sqrt[3]{x^2} + \sqrt[3]{x} + 1)} \\ = \lim_{x \rightarrow 1} \frac{(x-1)(\sqrt[3]{x^2} + \sqrt[3]{x} + 1)}{(x-1)(\sqrt{x}+1)} = \lim_{x \rightarrow 1} \frac{\sqrt[3]{x^2} + \sqrt[3]{x} + 1}{\sqrt{x} + 1} = \frac{3}{2}$$

$$6. \quad iz - 2 = 4i - z \rightarrow (1)$$

$$\text{Let } z = a + bi \rightarrow (2)$$

Using (2) in (1) we get:

$$i(a + bi) - 2 = 4i - (a + bi) \Rightarrow ai + bi^2 - 2 = 4i - a - bi$$

$$\Rightarrow (-b - 2) + ai = -a + (4 - b)i \Rightarrow \begin{cases} -b - 2 = -a \\ a = 4 - b \end{cases} \Rightarrow \begin{cases} a - b = 2 \\ a + b = 4 \end{cases} \Rightarrow \begin{cases} a = 3 \\ b = 1 \end{cases}$$

$$z = 3 + 1i$$

$$7. \quad \frac{\sin 2x + \sin 2x}{1 + \cos x + \cos 2x} = \frac{\sin x + 2 \sin x \cos x}{1 + \cos x + \cos^2 x - \sin^2 x} \\ = \frac{\sin x + 2 \sin x \cos x}{\cos x + 2 \cos^2 x} = \frac{\sin x(1 + 2 \cos x)}{\cos x(1 + 2 \cos x)} = \frac{\sin x}{\cos x} = \tan x$$

$$8. \quad \lim_{n \rightarrow +\infty} \frac{1 + 2 + 3 + 4 + \dots + n}{n^2} = \lim_{n \rightarrow +\infty} \frac{n(n+1)}{2n^2} = \lim_{n \rightarrow +\infty} \frac{n+1}{2n} = \frac{1}{2}$$

$$9. \quad y = \ln(4x - 11), \quad x_0 = 3$$

$$T \equiv y - y_0 = y'(x - x_0)$$

$$\text{where: } \begin{cases} y_0 = y(x_0) = y(3) \\ y'_0 = y'(x_0) = y'(3) \end{cases}$$

or

$$y = \ln(4x - 11) \Rightarrow y' = \frac{4}{4x - 11}$$

$$y(3) = 0, y'(3) = 4$$

Then, $T \equiv y = 4(x-3) \Rightarrow T \equiv y = 4x - 12$

10. $f(x) = \ln \frac{x+1}{x-1}$, f is defined if: $\frac{x+1}{x-1} > 0$

x	$-\infty$	-1	1	$+\infty$
$x+1$	-	0	+	+
$x-1$	-		-	0
$\frac{x+1}{x-1}$	+	0	-	+

$Domf =]-\infty, -1[\cup]1, +\infty[$

11. $F(x) = \frac{x^2}{2} + x - x \ln x \Rightarrow F'(x) = x + 1 - \left(\ln x + x \cdot \frac{1}{x} \right) = x - \ln x$

12. a) $e^x e^{x-1} = e \Rightarrow e^{2x-1} = e^1 \Rightarrow 2x-1=1 \Rightarrow 2x=2 \Rightarrow x=1$

$S = \{1\}$

b) $e^{2x-2} + e^{x-2} = 6e^{-2}$

$\Rightarrow e^{2x} e^{-2} + e^x e^{-2} = 6e^{-2} \Rightarrow e^{-2} (e^{2x} + e^x) = 6e^{-2} \Rightarrow e^{2x} + e^x - 6 = 0$

Let $t = e^x, (t > 0)$

$\Rightarrow t^2 + t - 6 = 0 = (t-2)(t+3) \Rightarrow t = 2 \text{ or } t = -3$

$t = -3$ is to be rejected since $t > 0$

For $t = 2 \Rightarrow e^x = 2 \Rightarrow \ln e^x = \ln 2 \Rightarrow x = \ln 2$

$S = \{\ln 2\}$

13. $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$

Let: $f^{-1}(x) = u(x)$ be the inverse of $f(x)$

$f(u) = x$

$\Rightarrow \frac{e^u - e^{-u}}{e^u + e^{-u}} = x$

$\Rightarrow \frac{e^{2u} - 1}{e^u} = \frac{xe^{2u} + x}{e^u}$

$\Rightarrow e^u - e^{-u} = x(e^u + e^{-u})$

$\Rightarrow e^{2u} - 1 = xe^{2u} + x$

$\Rightarrow e^u - \frac{1}{e^u} = xe^u + \frac{x}{e^u}$

$\Rightarrow e^{2u} (1-x) = x+1$


$\Rightarrow e^{2u} = \frac{x+1}{1-x} \Rightarrow 2u = \ln \frac{x+1}{1-x} \Rightarrow u = \frac{1}{2} \left(\ln \frac{x+1}{1-x} \right)$

The inverse function of $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$ is

$$f^{-1}(x) = \frac{1}{2} \ln \frac{x+1}{1-x}$$

14. $5 \log_2 y - 3 \log_2 (x+4) = 2 \log_2 y + 3 \log_2 x$
 $\Rightarrow 5 \log_2 y - 2 \log_2 y = 3 \log_2 (x+4) + 3 \log_2 x$
 $\Rightarrow \log_2 y^5 - \log_2 y^2 = \log_2 (x+4)^3 + \log_2 x^3$
 $\Rightarrow \log_2 \frac{y^5}{y^2} = \log_2 x^3 (x+4)^3$
 $\Rightarrow y^3 = [x(x+4)]^3 \Rightarrow y = x(x+4)$

15. Point P is 90 m away from a vertical flagpole, which is 11 m high



From the above figure, $\tan \hat{P} = \frac{11}{90}$

$$\hat{P} = \tan^{-1} \left(\frac{11}{90} \right) \approx 6.9^\circ$$

Thus, the angle of elevation is about 6.9°

16. Solving equations

a) $z^4 - (8i - 1)z^2 - 8i = 0$ (1)

Let $z^2 = y$, equation (1) can be written as

$$y^2 - (8i - 1)y - 8i = 0$$

$$\Delta = [-(8i - 1)]^2 - 4(-8i)$$

$$= -63 + 16i$$

Finding square roots of Δ

Let $(a + bi)^2 = -63 + 16i$

$$\Leftrightarrow \begin{cases} a^2 - b^2 = -63 \\ 2ab = 16 \\ a^2 + b^2 = \sqrt{(-63)^2 + (16)^2} \end{cases} \Leftrightarrow \begin{cases} a^2 - b^2 = -63 \\ 2ab = 16 \\ a^2 + b^2 = 65 \end{cases}$$

$$\Leftrightarrow \begin{cases} a^2 - b^2 = -63 \\ a^2 + b^2 = 65 \end{cases}$$

$$2a^2 = 2 \Rightarrow a^2 = 1 \Rightarrow a = \pm 1$$

$$\Leftrightarrow \begin{cases} -a^2 + b^2 = 63 \\ a^2 + b^2 = 65 \end{cases}$$

$$2b^2 = 128 \Rightarrow b^2 = 64 \Rightarrow b = \pm 8$$

Square roots of Δ are $\pm(1+8i)$

$$y_1 = \frac{8i-1+1+8i}{2} = 8i \Rightarrow z^2 = 8i \Rightarrow z = \pm(2+2i)$$

$$y_2 = \frac{8i-1-1-8i}{2} = -1 \Rightarrow z^2 = -1 \Rightarrow z = \pm i$$

$$\text{b) } \begin{cases} 1 + \log_2(-x+2y) = \log_2\{2x-3y\} \\ 3^{5x+y} = \frac{81}{3^{-x-7y}} \end{cases}$$

$$\Rightarrow \begin{cases} \log_2 2 + \log_2(-x+2y) = \log_2(2x-3y) \\ 3^{5x+y} = 3^4 \times 3^{x+7y} \end{cases}$$

$$\Rightarrow \begin{cases} \log_2 2(-x+2y) = \log_2(2x-3y) \\ 3^{5x+y} = 3^{4+x+7y} \end{cases}$$

$$\Rightarrow \begin{cases} 2(-x+2y) = 2x-3y \\ 5x+y = 4+x+7y \end{cases} \Rightarrow \begin{cases} -4x+7y = 0 \\ 4x-6y = 4 \end{cases}$$

$$\begin{cases} y = 4 \\ x = 7 \end{cases}$$

$$S = \{(7, 4)\}$$

17. Given $f(x) = \frac{x^2 - 1}{x^2 - 4}$

a) Domain of definition

$$\text{Dom}f = \{x \in \mathbb{R} : x^2 - 4 \neq 0\} = \mathbb{R} \setminus \{-2, 2\}$$

b) Limits at boundaries

$$\lim_{x \rightarrow \pm\infty} \frac{x^2 - 1}{x^2 - 4} = \lim_{x \rightarrow \pm\infty} \frac{x^2 \left(1 - \frac{1}{x^2}\right)}{x^2 \left(1 - \frac{4}{x^2}\right)} = \lim_{x \rightarrow \pm\infty} \frac{1 - \frac{1}{x^2}}{1 - \frac{4}{x^2}} = 1$$

x	$-\infty$	-2	-1	1	2	$+\infty$		
$x^2 - 1$		$+$	0	$-$	0	$+$		
$x^2 - 4$		$+$	0	$-$	0	$+$		
$\frac{x^2 - 1}{x^2 - 4}$		$+$	\parallel	$-$	0	$-$	\parallel	$+$

$$\lim_{x \rightarrow -2^-} \frac{x^2 - 1}{x^2 - 4} = +\infty, \quad \lim_{x \rightarrow -2^+} \frac{x^2 - 1}{x^2 - 4} = -\infty$$

$$\lim_{x \rightarrow 2^-} \frac{x^2 - 1}{x^2 - 4} = -\infty, \quad \lim_{x \rightarrow 2^+} \frac{x^2 - 1}{x^2 - 4} = +\infty$$

c) Asymptotes

Vertical asymptotes: $x = -2$ and $x = 2$

Horizontal asymptote: $y = 1$

d) Variation table

$$f'(x) = \frac{2x(x^2 - 4) - 2x(x^2 - 1)}{(x^2 - 4)^2}$$

$$= \frac{2x^3 - 8x - 2x^3 + 2x}{(x^2 - 4)^2} = \frac{-6x}{(x^2 - 4)^2}$$

$$f'(x) = 0 \Rightarrow x = 0$$

Variation table

x	$-\infty$	-2	0	2	$+\infty$
$f'(x)$		$+$	$+$ 0 $-$		$+$
$f(x)$		$+\infty$	$\frac{1}{4}$	$+\infty$	1


e) x intercepts: $f(x) = 0 \Rightarrow x^2 - 1 = 0 \Rightarrow x = \pm 1$

$$y \text{ intercept: } f(0) = \frac{1}{4}$$

f) Curve

Additional points

x	-5	-4.2	-3.6	-3.4	-3.2	-3	-2.8	-2.6	-2.4	-2.2	-1.8	-1.6	-1.4	-1.2	-1	-0.8	0.6
y	1.1	1.2	1.3	1.4	1.5	1.6	1.8	2.1	2.7	4.6	-2.9	-1.1	-0.5	-0.2	0.0	0.1	0.2
x	0.8	1	1.2	1.4	1.6	1.8	2.2	2.4	2.6	2.8	3	3.2	3.4	3.6	4.2	5	
y	0.1	0.0	-0.2	-0.5	-1.1	-2.9	4.6	2.7	2.1	1.8	1.6	1.5	1.4	1.3	1.2	1.1	



18. a) 4men and 5 women
- Number of possible committee with no restrictions is ${}^9C_3 = 84$
 - Number of possible committee with 1 man and 2 women is ${}^4C_1 \times {}^5C_2 = 40$
 - Number of possible committee with 2 men and 1 woman if a certain man must be in the committee is ${}^3C_1 \times {}^5C_1 = 15$
- b) (i) There are ${}^9C_3 = 84$ of selecting 3 books from 9 books. There are ${}^8C_2 \times {}^1C_1 = 28$ of selecting 1 dictionary and other 2 books.
- So, required probability is $\frac{28}{84} = \frac{1}{3}$.
- There are ${}^5C_2 \times {}^3C_1 = 30$ of selecting 2 novels and 1 poem book.

So, required probability is $\frac{30}{84} = \frac{5}{14}$.

19. a) Given $f(x) = \frac{x^2 + x + 2}{x + 1}$

$$ax + b + \frac{c}{x + 1} = \frac{x^2 + x + 2}{x + 1}$$

$$\Leftrightarrow \frac{ax(x + 1) + b(x + 1) + c}{x + 1} = \frac{x^2 + x + 2}{x + 1}$$

$$\Leftrightarrow \frac{ax^2 + ax + bx + b + c}{x + 1} = \frac{x^2 + x + 2}{x + 1}$$

$$\Leftrightarrow \frac{ax^2 + (a + b)x + b + c}{x + 1} = \frac{x^2 + x + 2}{x + 1}$$

$$\Rightarrow ax^2 + (a + b)x + b + c = x^2 + x + 2$$

$$\Rightarrow \begin{cases} a = 1 \\ a + b = 1 \\ b + c = 2 \end{cases} \Rightarrow \begin{cases} a = 1 \\ b = 0 \\ c = 2 \end{cases}$$

Then,

$$\int f(x) dx = \int \left(x + \frac{2}{x + 1} \right) dx = \int x dx + 2 \int \frac{1}{x + 1} dx = \frac{x^2}{2} + 2 \ln|x + 1| + c$$

b) $\frac{2y}{x} \frac{dy}{dx} = \frac{y^2}{x^2} - 1 \Rightarrow \frac{dy}{dx} = \left(\frac{y^2}{x^2} - 1 \right) \frac{x}{2y} \Rightarrow \frac{dy}{dx} = \frac{y}{2x} - \frac{x}{2y}$

$$f(x, y) = \frac{y}{2x} - \frac{x}{2y}$$

$$f(tx, ty) = \frac{ty}{2tx} - \frac{tx}{2ty} \Rightarrow f(tx, ty) = \frac{t^0 y}{2x} - \frac{t^0 x}{2y}$$

$$\Rightarrow f(tx, ty) = t^0 \left(\frac{y}{2x} - \frac{x}{2y} \right) \Rightarrow f(tx, ty) = t^0 f(x, y)$$

Then, $f(x, y)$ is homogeneous function of degree 0

To solve the given equation, put $\frac{y}{x} = z \Rightarrow y = zx$

$$\frac{dy}{dx} = z + x \frac{dz}{dx}$$

Then,

$$z + x \frac{dz}{dx} = \frac{zx}{2x} - \frac{x}{2zx} \Leftrightarrow z + x \frac{dz}{dx} = \frac{z^2 x - x}{2xz}$$

$$\begin{aligned} \Leftrightarrow z + x \frac{dz}{dx} &= \frac{z^2 - 1}{2z} \Leftrightarrow x \frac{dz}{dx} = \frac{z^2 - 1}{2z} - z \\ \Leftrightarrow x \frac{dz}{dx} &= \frac{z^2 - 1 - 2z^2}{2z} \Leftrightarrow x \frac{dz}{dx} = \frac{-z^2 - 1}{2z} \\ \Leftrightarrow \frac{2z}{-z^2 - 1} dz &= \frac{1}{x} dx \Leftrightarrow -\int \frac{2z}{z^2 + 1} dz = \int \frac{1}{x} dx \\ \Leftrightarrow -\ln|z^2 + 1| &= \ln|x| + \ln k \Leftrightarrow \ln|z^2 + 1| = -\ln|kx| \\ \Leftrightarrow \ln|z^2 + 1| &= \ln|(kx)^{-1}| \Leftrightarrow \left(\frac{y}{x}\right)^2 + 1 = k^{-1}x^{-1}, \text{ since } z = \frac{y}{x} \\ \Leftrightarrow \frac{y^2}{x^2} + 1 &= cx^{-1}, \quad c = k^{-1} \Leftrightarrow \frac{y^2 + x^2}{x^2} = cx^{-1} \\ \Rightarrow y^2 + x^2 &= cx \end{aligned}$$

20. a) Let X represent the random variable “the number of calls between 09:00 hrs and 10:00 hrs on weekday”. Then $X \sim Po(X)$ and

$$P(X = x) = e^{-\lambda} \frac{\lambda^x}{x!}, \quad x = 0, 1, 2, 3, \dots$$

The probability that the office receives 6 calls between 09:00 hrs and 10:00 hrs on this

$$\text{Wednesday is } P(X = 6) = e^{-5} \frac{5^6}{6!} = 0.146$$

- b) The average number of calls between 09:15 hrs and 09:30 hrs on weekday is 1.25. Let Y represent the random variable “the number of calls in the given 15 minutes”

Then, the probability that the office will receive exactly 3 calls between 09:15 hrs and 09:30 hrs is

$$P(Y = 3) = e^{-1.25} \frac{(1.25)^3}{3!} = 0.0933$$

- c) The required probability is

$${}^5C_2 (0.09326)^2 (0.90674)^3 = 0.0648$$

Answers for Summative Evaluation Three

Learner's Book pages 500 - 502

1. $3 - 5x - x^2 \geq 0$

$$\Delta = (-5)^2 - 4(-1)(3) = 37$$

$$x_1 = \frac{-5 + \sqrt{37}}{2}, \quad x_2 = \frac{-5 - \sqrt{37}}{2}$$

Sign table

x	$-\infty$	$\frac{-5 - \sqrt{37}}{2}$	$\frac{-5 + \sqrt{37}}{2}$	$+\infty$	
$3 - 5x - x^2$	-	0	+	0	-

$$S = \left[\frac{-5 - \sqrt{37}}{2}, \frac{-5 + \sqrt{37}}{2} \right]$$

2. Equation of a circle passing through the points
(0,1), (4,3) and (1,-1)

General equation of a circle is $x^2 + y^2 + ax + by + c = 0$

Using the three points, we have

$$\begin{cases} 1 + b + c = 0 \\ 16 + 9 + 4a + 3b + c = 0 \\ 1 + 1 + a - b + c = 0 \end{cases} \Leftrightarrow \begin{cases} b + c = -1 \Rightarrow b = -1 - c \\ 4a + 3b + c = -25 \\ a - b + c = -2 \end{cases}$$

$$\Leftrightarrow \begin{cases} b = -1 - c \\ 4a + 3(-1 - c) + c = -25 \\ a - (-1 - c) + c = -2 \end{cases} \Leftrightarrow \begin{cases} b = -1 - c \\ 4a - 3 - 3c + c = -25 \\ a + 1 + c + c = -2 \end{cases}$$

$$\Leftrightarrow \begin{cases} 4a - 2c = -22 \\ a + 2c = -3 \end{cases}$$

$$5a = -25 \Rightarrow a = -5$$

$$a + 2c = -3$$

$$\Rightarrow -5 + 2c = -3$$

$$\Rightarrow 2c = 2 \Rightarrow c = 1$$

$$b = -1 - c = -1 - 1 = -2$$

Then, the equation is $x^2 + y^2 - 5x - 2y + 1 = 0$

$$3. \quad \frac{x^2 - x + 1}{x - 1} = k \Leftrightarrow x^2 - x + 1 = kx - k$$

$$\Leftrightarrow x^2 + (-1 - k)x + 1 + k = 0$$

This equation has repeated roots if the discriminant is zero; $\Delta = 0$

$$\Delta = (-1 - k)^2 - 4(1 + k) = 1 + 2k + k^2 - 4 - 4k = k^2 - 2k - 3$$

$$k^2 - 2k - 3 = 0$$

$$\Delta = (-2)^2 - 4(-3) = 4 + 12 = 16$$

$$k_1 = \frac{2 + 4}{2} = 3 \text{ or } k_1 = \frac{2 - 4}{2} = -1$$

Thus, the given equation has repeated roots if $k \in \{-1, 3\}$

4. Consider the following augmented matrix

$$\left(\begin{array}{ccc|c} 1 & 1 & -1 & -1 \\ 3 & -2 & 1 & 0 \\ 2 & 3 & -3 & -3 \end{array} \right) \quad \begin{array}{l} r_2 = r_2 - 3r_1 \\ r_3 = r_3 - 2r_1 \end{array}$$

$$\left(\begin{array}{ccc|c} 1 & 1 & -1 & -1 \\ 0 & -5 & 4 & 3 \\ 0 & 1 & -1 & -1 \end{array} \right) \quad r_3 = 5r_3 + r_2$$

$$\left(\begin{array}{ccc|c} 1 & 1 & -1 & -1 \\ 0 & -5 & 4 & 3 \\ 0 & 0 & -1 & -2 \end{array} \right)$$

The simplified system is

$$\begin{cases} x + y - z = -1 \\ -5y + 4z = 3 \\ -z = -2 \Rightarrow z = 2 \end{cases}$$

$$-5y + 4z = 3 \Rightarrow -5y + 8 = 3 \Rightarrow y = 1$$

$$x + y - z = -1 \Rightarrow x + 1 - 2 = -1 \Rightarrow x = 0$$

Hence, $S = \{(0, 1, 2)\}$

$$5. \quad \lim_{x \rightarrow -\infty} \frac{\sqrt{x^2 + 2}}{3x - 6} = \frac{\infty}{\infty} \text{ I.C}$$

$$\lim_{x \rightarrow -\infty} \frac{\sqrt{x^2 + 2}}{3x - 6} = \lim_{x \rightarrow -\infty} \frac{-x\sqrt{1 + \frac{2}{x^2}}}{x\left(3 - \frac{6}{x}\right)} = \lim_{x \rightarrow -\infty} \frac{-\sqrt{1 + \frac{2}{x^2}}}{3 - \frac{6}{x}} = -\frac{1}{3}$$

6. $U_n = 72 - 6n$, $S_n = 378$

$$U_1 = 72 - 6 = 66$$

$$S_n = \frac{n}{2}(U_1 + U_n) = \frac{n}{2}(66 + 72 - 6n) = 69n - 3n^2$$

But $S_n = 378$, then $69n - 3n^2 = 378$

$$\Rightarrow 3n^2 - 69n + 378 = 0 \Rightarrow n^2 - 23n + 126 = 0$$

$$\Rightarrow (n - 14)(n - 9) = 0$$

Then, $n = 9$ or $n = 14$

7. $(x - 1)(x - 2) + (y + 3)(y - 4) + (z + 1)(z - 1) = 0$

$$\Leftrightarrow x^2 - 2x - x + 2 + y^2 - 4y + 3y - 12 + z^2 - z + z - 1 = 0$$

$$\Leftrightarrow x^2 - 3x + 2 + y^2 - y - 12 + z^2 - 1 = 0$$

$$\Leftrightarrow x^2 - 3x + \frac{9}{4} - \frac{9}{4} + 2 + y^2 - y + \frac{1}{4} - \frac{1}{4} - 12 + z^2 - 0z - 1 = 0$$

$$\Leftrightarrow \left(x^2 - 3x + \frac{9}{4}\right) - \frac{9}{4} + 2 + \left(y^2 - y + \frac{1}{4}\right) - \frac{1}{4} - 12 + (z^2 - 0z) - 1 = 0$$


$$\Leftrightarrow \left(x - \frac{3}{2}\right)^2 + \frac{-9 + 8}{4} + \left(y - \frac{1}{2}\right)^2 + \frac{-1 - 48}{4} + (z - 0)^2 - 1 = 0$$

$$\Leftrightarrow \left(x - \frac{3}{2}\right)^2 + \left(y - \frac{1}{2}\right)^2 + (z - 0)^2 + \frac{-9 + 8}{4} + \frac{-1 - 48}{4} - \frac{4}{4} = 0$$

$$\Leftrightarrow \left(x - \frac{3}{2}\right)^2 + \left(y - \frac{1}{2}\right)^2 + (z - 0)^2 = \frac{54}{4}$$

Centre is $\left(\frac{3}{2}, \frac{1}{2}, 0\right)$ and radius is $\sqrt{\frac{54}{4}} = \frac{3\sqrt{6}}{2}$

8. This argument is not valid. The conclusion is false because not only human beings are mortal. It is a converse error.



9. $f(x) = \sin^2 x \tan x$

$$\begin{aligned} f'(x) &= 2 \sin x \cos x \tan x + \frac{\sin^2 x}{1+x^2} = 2 \sin x \cancel{\cos x} \frac{\sin x}{\cancel{\cos x}} + \frac{\sin^2 x}{1+x^2} \\ &= 2 \sin^2 x + \frac{\sin^2 x}{1+x^2} = \sin^2 x \left(\frac{2+2x^2+1}{1+x^2} \right) \\ &= \sin^2 x \left(\frac{2x^2+3}{1+x^2} \right) \end{aligned}$$

10. a) Equation of the line joining the points $A(3,4,1)$ and $B(5,1,6)$

Direction vector is $\overline{AB} = (2, -3, 5)$

Parametric equations

$$\begin{cases} x = 3 + 2r \\ y = 4 - 3r \\ z = 1 + 5r \end{cases} \text{ where } r \text{ is a parameter}$$

Or symmetric equations

$$\frac{x-3}{2} = \frac{4-y}{3} = \frac{z-1}{5}$$

b) If $z = 0, 1 + 5r = 0 \Rightarrow r = -\frac{1}{5}$

$$\text{and } \begin{cases} x = 3 + 2\left(-\frac{1}{5}\right) = \frac{13}{5} \\ y = 4 - 3\left(-\frac{1}{5}\right) = \frac{23}{5} \\ z = 0 \end{cases}$$

Then, the point is $\left(\frac{13}{5}, \frac{23}{5}, 0\right)$

11. Given the function $f(x) = \cos 3x$

$$\begin{aligned} f(x) &= \cos x & f(0) &= 1 \\ f'(x) &= -3 \sin 3x & f'(0) &= 0 \\ f''(x) &= -9 \cos 3x & f''(0) &= -9 \\ f'''(x) &= 27 \sin 3x & f'''(0) &= 0 \\ f^{(4)}(x) &= 81 \cos 3x & f^{(4)}(0) &= 81 \\ f^{(5)}(x) &= -243 \sin 3x & f^{(5)}(0) &= 0 \\ f^{(6)}(x) &= -729 \cos 3x & f^{(6)}(0) &= -729 \\ &\vdots & & \end{aligned}$$

$$\begin{aligned} \cos x &= 1 + \frac{0}{1!}x + \frac{-9}{2!}x^2 + \frac{0}{3!}x^3 + \frac{81}{4!}x^4 + \frac{0}{5!}x^5 + \frac{-729}{6!}x^6 + \dots \\ &= 1 - \frac{9x^2}{2} + \frac{27x^4}{8} - \frac{81x^6}{80} + \dots \end{aligned}$$

12. $\frac{x^2 + 1}{x^3 + 4x^2 + 3x} = x(x^2 + 4x + 3) = x(x+1)(x+3)$

$$\frac{x^2 + 1}{x^3 + 4x^2 + 3x} = \frac{x^2 + 1}{x(x+1)(x+3)}$$

$$\frac{x^2 + 1}{x(x+1)(x+3)} = \frac{A}{x} + \frac{B}{x+1} + \frac{C}{x+3}$$

$$= \frac{A(x+1)(x+3) + Bx(x+3) + Cx(x+1)}{x(x+1)(x+3)}$$

$$x^2 + 1 = A(x+1)(x+3) + Bx(x+3) + Cx(x+1)$$

Let $x = 0$, $\Rightarrow 1 = 3A \Rightarrow A = \frac{1}{3}$

Let $x = -1$, $\Rightarrow 2 = -2B \Rightarrow B = -1$

Let $x = -3$, $\Rightarrow 10 = 6C \Rightarrow C = \frac{5}{3}$

Then, $\frac{x^2 + 1}{x^3 + 4x^2 + 3x} = \frac{1}{3x} - \frac{1}{x+1} + \frac{5}{3x+9}$

Hence; $\int \frac{x^2 + 1}{x^3 + 4x^2 + 3x} dx = \int \frac{1}{3x} dx - \int \frac{1}{x+1} dx + \int \frac{5}{3x+9} dx$

$$\begin{aligned}
 &= \frac{1}{3} \int \frac{1}{x} dx - \int \frac{1}{x+1} dx + \frac{5}{3} \int \frac{1}{x+3} dx \\
 &= \frac{1}{3} \ln|x| - \ln|x-1| + \frac{5}{3} \ln|x+3| + c
 \end{aligned}$$

$$\begin{aligned}
 13. \quad z &= \frac{2-2i}{1+i} \\
 \Rightarrow z &= \frac{(2-2i)(1-i)}{(1+i)(1-i)} \Rightarrow z = \frac{2-2i-2i-2}{2} \Rightarrow z = -2i
 \end{aligned}$$

$$|z| = 2, \quad \arg(z) = -\frac{\pi}{2}$$

$$\text{Then, } z = 2 \operatorname{cis}\left(-\frac{\pi}{2}\right)$$

$$14. \quad \text{Given that } I = \int_0^{\ln 16} \frac{e^x + 3}{e^x + 4} dx \text{ and } J = \int_0^{\ln 16} \frac{dx}{e^x + 4}$$

$$I + J = \int_0^{\ln 16} \left(\frac{e^x + 3}{e^x + 4} + \frac{1}{e^x + 4} \right) dx = \int_0^{\ln 16} \left(\frac{e^x + 4}{e^x + 4} \right) dx = \int_0^{\ln 16} dx = [x]_0^{\ln 16} = \ln 16$$

$$\begin{aligned}
 I - 3J &= \int_0^{\ln 16} \left(\frac{e^x + 3}{e^x + 4} - \frac{3}{e^x + 4} \right) dx = \int_0^{\ln 16} \left(\frac{e^x}{e^x + 4} \right) dx \\
 &= \left[\ln(e^x + 4) \right]_0^{\ln 16} = \ln(e^{\ln 16} + 4) - \ln(e^0 + 4) \\
 &= \ln(16 + 4) - \ln 4 = \ln 20 - \ln 4 \\
 &= \ln \frac{20}{4} = \ln 5
 \end{aligned}$$

$$\begin{aligned}
 15. \quad U &= \{(a, b, c, d) : b + c + d = 0\} \text{ and} \\
 W &= \{(a, b, c, d) : a + b = 0, c = 2d\}
 \end{aligned}$$

We need to solve

$$\begin{cases}
 b + c + d = 0 \\
 a + b = 0 \Rightarrow a = -b \\
 c = 2d
 \end{cases}$$

$$b + c + d = 0$$

$$\Rightarrow b + 2d + d = 0$$

$$\Rightarrow b = -3d$$

$$a = -b \Rightarrow a = 3d$$

Then $U \cap W = \{(3d, -3d, 2d, d) : d \in \mathbb{R}\}$ and $\dim(U \cap W) = 1$

16. The quarterly, monthly,....rates of interest are found by dividing the nominal annual rate by 4,12,.....

Interest rate	Number of compounding	Value of investment after one year in Frw
a) Annually	1	$100,000 \times (1 + 0.08) = 108,000$
b) Quarterly	4	$100,000 \times \left(1 + \frac{0.08}{4}\right)^4 = 100,000 \times 1.02^4 = 108,240$
c) Monthly	12	$100,000 \times \left(1 + \frac{0.08}{12}\right)^{12} = 100,000 \times 1.0067^{12} = 108,300$
d) Weekly	52	$100,000 \times \left(1 + \frac{0.08}{52}\right)^{52} = 100,000 \times 1.0015^{52} = 108,320$
e) Daily	365	$100,000 \times \left(1 + \frac{0.08}{365}\right)^{365} = 100,000 \times 1.0002^{365} = 108,330$

17. Advertisement sports (x_i) and volume of sales in hundreds (y_i)

x_i	y_i	x_i^2	y_i^2	$x_i y_i$
1	41	1	1681	41
2	50	4	2500	100
3	54	9	2916	162
4	54	16	2916	216
5	57	25	3249	285
6	63	36	3969	378
$\sum_{i=1}^6 x_i = 21$	$\sum_{i=1}^6 y_i = 319$	$\sum_{i=1}^6 x_i^2 = 91$	$\sum_{i=1}^6 y_i^2 = 17231$	$\sum_{i=1}^6 x_i y_i = 1182$

a) Mean: $\bar{x} = \frac{21}{6} = \frac{7}{2}$, $\bar{y} = \frac{319}{6}$

$$\sigma_x^2 = \frac{91}{6} - \left(\frac{7}{2}\right)^2 = \frac{35}{12}$$

Standard deviation for x_i is $\sigma_x = \sqrt{\frac{35}{12}} = \frac{\sqrt{105}}{6}$

$$\sigma_y^2 = \frac{17231}{6} - \left(\frac{319}{6}\right)^2 = \frac{1625}{36}$$

Standard deviation for y_i is $\sigma_y = \sqrt{\frac{1625}{36}} = \frac{5\sqrt{65}}{6}$.

b) Correlation coefficient is given by $r = \frac{\text{cov}(x,y)}{\sigma_x \sigma_y}$

$$\text{cov}(x,y) = \frac{\sum_{i=1}^6 x_i y_i}{6} - \bar{x} \bar{y}$$

$$\text{cov}(x,y) = \frac{1182}{6} - \left(\frac{21}{6}\right)\left(\frac{319}{6}\right) = \frac{7092 - 6699}{36} = \frac{393}{36} = \frac{131}{12}$$

Then,

$$r = \frac{\frac{131}{12}}{\frac{\sqrt{105}}{6} \times \frac{5\sqrt{65}}{6}} = \frac{131}{12} \times \frac{36}{5\sqrt{6825}} = \frac{393}{5\sqrt{6825}} \approx 0.95$$

c) Regression line for y on x

$$y - \bar{y} = \frac{\text{cov}(x,y)}{\sigma_x^2} (x - \bar{x}) \Leftrightarrow y - \frac{319}{6} = \frac{\frac{131}{12}}{\frac{35}{12}} \left(x - \frac{21}{6}\right)$$

$$\Leftrightarrow y - \frac{319}{6} = \frac{131}{35} \left(x - \frac{21}{6}\right) \Leftrightarrow y = \frac{131}{35}x - \frac{131}{35} \times \frac{21}{6} + \frac{319}{6}$$

$$\Leftrightarrow y = \frac{131}{35}x + \frac{601}{15}$$

d) If $x = 7$, $y = \frac{131}{35} \times 7 + \frac{601}{15} \approx 66$

18. Given the vertices of the triangle: $A(1,2,3)$,

$B(-2,1,-4)$ and $C(3,4,-2)$

a) (i) $\overline{AB} = (-3, -1, -7) \Rightarrow \|\overline{AB}\| = \sqrt{9+1+49} = \sqrt{59}$

$$\overline{AC} = (2, 2, -5) \Rightarrow \|\overline{AC}\| = \sqrt{4+4+25} = \sqrt{33}$$

$$\overline{BC} = (5, 3, 2) \Rightarrow \|\overline{BC}\| = \sqrt{25+9+4} = \sqrt{38}$$

The perimeter is $\sqrt{59} + \sqrt{33} + \sqrt{38}$ units of length

$$\text{b) Centre of gravity } \frac{1}{3}(A+B+C) = \frac{1}{3}(2, 7, -3) = \left(\frac{2}{3}, \frac{7}{3}, -1\right)$$

$$\begin{aligned} \text{c) } \angle(\overrightarrow{AB}, \overrightarrow{AC}) &= \cos^{-1}\left(\frac{-6-2+35}{\sqrt{59 \times 33}}\right) \\ &= \cos^{-1}\left(\frac{-32}{\sqrt{2242}}\right) = 132.5^\circ = \cos^{-1}\left(\frac{27}{\sqrt{1947}}\right) = 52.3^\circ \end{aligned}$$

$$\text{Thus, } \theta_1 = 52.3^\circ$$

$$\begin{aligned} \angle(\overrightarrow{AB}, \overrightarrow{BC}) &= \cos^{-1}\left(\frac{-15-3-14}{\sqrt{59 \times 38}}\right) = \cos^{-1}\left(\frac{-32}{\sqrt{2242}}\right) = 132.5^\circ \\ &= \cos^{-1}\left(\frac{-32}{\sqrt{2242}}\right) = 132.5^\circ \end{aligned}$$

$$\text{Therefore, } \theta_2 = 47.5$$

$$\angle(\overrightarrow{AC}, \overrightarrow{BC}) = \cos^{-1}\left(\frac{10+6-10}{\sqrt{33 \times 38}}\right) = \cos^{-1}\left(\frac{10}{\sqrt{1254}}\right) = 80.2^\circ$$

$$\text{Therefore, } \theta_3 = 80.2^\circ$$

$$\text{(or } \theta_3 = 180^\circ - 52.3^\circ - 47.5^\circ = 80.2^\circ \text{).}$$

$$\text{d) The area of triangle } ABC \text{ is given by } \frac{1}{2} \|\overrightarrow{AB} \times \overrightarrow{AC}\|$$

$$\begin{aligned} \frac{1}{2} \|\overrightarrow{AB} \times \overrightarrow{AC}\| &= \frac{1}{2} \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -3 & -1 & -7 \\ 2 & 2 & -5 \end{vmatrix} \\ &= \frac{1}{2} \|(5+14)\vec{i} - (15+14)\vec{j} + (-6+2)\vec{k}\| \\ &= \frac{1}{2} \|19\vec{i} - 29\vec{j} - 4\vec{k}\| = \frac{\sqrt{361+841+16}}{2} = \frac{\sqrt{1218}}{2} \text{ sq. units} \end{aligned}$$

$$19. \quad f(x) = x + |x| + 1 - \frac{1}{x+2}$$

a) Domain of definition

$$\text{Existence condition: } x+2 \neq 0 \Leftrightarrow x \neq -2$$

$$\text{Then, } \text{Dom}f = \mathbb{R} \setminus \{-2\} \text{ or } \text{Dom}f =]-\infty, -2[\cup]-2, +\infty[$$

b) $f(x)$ without the symbol of absolute value

$$f(x) = \begin{cases} x + x + 1 - \frac{1}{x+2}, & x \geq 0 \\ x - x + 1 - \frac{1}{x+2}, & x < 0 \text{ or } x \neq -2 \end{cases}$$

$$\Leftrightarrow f(x) = \begin{cases} 2x + 1 - \frac{1}{x+2}, & x \geq 0 \\ 1 - \frac{1}{x+2}, & x < 0 \text{ or } x \neq -2 \end{cases}$$

$$\text{Or } f(x) = \begin{cases} 2x + 1 - \frac{1}{x+2}, & x \in [0, +\infty[\\ 1 - \frac{1}{x+2}, & x \in]-\infty, -2[\cup]-2, 0[\end{cases}$$

c) Limits on boundaries of domain of definition and asymptotes

$$\lim_{x \rightarrow -2} f(x) = \lim_{x \rightarrow -2} 1 - \frac{1}{x+2} = \infty$$

Thus, $V.A. \equiv x = -2$

Table of sign for determining sided limits:

x	$-\infty$	-2	$+\infty$
$x+2$		$-$	$+$
$-\frac{1}{x+2}$	$+$		$-$

From table of sign, we deduce that

$$\lim_{x \rightarrow -2^-} f(x) = +\infty \text{ and } \lim_{x \rightarrow -2^+} f(x) = -\infty$$

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{x \rightarrow -\infty} 1 - \frac{1}{x+2} = 1$$

Hence, for $x \rightarrow -\infty$, there is $H.A. \equiv y = 1$

$$\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow +\infty} \left(2x + 1 - \frac{1}{x+2} \right) = +\infty$$

Thus, for $x \rightarrow +\infty$, there is no horizontal asymptote.

Let us check if there is an oblique asymptote

For $x \rightarrow +\infty$, $f(x) = 2x + 1 - \frac{1}{x+2}$;

As $\lim_{x \rightarrow +\infty} \frac{1}{x+2} = 0$, $y = 2x + 1$ is oblique asymptote.

$$\lim_{x \rightarrow 0^+} f(x) = \lim_{x \rightarrow 0} \left(2x + 1 - \frac{1}{x+2} \right) = \frac{1}{2} = \lim_{x \rightarrow 0^-} f(x) = \lim_{x \rightarrow 0} \left(1 - \frac{1}{x+2} \right)$$

Therefore, f is continuous at $x = 0$

d) Interval of increasing

$$f(x) = \begin{cases} 2x + 1 - \frac{1}{x+2}, & x \in [0, +\infty[\\ 1 - \frac{1}{x+2}, & x \in]-\infty, -2[\cup]-2, 0[\end{cases} \Rightarrow f'(x) = \begin{cases} 2 + \frac{1}{(x+2)^2}, & x \in [0, +\infty[\\ \frac{1}{(x+2)^2}, & x \in]-\infty, -2[\cup]-2, 0[\end{cases}$$



As $f'(x) > 0, \forall x \in \text{Dom}f$, f is increasing on its domain of definition.

e) Concavity

$$f'(x) = \begin{cases} 2 + \frac{1}{(x+2)^2}, & x \in [0, +\infty[\\ \frac{1}{(x+2)^2}, & x \in]-\infty, -2[\cup]-2, 0[\end{cases}$$

$$\Rightarrow f''(x) = -\frac{x+2}{(x+2)^4} = -\frac{1}{(x+2)^3}, \forall x \in \text{Dom}f$$

f) Table of variation

x	$-\infty$	-2	0	$+\infty$
$f'(x)$	+		+	+
$f''(x)$	+		-	
$f(x)$	 $H.A. \equiv x = 1$		 $V.A. \equiv x = -2$ $O.A. \equiv y = 2x + 1$	

g) Curve sketching

Additional points:


For $x < -2$

x	-5	-4.5	-4	-3.5	-3	-2.5	-2.2
y	1.3	1.4	1.5	1.7	2	3	6

For $x > -2$

x	-1.8	-1.5	-1	-0.5	0	0.5	1	1.5	2	2.5	3
y	-4	-1	0	-0.7	0.5	1.6	2.7	3.7	4.8	5.8	6.8

Curve



20. a) Let $P(x, y)$ be any point on the parabola using focus-directrix ($F - M$) property of the parabola: $\overline{FP} = \overline{PM}$.

$$\text{Therefore, } \sqrt{(x+1)^2 + (y+2)^2} = \frac{|x-2y+3|}{\sqrt{1^2 + (-2)^2}}$$

$$\Leftrightarrow (x+1)^2 + (y+2)^2 = \frac{(x-2y+3)^2}{5}$$

$$\Leftrightarrow x^2 + 2x + 1 + y^2 + 4y + 4 = \frac{x^2 + 4y^2 + 9 - 4xy + 6x - 12y}{5}$$

$$\Leftrightarrow 5x^2 + 10x + 5y^2 + 20y + 25 = x^2 + 4y^2 + 9 - 4xy + 6x - 12y$$

$\Leftrightarrow 4x^2 + y^2 + 4xy + 4x + 32y + 16 = 0$ which is the required equation of the parabola.

- b) Let $P(x, y)$ be any point of focus and the given point $(0, 4)$ be denoted by A .

Then, $PA = \frac{2}{3} \times$ distance of P from the line $y = 9$.

$$\Leftrightarrow \sqrt{x^2 + (y-4)^2} = \frac{2}{3} \times \frac{|y-9|}{\sqrt{0^2+1^2}} \Leftrightarrow x^2 + (y-4)^2 = \frac{4}{9} \times (y-9)^2$$

$$\Leftrightarrow x^2 + y^2 - 8y + 16 = \frac{4}{9}(y^2 - 18y + 81)$$

$$\Leftrightarrow 9x^2 + 9y^2 - 72y + 144 = 4y^2 - 72y + 324$$

$\Leftrightarrow 9x^2 + 5y^2 - 180 = 0$ which is the required equation of locus.

- c) The equation of hyperbola is $x^2 - 4y^2 = 4 \Leftrightarrow \frac{x^2}{4} - \frac{y^2}{1} = 1$

Here, $a^2 = 4$, $b^2 = 1$ then, $a = 2$, $b = 1$.

Therefore, axes are 4 and 2.

$$b^2 = a^2(e^2 - 1) \text{ or } 1 = 4(e^2 - 1) \text{ which gives } e = \frac{\sqrt{5}}{2}$$

$$\text{Thus, Eccentricity} = \frac{\sqrt{5}}{2}$$

Since, coordinates of foci are given by $(\pm ae, 0)$, then

$$\text{they are } \left(\pm 2 \times \frac{\sqrt{5}}{2}, 0 \right) \text{ or } (\pm \sqrt{5}, 0)$$

$$\text{Length of latus rectum is } \frac{2b^2}{a} = \frac{2 \times 1}{2} = 1$$

Alternative method:

$$\text{From } \frac{x^2}{4} - \frac{y^2}{1} = 1, \quad a = 2, \quad b = 1;$$

$$\text{For hyperbola } c^2 = a^2 + b^2$$

$$\text{Here, } c^2 = 4 + 1 = 5 \text{ or } c = \sqrt{5}$$

► Axes are $2a = 4$ and $2b = 2$

► Coordinates of foci $F(\pm c, 0) = (\pm \sqrt{5}, 0)$

► Eccentricity $e = \frac{c}{a} = \frac{\sqrt{5}}{2}$

► Length of latus rectum is equal to $\frac{2b^2}{a} = \frac{2 \times 1}{2} = 1$

Answers for Summative Evaluation Four

Learner's Book pages 503 - 506

$$1. \quad \left(x + \frac{1}{x}\right)^{20} = (x + x^{-1})^{20}$$

$$= {}^{20}C_r x^r (x^{-1})^{20-r} = {}^{20}C_r x^r x^{-20+r} = {}^{20}C_r x^{2r-20}$$

For the independent term, $2r - 20 = 0$ or $r = 10$

Then, the independent term is ${}^{20}C_{10} = \frac{20!}{10!10!} = 184756$

$$2. \quad \text{If } 6x^3 + 7x^2 + ax + b \text{ is divisible by } x - 2, \text{ the remainder}$$

$$\text{is } 6(2)^3 + 7(2)^2 + 2a + b = 72$$

$$48 + 28 + 2a + b = 72$$

$$\Rightarrow 2a + b = -4$$

Also, $6x^3 + 7x^2 + ax + b$ is exactly divisible by $x + 1$ then

$$6(-1)^3 + 7(-1)^2 - a + b = 0$$

$$-6 + 7 - a + b = 0$$

$$\Rightarrow -a + b = -1$$

$$\begin{cases} 2a + b = -4 \\ -a + b = -1 \end{cases} \Rightarrow \begin{cases} 2a + b = -4 \\ \underline{a - b = 1} \end{cases}$$

$$3a = -3 \Rightarrow a = -1$$

$$-a + b = -1 \Rightarrow 1 + b = -1 \Rightarrow b = -2 \quad \begin{cases} a = -1 \\ b = -2 \end{cases}$$

$$3. \quad \sin x + \sqrt{3} \cos x = 1$$

$$\text{Let } \sqrt{3} = \tan \alpha \Rightarrow \alpha = \frac{\pi}{3}$$

$$\sin x + \tan \alpha \cos x = 1 \Rightarrow \sin x + \frac{\sin \alpha}{\cos \alpha} \cos x = 1$$

$$\Rightarrow \sin x \cos \alpha + \sin \alpha \cos x = \cos \alpha$$

$$\Rightarrow \sin(x + \alpha) = \cos \alpha \Rightarrow \sin\left(x + \frac{\pi}{3}\right) = \cos \frac{\pi}{3}$$

$$\Rightarrow \sin\left(x + \frac{\pi}{3}\right) = \frac{1}{2}$$

$$x + \frac{\pi}{3} = \begin{cases} \frac{\pi}{6} + 2k\pi \\ \frac{5\pi}{6} + 2k\pi \end{cases}, \quad k \in \mathbb{Z} \qquad x = \begin{cases} -\frac{\pi}{6} + 2k\pi \\ \frac{\pi}{2} + 2k\pi \end{cases}$$

$$S = \left\{ -\frac{\pi}{6} + 2k\pi, \frac{\pi}{2} + 2k\pi \right\}, \quad k \in \mathbb{Z}$$

4. A matrix has no inverse if its determinant is zero

$$\begin{vmatrix} 11-x & 2 & 8 \\ 2 & 2-x & -10 \\ 8 & -10 & 5-x \end{vmatrix} = 0$$

$$\Rightarrow (11-x)(2-x)(5-x) - 160 - 160 - 64(2-x) - 100(11-x) - 4(5-x) = 0$$

$$\Rightarrow 110 - 65x + 5x^2 - 22x + 13x^2 - x^3 - 320 - 128 + 64x - 1100 + 100x - 20 + 4x = 0$$

$$\Rightarrow -x^3 + 18x^2 + 81x - 1458 = 0$$

$$\Rightarrow x^3 - 18x^2 - 81x + 1458 = 0$$

9 is one of the roots

	1	-18	-81	1458
9		9	-81	-1458
	1	-9	-162	0

$$x^3 - 18x^2 - 81x + 1458 = (x-9)(x^2 - 9x - 162)$$

$$x^2 - 9x - 162 = 0 \Rightarrow (x+9)(x-18) = 0 \Rightarrow x = -9 \text{ or } x = 18$$

Thus, the given matrix is singular if $x \in \{-9, 9, 18\}$

5.
$$\begin{cases} \log(x+y) = 1 \\ \log_2 x + 2\log_4 y = 4 \end{cases}$$

$$\log(x+y) = 1 \Leftrightarrow \log(x+y) = \log 10 \Rightarrow x+y = 10$$

$$\log_2 x + 2\log_4 y = 4$$

$$\Leftrightarrow \log_2 x + 2\frac{\log_2 y}{\log_2 4} = 4$$

$$\Leftrightarrow \log_2 x + 2\frac{\log_2 y}{2} = 4$$

$$\Leftrightarrow \log_2 x + \log_2 y = 4\log_2 2$$

$$\Leftrightarrow \log_2 xy = \log_2 2^4$$

$$\Rightarrow xy = 16$$

Now,

$$\begin{cases} x+y=10 \Rightarrow x=10-y \\ xy=16 \end{cases}$$

$$(10 - y)y = 16 \Rightarrow 10y - y^2 - 16 = 0 \Rightarrow y^2 - 10y + 16 = 0$$

$$(y - 2)(y - 8) = 0$$

$$y - 2 = 0 \Rightarrow y = 2 \Rightarrow x = 10 - y = 10 - 2 = 8$$

$$y - 8 = 0 \Rightarrow y = 8$$

$$\Rightarrow x = 10 - 8 = 2$$

6. $x^2 - x - 3 = 0$

We know that for the equation of the form $ax^2 + bx + c = 0$,

if α and β are the roots, then, $\alpha + \beta = -\frac{b}{a}$ and $\alpha\beta = \frac{c}{a}$

Here, $\alpha + \beta = 1$ and $\alpha\beta = -3$

$$(\alpha + \beta)^3 = \alpha^3 + 3\alpha^2\beta + 3\alpha\beta^2 + \beta^3$$

$$\Leftrightarrow (\alpha + \beta)^3 - 3\alpha^2\beta - 3\alpha\beta^2 = \alpha^3 + \beta^3$$

$$\Leftrightarrow \alpha^3 + \beta^3 = (\alpha + \beta)^3 - 3\alpha\beta(\alpha + \beta)$$

$$\Leftrightarrow \alpha^3 + \beta^3 = (1)^3 - 3(-3)(1) = 10$$

Then, $\alpha^3 + \beta^3 = 10$

7. $x^2 + 4y^2 - 4x + 8y + 4 = 0$

$$\Leftrightarrow x^2 - 4x + 4y^2 + 8y + 4 = 0 \Leftrightarrow x^2 - 4x + 4 + 4(y^2 + 2y) = 0$$

$$\Leftrightarrow (x - 2)^2 + 4[(y + 1)^2 - 1] = 0 \Leftrightarrow (x - 2)^2 + 4(y + 1)^2 - 4 = 0$$

$$\Leftrightarrow (x - 2)^2 + 4(y + 1)^2 = 4 \Leftrightarrow \frac{(x - 2)^2}{4} + \frac{(y + 1)^2}{1} = 1$$

The centre is $(2, -1)$

$$a^2 = 4, b^2 = 1$$

The eccentricity is $e = \sqrt{\frac{a^2 - b^2}{a^2}} = \sqrt{\frac{4 - 1}{4}} = \frac{\sqrt{3}}{2}$

Foci are $(2e + 2, -1)$ and $(-2e + 2, -1)$

Or $(\sqrt{3} + 2, -1)$ and $(-\sqrt{3} + 2, -1)$

8. Tangent and normal line to the curve

$$3x^2 - xy - 2y^2 + 12 = 0 \text{ at point } (2, 3)$$

$$(3x^2 - xy - 2y^2 + 12)' = 6x - (y + xy') - 4yy'$$

$$6x - y - xy' - 4yy' = 0 \Rightarrow xy' + 4yy' = 6x - y$$

$$\Rightarrow y'(x + 4y) = 6x - y \Rightarrow y' = \frac{6x - y}{x + 4y}$$

$$y'_{(2,3)} = \frac{6(2) - 3}{2 + 4(3)} = \frac{9}{14}$$

$$T \equiv y - 3 = \frac{9}{14}(x - 2)$$

$$\equiv y = \frac{9}{14}x - \frac{9}{7} + 3$$

$$\equiv y = \frac{9}{14}x + \frac{12}{7}$$

$$N \equiv y - 3 = -\frac{14}{9}(x - 2)$$

$$\equiv y = -\frac{14}{9}x + \frac{28}{9} + 3$$

$$\equiv y = -\frac{14}{9}x + \frac{55}{9}$$

$$9. \int_0^1 \frac{1}{(2x+k)^2} dx = \frac{1}{3}$$

$$\int_0^1 \frac{1}{(2x+k)^2} dx = -\frac{1}{2} \left[\frac{1}{2x+k} \right]_0^1 = -\frac{1}{2} \left(\frac{1}{2+k} - \frac{1}{k} \right) = -\frac{1}{4+2k} + \frac{1}{2k}$$

$$\Rightarrow -\frac{1}{4+2k} + \frac{1}{2k} = \frac{1}{3}$$

$$\Rightarrow \frac{-2k + 4 + 2k}{2k(4+2k)} = \frac{1}{3} \Rightarrow \frac{4}{2k(4+2k)} = \frac{1}{3}$$

$$\Rightarrow 4k^2 + 8k = 12 \Rightarrow 4k^2 + 8k - 12 = 0 \Rightarrow k^2 + 2k - 3 = 0$$

$$\Rightarrow (k+3)(k-1) = 0 \Rightarrow k = -3 \text{ or } k = 1$$

$$10. z^6 = 1$$

$$z_k = \text{cis} \frac{2k\pi}{6} = \text{cis} \frac{k\pi}{3}, k = 0, 1, 2, 3, 4, 5$$

$$z_0 = \text{cis} 0 = 1$$

$$z_1 = \text{cis} \frac{\pi}{3} = \frac{1}{2} + i \frac{\sqrt{3}}{2}$$

$$z_2 = \text{cis} \frac{2\pi}{3} = -\frac{1}{2} + i \frac{\sqrt{3}}{2}$$

$$z_3 = \text{cis} \frac{3\pi}{3} = -1$$

$$z_4 = \text{cis} \frac{4\pi}{3} = -\frac{1}{2} - i \frac{\sqrt{3}}{2}$$

$$z_5 = \text{cis} \frac{5\pi}{3} = \frac{1}{2} - i \frac{\sqrt{3}}{2}$$

$$S = \left\{ 1, \frac{1}{2} + i \frac{\sqrt{3}}{2}, -\frac{1}{2} + i \frac{\sqrt{3}}{2}, -1, -\frac{1}{2} - i \frac{\sqrt{3}}{2}, \frac{1}{2} - i \frac{\sqrt{3}}{2} \right\}$$

11. u_1 and d be the first term and the common difference respectively

$$u_1 + (u_1 + d) + (u_1 + 2d) + (u_1 + 3d) + (u_1 + 4d) + (u_1 + 5d) = 72$$

$$\Rightarrow 6u_1 + 15d = 72$$

But $u_2 = 7u_5$ or

$$u_1 + d = 7(u_1 + 4d) \Rightarrow u_1 + d - 7u_1 - 28d = 0 \Rightarrow -6u_1 - 27d = 0$$

$$\begin{cases} 6u_1 + 15d = 72 \\ -6u_1 - 27d = 0 \end{cases}$$

$$-12d = 72 \Rightarrow d = -6$$

$$6u_1 + 15d = 72 \Rightarrow 6u_1 - 90 = 72 \Rightarrow u_1 = 27$$

12. $\lim_{x \rightarrow \frac{\pi}{2}} (\tan x)^{\cos x} = \lim_{x \rightarrow \frac{\pi}{2}} e^{\ln(\tan x)^{\cos x}} = \lim_{x \rightarrow \frac{\pi}{2}} e^{\cos x \ln(\tan x)} = e^{\lim_{x \rightarrow \frac{\pi}{2}} \cos x \ln(\tan x)}$

$$\lim_{x \rightarrow \frac{\pi}{2}} \cos x \ln(\tan x) = 0 \times \infty \text{ IC}$$

$$\lim_{x \rightarrow \frac{\pi}{2}} \cos x \ln(\tan x) = \lim_{x \rightarrow \frac{\pi}{2}} \frac{\ln(\tan x)}{\frac{1}{\cos x}} = \lim_{x \rightarrow \frac{\pi}{2}} \frac{(\tan x)'}{\frac{\sin x}{\cos^2 x}} \quad \text{L'Hôpital's rule}$$

$$= \lim_{x \rightarrow \frac{\pi}{2}} \frac{\cos^2 x}{\cos^2 x \tan x \sin x}$$

$$= \lim_{x \rightarrow \frac{\pi}{2}} \frac{1}{\tan x \sin x} = \frac{1}{\tan \frac{\pi}{2} \sin \frac{\pi}{2}} = 0$$

Then, $\lim_{x \rightarrow \frac{\pi}{2}} (\tan x)^{\cos x} = e^0 = 1$

13. $\bar{x} = 6.2$, $\sigma_x = 3.03315$, $\bar{y} = 2.04$, $\sigma_y = 0.461519$ and $r_{xy} = 0.957241$

The regression line of y on x is $y - \bar{y} = \frac{\text{cov}(x, y)}{\sigma_x^2} (x - \bar{x})$

But $r_{xy} = \frac{\text{cov}(x, y)}{\sigma_x \sigma_y} \Leftrightarrow \text{cov}(x, y) = r_{xy} \sigma_x \sigma_y$

Then, $y - \bar{y} = \frac{r_{xy} \sigma_x \sigma_y}{\sigma_x^2} (x - \bar{x})$ or

$$y - \bar{y} = \frac{r_{xy}\sigma_y}{\sigma_x}(x - \bar{x}) \Leftrightarrow y = \frac{r_{xy}\sigma_y}{\sigma_x}(x - \bar{x}) + \bar{y}$$

$$\Rightarrow y = \frac{0.957241 \times 0.461519}{3.03315}(x - 6.2) + 2.04$$

$$\Rightarrow y = 0.145x + 1.141$$

14. Let S be the sample space, then $n(S) = {}^{52}C_2 = 1326$

Let E be the event of getting two Kings.

$$n(E) = {}^4C_2 = 6$$

$$\text{Then, } P(E) = \frac{6}{1326} = \frac{1}{221}$$

15. Let θ be the angle between vectors $(2, 5)$ and $(-1, 3)$

$$\cos \theta = \frac{(2,5) \cdot (-1,3)}{\|(2,5)\| \|(-1,3)\|} = \frac{-2+15}{\sqrt{4+25}\sqrt{1+9}} = \frac{13\sqrt{290}}{290}$$

$$\theta = \cos^{-1}\left(\frac{13\sqrt{290}}{290}\right) = 40.24 \text{ deg} = 0.70 \text{ rad}$$


16. $y = x^2 - 5x + 4$ and $y = -2x^2 + 5x + 1$

Intersection:

$$x^2 - 5x + 4 = -2x^2 + 5x + 1 \Rightarrow 3x^2 - 10x + 3 = 0$$

$$\Rightarrow (x-3)(3x-1) = 0 \Rightarrow x = 3 \text{ or } x = \frac{1}{3}$$

The curves intersect at $x = 3$ and $x = \frac{1}{3}$



The area of the region enclosed between the two curves is:

$$\int_{1/3}^3 (-2x^2 + 5x + 1 - x^2 + 5x - 4) dx = \int_{1/3}^3 (-3x^2 + 10x - 3) dx$$

$$= \left[-x^3 + 5x^2 - 3x \right]_{1/3}^3$$

$$= -3^3 + 5(3)^2 - 3(3) + \left(\frac{1}{3}\right)^3 - 5\left(\frac{1}{3}\right)^2 + 3\left(\frac{1}{3}\right)$$

$$= -27 = 45 - 9 + \frac{1}{27} - \frac{5}{9} + 1$$

$$= \frac{256}{27} \text{ sq. units}$$

17. a) i) $P(A \cup C) = P(A) + P(C) - P(A \cap C)$

But $P(A \cap C) = 0$ since A and C are mutually exclusive events

$$P(A \cup C) = P(A) + P(C) = \frac{2}{3} + \frac{1}{5} = \frac{13}{15}$$

Since A and B are independent events,

$$P(A \cap B) = P(A)P(B) \Rightarrow P(B) = \frac{P(A \cap B)}{P(A)}$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cup B) = P(A) + \frac{P(A \cap B)}{P(A)} - P(A \cap B)$$

$$\frac{4}{5} = \frac{2}{3} + \frac{P(A \cap B)}{\frac{2}{3}} - P(A \cap B)$$

$$\Rightarrow \frac{4}{5} - \frac{2}{3} = \frac{3}{2} P(A \cap B) - P(A \cap B)$$

$$\Rightarrow \frac{2}{15} = \frac{P(A \cap B)}{2}$$

$$\Rightarrow P(A \cap B) = \frac{4}{15}$$

$$P(B) = \frac{P(A \cap B)}{P(A)} = \frac{\frac{4}{15}}{\frac{2}{3}} = \frac{2}{5}$$

ii) B and C are independent events if

$$P(B \cap C) = P(B)P(C)$$

$$P(B)P(C) = \frac{2}{5} \times \frac{1}{5} = \frac{2}{25}$$

$$\begin{aligned} P(B \cap C) &= P(B) + P(C) - P(B \cup C) \\ &= \frac{2}{5} + \frac{1}{5} - \frac{13}{25} = \frac{2}{25} \end{aligned}$$

$$P(B \cap C) = P(B)P(C)$$

Thus, B and C are independent events.

b) Let $X_i, i=1,2,3$ be the event "patient have the virus" and let D be the event "selected patient recovers".

We need $P(X_3 | D)$

$$P(X_3 | D) = \frac{P(D | X_3)P(X_3)}{\sum_{i=1}^3 P(D | X_i)} = \frac{\frac{1}{8} \times \frac{1}{8}}{\frac{1}{2} \times \frac{1}{2} + \frac{3}{8} \times \frac{3}{8} + \frac{1}{8} \times \frac{1}{8}} = \frac{1}{26}$$

18. a) Given the points $A(2, -3, -1), B(3, -4, 2)$ and $C(4, -5, 2)$

(i) $\overrightarrow{AB} = (1, -1, 3), \overrightarrow{AC} = (2, -2, 3)$

$$\begin{aligned} \overrightarrow{AB} \times \overrightarrow{AC} &= \begin{vmatrix} \vec{i} & 1 & 2 \\ \vec{j} & -1 & -2 \\ \vec{k} & 3 & 3 \end{vmatrix} \\ &= \vec{i} \begin{vmatrix} -1 & -2 \\ 3 & 3 \end{vmatrix} - \vec{j} \begin{vmatrix} 1 & 2 \\ 3 & 3 \end{vmatrix} + \vec{k} \begin{vmatrix} 1 & 2 \\ -1 & -2 \end{vmatrix} \\ &= \vec{i}(-3+6) - \vec{j}(3-6) + \vec{k}(-2+2) \\ &= 3\vec{i} + 3\vec{j} \end{aligned}$$

(ii) The area of triangle ABC is given by $\frac{1}{2} \|\overrightarrow{AB} \times \overrightarrow{AC}\|$

$$\frac{1}{2} \|\overrightarrow{AB} \times \overrightarrow{AC}\| = \frac{1}{2} \|3\vec{i} + 3\vec{j}\| = \frac{1}{2} \sqrt{9+9} = \frac{3\sqrt{2}}{2} \text{ sq. units}$$

b) We have points $A(2,1,1)$, $B(0,5,3)$

(i) Direction vector is $\overline{AB} = (-2, 4, 2)$

Parametric equations:

$$\begin{cases} x = 2 - 2r \\ y = 1 + 4r \\ z = 1 + 2r \end{cases}, \quad r \text{ is a parameter}$$

(ii) Given $C(5, -4, 2)$

Vectors \overline{CD} is perpendicular to vector \overline{AB} if

$$\overline{CD} \cdot \overline{AB} = 0$$

Let $D(x, y, z)$ be the point on line AB then

$$\overline{CD} = (x - 5, y + 4, z - 2) \text{ and}$$

$$\begin{aligned} \overline{CD} \cdot \overline{AB} &= -2(x - 5) + 4(y + 4) + 2(z - 2) \\ &= -2x + 10 + 4y + 16 + 2z - 4 \\ &= -2x + 4y + 2z + 22 \end{aligned}$$

But

$$\begin{cases} x = 2 - 2r \\ y = 1 + 4r \\ z = 1 + 2r \end{cases}$$

Then

$$\begin{aligned} -2(2 - 2r) + 4(1 + 4r) + 2(1 + 2r) + 22 &= 0 \\ \Rightarrow -4 + 4r + 4 + 16r + 2 + 4r + 22 &= 0 \\ \Rightarrow 24r + 24 &= 0 \\ \Rightarrow r &= -1 \end{aligned} \quad \begin{cases} x = 4 \\ y = -3 \\ z = -1 \end{cases}$$

And point D has coordinates $(4, -3, -1)$

(iii) If the plane π contains the line AB , the vector \overline{CD} is perpendicular to the plane π since this vector is also perpendicular to the line AB . So this is a contradiction, no plane can contain the line AB and be parallel to CD .

19. $U = \operatorname{cis} \frac{2\pi}{5} = \cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5}$

$$\begin{aligned} \text{a) } \frac{1}{U} &= \frac{1}{\cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5}} = \cos \frac{2\pi}{5} - i \sin \frac{2\pi}{5} \\ \frac{1}{2} \left(U + \frac{1}{U} \right) &= \frac{1}{2} \left(\cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5} + \cos \frac{2\pi}{5} - i \sin \frac{2\pi}{5} \right) \\ &= \frac{1}{2} \left(2 \cos \frac{2\pi}{5} \right) = \cos \frac{2\pi}{5} \quad \text{as required} \end{aligned}$$

$$\begin{aligned} \text{b) } U^5 &= \left(\cos \frac{2\pi}{5} + i \sin \frac{2\pi}{5} \right)^5 \\ &= \cos \frac{5 \times 2\pi}{5} + i \sin \frac{5 \times 2\pi}{5} = \cos 2\pi + i \sin 2\pi = 1 \end{aligned}$$

$$\begin{aligned} \text{c) } \text{Since } U &= \text{cis} \frac{2\pi}{5} \\ U^0 &= \text{cis} \frac{0 \times 2\pi}{5}, U^1 = \text{cis} \frac{1 \times 2\pi}{5} = 1, U^2 = \text{cis} \frac{2 \times 2\pi}{5} = \text{cis} \frac{4\pi}{5}, \\ U^3 &= \text{cis} \frac{3 \times 2\pi}{5} = \text{cis} \frac{6\pi}{5}, U^4 = \text{cis} \frac{4 \times 2\pi}{5} = \text{cis} \frac{8\pi}{5} \end{aligned}$$

These are five fifth roots of unit. Then, their sum must be zero. Hence, $U^4 + U^3 + U^2 + U + 1 = 0$

$$\begin{aligned} \text{d) } U^4 + U^3 + U^2 + U + 1 &= \text{cis} \frac{8\pi}{5} + \text{cis} \frac{6\pi}{5} + \text{cis} \frac{4\pi}{5} + \text{cis} \frac{2\pi}{5} + 1 \\ U^4 + U^3 + U^2 + U + 1 &= \cos \frac{8\pi}{5} + \cos \frac{6\pi}{5} + \cos \frac{4\pi}{5} \\ &+ \cos \frac{2\pi}{5} + 1 + i \left(\sin \frac{8\pi}{5} + \sin \frac{6\pi}{5} + \sin \frac{4\pi}{5} + \sin \frac{2\pi}{5} \right) \end{aligned}$$

Take the real party

$$\cos \frac{8\pi}{5} + \cos \frac{6\pi}{5} + \cos \frac{4\pi}{5} + \cos \frac{2\pi}{5} + 1 = 0$$

We know that $\cos \alpha = \cos(2\pi - \alpha)$, then

$$\cos \frac{8\pi}{5} = \cos \left(2\pi - \frac{8\pi}{5} \right) = \cos \frac{2\pi}{5}$$

$$\cos \frac{6\pi}{5} = \cos \left(2\pi - \frac{6\pi}{5} \right) = \cos \frac{4\pi}{5}$$

$$\text{Also, } \cos 2\alpha = 2 \cos^2 \alpha - 1 \Rightarrow \cos \frac{4\pi}{5} = 2 \cos^2 \frac{2\pi}{5} - 1$$

$$\cos \frac{2\pi}{5} + 2 \cos^2 \frac{2\pi}{5} - 1 + 2 \cos^2 \frac{2\pi}{5} - 1 + \cos \frac{2\pi}{5} + 1 = 0$$

$$\Rightarrow 4 \cos^2 \frac{2\pi}{5} + 2 \cos \frac{2\pi}{5} - 1 = 0$$

$$\text{But } x = U + \frac{1}{U} = 2 \cos \frac{2\pi}{5}$$

$$\text{Then, } 4 \cos^2 \frac{2\pi}{5} + 2 \cos \frac{2\pi}{5} - 1 = x^2 + x - 1 \text{ or } x^2 + x - 1 = 0$$

e) $x^2 + x - 1 = 0$

$$\Delta = 1 + 4 = 5$$

$$x_1 = \frac{-1 + \sqrt{5}}{2}, \quad x_2 = \frac{-1 - \sqrt{5}}{2}$$

$$x_2 = \frac{-1 - \sqrt{5}}{2} \text{ is to be rejected.}$$

$$\text{For } x_1 = \frac{-1 + \sqrt{5}}{2}, \quad 2 \cos \frac{2\pi}{5} = \frac{-1 + \sqrt{5}}{2} \Rightarrow \cos \frac{2\pi}{5} = \frac{-1 + \sqrt{5}}{4}$$

20. a) Taking 5 men together, 4 women together, 3 children together we have $3!$ ways.

But 5 men can be permuted among them in $5!$ ways, 4 women can be permuted among them in $4!$ ways and 3 children can be permuted among them in $3!$ ways

The total ways is $3!5!4!3! = 103680$ ways

$$\text{b) } \begin{cases} {}^x C_y = {}^x C_{y+1} \\ 4({}^x C_y) = 5({}^x C_{y-1}) \end{cases}$$

$${}^x C_y = {}^x C_{y+1} \Leftrightarrow \frac{x!}{y!(x-y)!} = \frac{x!}{(y+1)!(x-y-1)!}$$

$$\Rightarrow (y+1)!(x-y-1)! = y!(x-y)!$$

$$\Rightarrow (y+1)y!(x-y-1)! = y!(x-y)(x-y-1)!$$

$$\Rightarrow y+1 = x-y$$

$$\Rightarrow -x + 2y = -1$$

$$4({}^x C_y) = 5({}^x C_{y-1}) \Leftrightarrow 4 \frac{x!}{y!(x-y)!} = 5 \frac{x!}{(y-1)!(x-y+1)!}$$

$$\Rightarrow 4(y-1)!(x-y+1)! = 5y!(x-y)!$$

$$\Rightarrow 4(y-1)!(x-y+1)(x-y)! = 5y(y-1)!(x-y)!$$

$$\Rightarrow 4(x-y+1) = 5y$$

$$\Rightarrow 4x - 4y + 4 = 5y$$

$$\Rightarrow 4x - 9y = -4$$

$$\begin{cases} -x + 2y = -1 \\ 4x - 9y = -4 \end{cases} \Leftrightarrow \begin{cases} -4x + 8y = -4 \\ 4x - 9y = -4 \end{cases}$$

$$-y = -8 \Rightarrow y = 8$$

$$-x + 2y = -1$$

$$\Rightarrow -x + 16 = -1$$

$$\Rightarrow x = 17$$

Thus, $S = \{(17, 8)\}$

c) ${}^{n-2}C_m + 2({}^{n-2}C_{m-1}) + {}^{n-2}C_{m-2} = {}^n C_m$

$$\frac{(n-2)!}{m!(n-m-2)!} + \frac{2(n-2)!}{(m-1)!(n-m-1)!} + \frac{(n-2)!}{(m-2)!(n-m)!}$$

$$= \frac{\frac{n!}{n(n-1)}}{m!(n-m)!} + \frac{2 \frac{n!}{n(n-1)}}{m!(n-m)!} + \frac{\frac{n!}{n(n-1)}}{m!(n-m)!}$$

$$\frac{(n-m)(n-m-1)}{(n-m)(n-m-1)} \quad \frac{2}{m} \quad \frac{1}{n-m} \quad \frac{1}{m(m-1)}$$

$$= \frac{n!(n-m)(n-m-1)}{m!(n-m)!n(n-1)} + \frac{2n!m(n-m)}{m!(n-m)!n(n-1)} + \frac{n!m(m-1)}{m!(n-m)!n(n-1)}$$

$$= \frac{n!}{m!(n-m)!} \left[\frac{(n-m)(n-m-1) + 2m(n-m) + m(m-1)}{n(n-1)} \right]$$

$$= \frac{n!}{m!(n-m)!} \left[\frac{n^2 - nm - n - mn + m^2 + m + 2mn - 2m^2 + m^2 - m}{n(n-1)} \right]$$

$$= \frac{n!}{m!(n-m)!} \left(\frac{n^2 - n}{n^2 - n} \right) = \frac{n!}{m!(n-m)!}$$

$$= {}^n C_m \quad \text{as required}$$

References

- [1] A. J. Sadler, D. W. S. Thorning: *Understanding Pure Mathematics*, Oxford University Press 1987.
- [2] Arthur Adam Freddy Goossens: Francis Lousberg. *Mathématisons 65*. DeBoeck, 3e edition 1991.
- [3] David Rayner, *Higher GCSE Mathematics*, Oxford University Press 2000
- [4] Frank Ebos, Dennis Hamaguchi, Barbana Morrison & John Klassen, *Mathematics Principles & Process*, Nelson Canada A Division of International Thomson Limited 1990
- [5] George B. Thomas, Maurice D. Weir & Joel R. Hass, *Thomas' Calculus Twelfth Edition*, Pearson Education, Inc. 2010
- [6] Geoff Mannall & Michael Kenwood, *Pure Mathematics 2*, Heinemann Educational Publishers 1995
- [7] H.K. DASS...*Engineering Mathematics*. New Delhi, S. CHAND&COMPANY LTD, thirteenth revised edition 2007.
- [8] Hubert Carnec, Genevieve Haye, Monique Nouet, ReneSeroux, Jacqueline Venard. *Mathématiques TS Enseignement obligatoire*. Bordas Paris 1994.
- [9] Jean Paul Beltramonde, VincentBrun, ClaudeFelloneau, LydiaMisset, Claude Talamoni. *Declic 1re S Mathématiques*. Hachette-education, Paris 2005.
- [10] JF Talber & HH Heing, *Additional Mathematics 6th Edition Pure & Applied*, Pearson Education South Asia Pte Ltd 1995
- [11] J.K. Backhouse, SPTHouldsworth B.E.D. Copper and P.J.F. Horril. *Pure Mathematics 2*. Longman, third edition 1985, fifteenth impression 1998.
- [12] M.Nelkon, P. Parker. *Advanced Level Pysics, Seventh Edition*. Heinemann 1995
- [13] Mukasonga Solange. *Mathématiques 12, Analyse Numérique*. KIE, Kigali 2006.
- [14] Ngezahayo Emmanuel, Icyingeneye Pacifique. *Advanced Mathematics for Rwanda Secondary Schools Learner's Book 4*, Fountain Publishers Ltd, 2016.
- [15] Ngezahayo Emmanuel, Icyingeneye Pacifique. *Subsidiary Mathematics for Rwanda Secondary Schools Learner's Book 4*, Fountain Publishers Ltd, 2016.
- [16] Ngezahayo Emmanuel, Icyingeneye Pacifique. *Advanced Mathematics for Rwanda Secondary Schools Learner's book Senior Five*, Fountain Publishers Ltd, 2017.

- [17] Ngezahayo Emmanuel, Icyingeneye Pacifique. *Subsidiary Mathematics for Rwanda Secondary Schools Learner's book Senior Five*, Fountain Publishers Ltd, 2017.
- [18] N. PISKOUNOV, *Calcul Différentiel et Integral tom II 9ème édition*. Editions MIR. Moscou, 1980.
- [19] Paule Faure- Benjamin Bouchon, *Mathématiques Terminales F*. Editions Nathan, Paris 1992.
- [20] Peter Smythe: *Mathematics HL & SL with HL options, Revised Edition*, Mathematics Publishing Pty. Limited, 2005.
- [21] Rwanda Education Board (2015), *Teacher training manual*, Ministry of Education: Rwanda.
- [22] Rwanda Education Board (2015), *Subsidiary Mathematics Syllabus S4-S6*, Ministry of Education: Rwanda.
- [23] Robert A. Adms & Christopher Essex, *Calculus A complete course* Seventh Edition, Pearson Canada Inc., Toronto, Ontario 2010
- [24] Seymour Lipschutz. *Schaum's outline of Theory and Problems of linear algebra*. McGraw-Hill 1968.
- [25] Shampiyona Aimable : *Mathématiques 6*. Kigali, Juin 2005.
- [26] Tom Duncan, *Advanced Physics, Fourth edition*. John Murray (Publishers) Ltd, London 1998
- [27] Yves Noirot, Jean–Paul Parisot, Nathalie Brouillet. *Cours de Physique Mathématiques pour la Physique*. Paris, DUNOD, 1997.

